

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 February 2006 (09.02.2006)

PCT

(10) International Publication Number
WO 2006/015389 A2

(51) International Patent Classification:
C12N 15/11 (2006.01)

(21) International Application Number:
PCT/US2005/029011

(22) International Filing Date: 4 August 2005 (04.08.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
10/912,440 4 August 2004 (04.08.2004) US

(71) Applicant (*for all designated States except US*): **UNIVERSITY OF MASSACHUSETTS** [US/US]; 225 Franklin Street, 12th Floor, Boston, MA 02110 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **ZAMORE, Phillip, D.** [US/US]; 500 Green Street, Northboro, MA 01532 (US). **HUTVAGNER, Gyorgy** [US/US]; 4 Browker Street #3, Worcester, MA 01604 (US). **SCHWARZ, Dianne** [US/US]; 72 Harley Drive, Apt. #3, Worcester, MA 01606 (US). **SIMARD, Martin** [CA/CA]; 150 Chemin du Roy, Deschambault, Quebec G0A 1S0 (CA).

(74) Agents: **MILASINCIC, Debra, J.** et al.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS AND COMPOSITIONS FOR ENHANCING THE EFFICACY AND SPECIFICITY OF RNA SILENCING

(57) Abstract: The present invention provides methods of enhancing the efficacy and specificity of RNA silencing. The invention also provides compositions for mediating RNA silencing. In particular, the invention provides siRNAs, siRNA-like molecules, shRNAs, vectors and transgenes having improved specificity and efficacy in mediating silencing of a target gene. Therapeutic methods are also featured.



WO 2006/015389 A2

METHODS AND COMPOSITIONS FOR ENHANCING THE EFFICACY AND SPECIFICITY OF RNA SILENCING

Government Rights

- 5 This invention was made at least in part with government support under grant nos. R01 GM62862-01, GM65236-01, and R21 NS44952-01 awarded by the National Institutes of Health. The government may have certain rights in this invention.

Related Applications

- 10 This application claims priority to U.S. Utility Application Serial No. 10/912,440, entitled "Methods and Compositions for Enhancing the Efficacy and Specificity of RNA Silencing" (filed August 4, 2004). The entire content of the above-referenced patent application is incorporated herein by this reference.

15 Background of the Invention

- Two types of ~21 nt RNAs trigger post-transcriptional gene silencing in animals: small interfering RNAs (siRNAs) and microRNAs (miRNAs). Both siRNAs and miRNAs are produced by the cleavage of double-stranded RNA (dsRNA) precursors by Dicer, a nuclease of the RNase III family of dsRNA-specific endonucleases (Bernstein et al., 2001; Billy et al., 2001; Grishok et al., 2001; Hutvagner et al., 2001; Ketting et al., 2001; Knight and Bass, 2001; Paddison et al., 2002; Park et al., 2002; Provost et al., 2002; Reinhart et al., 2002; Zhang et al., 2002; Doi et al., 2003; Myers et al., 2003). siRNAs result when transposons, viruses or endogenous genes express long dsRNA or when dsRNA is introduced experimentally into plant or animal cells to trigger gene silencing, a process known as RNA interference (RNAi) (Fire et al., 1998; Hamilton and Baulcombe, 1999; Zamore et al., 2000; Elbashir et al., 2001a; Hammond et al., 2001; Sijen et al., 2001; Catalanotto et al., 2002). In contrast, miRNAs are the products of endogenous, non-coding genes whose precursor RNA transcripts can form small stem-loops from which mature miRNAs are cleaved by Dicer (Lagos-Quintana et al., 2001; Lau et al., 2001; Lee and Ambros, 2001; Lagos-Quintana et al., 2002; Mourelatos et al., 2002; Reinhart et al., 2002; Ambros et al., 2003; Brennecke et al., 2003; Lagos-Quintana

et al., 2003; Lim et al., 2003a; Lim et al., 2003b). miRNAs are encoded by genes distinct from the mRNAs whose expression they control.

siRNAs were first identified as the specificity determinants of the RNA interference (RNAi) pathway (Hamilton and Baulcombe, 1999; Hammond et al., 2000),
5 where they act as guides to direct endonucleolytic cleavage of their target RNAs (Zamore et al., 2000; Elbashir et al., 2001a). Prototypical siRNA duplexes are 21 nt, double-stranded RNAs that contain 19 base pairs, with two-nucleotide, 3' overhanging ends (Elbashir et al., 2001a; Nykänen et al., 2001; Tang et al., 2003). Active siRNAs contain 5' phosphates and 3' hydroxyls (Zamore et al., 2000; Boutla et al., 2001;
10 Nykänen et al., 2001; Chiu and Rana, 2002). Similarly, miRNAs contain 5' phosphate and 3' hydroxyl groups, reflecting their production by Dicer (Hutvagner et al., 2001; Mallory et al., 2002).

In plants, miRNAs regulate the expression of developmentally important proteins, often by directing mRNA cleavage (Rhoades et al., 2002; Reinhart et al., 2002;
15 Llave et al., 2002a; Llave et al., 2002b; Xie et al., 2003; Kasschau et al., 2003; Tang et al., 2003; Chen, 2003). Whereas plant miRNA's show a high degree of complementarity to their mRNA targets, animal miRNA's have only limited complementarity to the mRNAs whose expression they control (Lee et al., 1993; Wightman et al., 1993; Olsen and Ambros, 1999; Reinhart et al., 2000; Slack et al., 2000; Abrahante et al., 2003;
20 Brennecke et al., 2003; Lin et al., 2003; Xu et al., 2003). Animal miRNAs are thought to repress mRNA translation, rather than promote target mRNA destruction (Lee et al., 1993; Wightman et al., 1993; Olsen and Ambros, 1999; Brennecke et al., 2003). Recent evidence suggests that the two classes of small RNAs are functionally interchangeable, with the choice of mRNA cleavage or translational repression
25 determined solely by the degree of complementarity between the small RNA and its target (Schwarz and Zamore, 2002; Hutvagner and Zamore, 2002; Zeng et al., 2003; Doench et al., 2003). Furthermore, siRNAs and miRNAs are found in similar, if not identical complexes, suggesting that a single, bifunctional complex –the RNA-induced silencing complex (RISC)- mediates both cleavage and translational control (Mourelatos
30 et al., 2002; Hutvagner and Zamore, 2002; Caudy et al., 2002; Martinez et al., 2002). Nonetheless, studies in both plants and animals show that at steady-state, siRNAs and miRNAs differ in at least one crucial respect: *in vivo* and *in vitro*, siRNAs are double-stranded, whereas miRNAs are single-stranded (Lee et al., 1993; Hamilton and

Baulcombe, 1999; Pasquinelli et al., 2000; Reinhart et al., 2000; Elbashir et al., 2001a; Djikeng et al., 2001; Nykänen et al., 2001; Lagos-Quintana et al., 2001; Lau et al., 2001; Lee and Ambros, 2001; Lagos-Quintana et al., 2002; Reinhart et al., 2002; Llave et al., 2002a; Silhavy et al., 2002; Llave et al., 2002b; Tang et al., 2003).

- 5 siRNA duplexes can assemble into RISC in the absence of target mRNA, both in vivo and in vitro (Tuschl et al., 1999; Hammond et al., 2000; Zamore et al., 2000). Each RISC contains only one of the two strands of the siRNA duplex (Martinez et al., 2002). Since siRNA duplexes have no foreknowledge of which siRNA strand will guide target cleavage, both strands must assemble with the appropriate proteins to form a RISC.
- 10 Previously, we and others showed that both siRNA strands are competent to direct RNAi (Tuschl et al., 1999; Hammond et al., 2000; Zamore et al., 2000; Elbashir et al., 2001b; Elbashir et al., 2001a; Nykänen et al., 2001). That is, the anti-sense strand of an siRNA can direct cleavage of a corresponding sense RNA target, whereas the sense siRNA strand directs cleavage of an anti-sense target. In this way, siRNA duplexes appear to be
- 15 functionally symmetric. The ability to control which strand of an siRNA duplex enters into the RISC complex to direct cleavage of a corresponding RNA target would provide a significant advance for both research and therapeutic applications of RNAi technology.

Summary of the Invention

- 20 A key step in RNA interference (RNAi) is the assembly of a catalytically active protein-RNA complex, the RNA-induced silencing complex (RISC), that mediates target RNA cleavage. The instant invention is based, at least in part, on the discovery that the two strands of a siRNA duplex do not contribute equally to RISC assembly. Rather, both the absolute and the relative stabilities of the base pairs at the 5' ends of the two siRNA
- 25 strands determines the degree to which each strand participates in the RNAi pathway. In fact, siRNA can be functionally asymmetric, with only one of the two strands able to trigger RNAi. The present invention is also based on the discovery that single stranded miRNAs are initially generated as siRNA-like duplexes whose structures predestine one strand to enter the RISC and the other strand to be destroyed. This finding helps to
- 30 explain the biogenesis of single-stranded miRNAs; the miRNA strand of a short-lived, siRNA duplex-like intermediate is assembled into a RISC complex, causing miRNAs to accumulate in vivo as single-stranded RNAs.

The present invention is further based on the discovery that RISC can cleave RNA targets with up to five contiguous mismatches at the siRNA 5' end and eight mismatches at the siRNA 3' end, indicating that 5' bases contribute disproportionately to target RNA binding, but do not play a role in determining the catalytic rate, *k_{cat}*. This finding explains how the 5', central and 3' sequences of the siRNA guide strand function to direct target cleavage.

The invention is further based on the discovery that the 3' bases of the siRNA contribute much less than 5' bases to the overall strength of binding, but instead help to establish the helical geometry required for RISC-mediated target cleavage, consistent with the view that catalysis by RISC requires a central A-form helix (Chiu et al., 2003). This finding indicates that complementarity is essential for translational repression by siRNAs designed to act like animal miRNAs, which typically repress translation (Doench et al., 2004).

The present invention is further based on the discovery that when an siRNA fails to pair with the first three, four or five nucleotides of the target RNA, the phosphodiester bond severed in the target RNA is unchanged; for perfectly matched siRNA, RISC measures the site of cleavage from the siRNA 5' end (Elbashir et al., 2001; Elbashir et al., 2001). This finding indicates that the identity of the scissile phosphate is determined prior to the encounter of the RISC with its target RNA, perhaps because the RISC endonuclease is positioned with respect to the siRNA 5' end during RISC assembly.

Accordingly, the instant invention features methods of enhancing the efficacy and specificity of RNAi. Also provided is a method of decreasing silencing of an inadvertent target by an RNAi agent. The invention further features compositions, including siRNAs, shRNAs, as well as vectors and transgenes, for mediating RNAi. The RNAi agents of the invention have improved specificity and efficacy in mediating silencing of a target gene.

The design rules for enhancing the efficacy and specificity of RNAi agents can also be applied to other RNA silencing agents, in particular agents that mediate translational repression of a target gene. Such silencing agents include, for example, siRNA-like duplexes that include a miRNA strand, as described *supra*. Methods featuring such silencing agents are also described herein.

Other features and advantages of the invention will be apparent from the following detailed description and claims.

Brief Description of the Drawings

- 5 *Figure 1.* RNAi mediated by asymmetric duplex and single-stranded siRNAs. (A) Schematic showing relevant portions of the sense and anti-sense target RNA sequences. (B) Schematic showing siRNA duplex sequence and graph depicting RNAi mediated by the antisense and sense strands. (C) Schematic showing siRNA sequences of individual single strands and graph depicting RNAi mediated by the single strands.
- 10 (D) Bar graph depicting fraction of total siRNA present as single-strand. (E) Schematic showing siRNA duplex sequence containing G:U wobble base pair and graph depicting RNAi mediated by the antisense and sense strands.

- Figure 2.* RNAi mediated by asymmetric duplex siRNAs. (A) Schematic showing relevant portions of the sense and anti-sense target RNA sequences. (B)
- 15 Schematic showing siRNA duplex sequence and graph depicting RNAi mediated by the antisense and sense strands. (C) Schematic showing siRNA duplex sequence containing A:U mismatch and graph depicting RNAi mediated by the antisense and sense strands. (D) Schematic showing siRNA duplex sequence containing G:U mismatch and graph depicting RNAi mediated by the antisense and sense strands. (E) Schematic showing
- 20 siRNA duplex sequence containing C:A mismatch and graph depicting RNAi mediated by the antisense and sense strands.

- Figure 3.* RNAi mediated by asymmetric duplex siRNAs. (A) Schematic showing relevant portions of the sense and anti-sense target RNA sequences. (B)
- 25 Schematic showing siRNA duplex sequence and graph depicting RNAi mediated by the antisense and sense strands. (C) Schematic showing siRNA duplex sequence containing A:G mismatch and graph depicting RNAi mediated by the antisense and sense strands. (D) Schematic showing siRNA duplex sequence containing C:U mismatch and graph depicting RNAi mediated by the antisense and sense strands. (E) Schematic showing siRNA duplex sequence containing A:U base pair and graph depicting RNAi mediated
- 30 by the antisense and sense strands. (F) Schematic showing siRNA duplex sequence containing A:G mismatch and graph depicting RNAi mediated by the antisense and

sense strands. (G) Schematic showing siRNA duplex sequence containing C:U mismatch and graph depicting RNAi mediated by the antisense and sense strands. (H) Schematic showing siRNA duplex sequence containing A:U base pair and graph depicting RNAi mediated by the antisense and sense strands. (I) Schematic of
 5 individual single-strands of siRNAs and graph depicting RNAi mediated by the individual single-strands.

Figure 4. RNAi mediated by asymmetric duplex siRNAs containing inosine. (A) Schematic showing siRNA duplex sequence having inosine at 5' end of sense strand and graph depicting RNAi mediated by the antisense and sense strands. (B) Schematic
 10 showing siRNA duplex sequence having inosine at 5' end of antisense strand and graph depicting RNAi mediated by the antisense and sense strands. (C) Schematic showing siRNA duplex sequence containing inosine in both strands and graph depicting RNAi mediated by the antisense and sense strands. (D) Schematic showing individual siRNA strands containing inosine and graph depicting RNAi mediated by the individual single-
 15 strands.

Figure 5. Symmetric cleavage of pre-*let-7* by Dicer. (A) Analysis of cleavage products produced on 5' side of precursor stem (*let-7*). (B) Analysis of cleavage products produced on 3' side of precursor stem (*let-7**). (C) Conceptual dicing of pre-*let-7* to a deduced pre-*let-7* siRNA.

20 *Figure 6.* Analysis of *Drosophila* miRNA genes for predicted miRNA and miRNA*. (A) Conceptual dicing of 26 published *Drosophila* miRNA genes to a deduced duplex siRNA. (B) Amounts of miR-10 and miR-10* detected *in vivo*.

Figure 7. Schematic representing mechanism of RISC assembly from pre-miRNA or dsRNA.

25 *Figure 8.* Reduction of off-target silencing by sense strand. (A) Sense and anti-sense *sod1* target RNA sequences. (B) Schematic showing siRNA duplex sequence and graph depicting RNAi mediated by the antisense and sense strands. (C) Schematic showing siRNA duplex sequence containing G:U wobble base pair and graph depicting RNAi mediated by the antisense and sense strands. (D) Schematic showing individual
 30 siRNA strands and graph depicting RNAi mediated by the individual single-strands. (E)

Thermodynamic analysis of siRNA strand 5' ends for the siRNA duplex in (B). ΔG (kcal/mole) was calculated in 1M NaCl at 37°C.

Figure 9. Enhancement of silencing by antisense strand. (A) Schematic showing relevant portions of the sense and anti-sense target RNA sequences. (B) Schematic showing siRNA duplex sequence and graph depicting RNAi mediated by the antisense and sense strands. (B) Schematic showing siRNA duplex sequence containing A:U base pair and graph depicting RNAi mediated by the antisense and sense strands. (C) Schematic showing siRNA duplex sequence containing A:G mismatch and graph depicting RNAi mediated by the antisense and sense strands. (D) Thermodynamic analysis of siRNA strand 5' ends for the siRNA duplex in (B). ΔG (kcal/mole) was calculated in 1M NaCl at 37°C.

Figure 10. The relative thermodynamic stability of the first four base pairs of the siRNA strands explains siRNA functional asymmetry. Thermodynamic analysis of siRNA strand 5' ends for the siRNAs in Figures 1B and 1E. ΔG (kcal/mole) was calculated in 1M NaCl at 37°C.

Figure 11. The first four base pairs of the siRNA duplex determine strand-specific activity. Internal, single-nucleotide mismatches (A-F) near the 5' ends of an siRNA strand generate functional asymmetry, but internal G:U wobble pairs (G-I) do not.

Figure 12. Increased rate of siRNA efficiency when duplexes have dTdT mismatched tails.

Figure 13. Product release limits the rate of catalysis by RISC. (a) ATP stimulates multiple rounds of RISC cleavage of the RNA target. siRNA was incubated with ATP in *Drosophila* embryo lysate, then NEM was added to quench RISC assembly and to disable the ATP-regenerating system. The energy regenerating system was either restored by adding additional creatine kinase (+ATP) or the reaction was ATP-depleted by adding hexokinase and glucose (-ATP). The target RNA concentration was 49 nM and the concentration of RISC was ~4 nM. The siRNA sequence and its target RNA is given in Figure 21A. (b) In the absence of ATP, cleavage by RISC produces a pre-steady state burst equal, within error, to the concentration of active RISC. The target concentration was 110 nM and the RISC concentration was ~4 nM. (c) Catalysis by

RISC is not enhanced by ATP under single-turnover conditions. RISC was present in ~8-fold excess over target. Each data point represents the average of two trials.

Figure 14. In the absence of ATP, mismatches between the 3' end of the siRNA guide strand and the target RNA facilitate product release, but reduce the rate of target cleavage. (a) Representative siRNA sequences are shown aligned with the target sequence. The siRNA guide strand is in color (5' to 3') and the mismatch with the target site is highlighted in yellow. A complete list of siRNA sequences and their corresponding target RNA sequences appear in Figure 21A. (b) The steady-state rate of cleavage in the presence and absence of ATP was determined for siRNAs with zero to four 3' mismatches with the target site. The target RNA concentration was 49 nM and the concentration of RISC was either ~4 nM (no mismatches) or ~6 nM (1 to 4 mismatches). The steady-state velocity with ATP, relative to the velocity without ATP is shown for each siRNA. (c) Time course of cleavage for perfectly matched (~16-fold excess of RISC relative to target) and mismatched (~80-fold excess of RISC) siRNA. (d) Data representative of those used in the analysis in (c) for target cleavage directed by siRNAs with zero, four, and five 3' mismatches.

Figure 15. Remarkable tolerance of RISC for 3' mismatches. (a) Each additional 3' mismatch further reduced the rate of cleavage by RISC. The steady-state rates of cleavage were determined for siRNA with zero, one, two, and four mismatches under multiple-turnover conditions (~49 nM target mRNA and ~4–6 nM RISC). (b) Analysis of siRNAs bearing zero to five 3' mismatches with the target RNA under conditions of slight enzyme excess (~2-fold more RISC than target). siRNA sequences and corresponding target RNA sequences used in (a) and (b) are shown in Figure 14A and Figure 21A. (c) Extended endpoint analysis of RISC cleavage under conditions of ~80-fold enzyme excess reveals that cleavage can occur for siRNAs with as many as eight mismatches to the target RNA. Note the different time scales in (c) versus (b). All reactions were under standard in vitro RNAi (+ATP) conditions. siRNA sequences and corresponding target RNA sequences used in (c) are shown in Figure 21B.

Figure 16. Limited tolerance of RISC for 5' mismatches. (a) RISC cleavage was analyzed as in Figure 21C using 5' mismatched siRNAs, whose sequences are given in Figure 21. The target RNA was the same for all siRNAs. siRNA sequences and the corresponding target RNA sequence are shown in Figure 21E. (b) RISC cleavage was

analyzed using a single siRNA sequence. Mismatches were created by altering the sequence of the target RNA. For the target containing compensatory mutations, the target concentration was 0.25 nM and the siRNA concentration was ~20 nM; RISC concentration was not determined. The asterisk denotes a 15 second time-point. siRNA sequences and corresponding target RNA sequence are shown in Figure 21F. (c) RISC cleavage was analyzed by incubating 50 nM siRNA with 0.5 nM target RNA. 3' mismatches were created by modifying the target sequence, and 5' mismatches by changing the siRNA. Target and siRNA sequences are given in Figure 21C. (d) Perfectly base-paired and 5' mismatched siRNAs direct cleavage at the same phosphodiester bond. Cleavage reactions were performed with ~20 nM RISC generated from 50 nM siRNA and 0.5 nM target RNA and analyzed on an 8% denaturing polyacrylamide sequencing gel. The target mRNA was 182 nt and 5' cleavage product was 148 nt. After RISC was assembled, the extract was treated with NEM to inactivate nucleases (Schwartz et al., 2004). After NEM treatment, the ATP regenerating system was restored by adding additional creatine kinase, then target RNA was added and the incubation continued for the indicated time. OH⁻ denotes a base hydrolysis ladder. The siRNA and corresponding target RNA sequences used in this experiment are shown in Figure 21E.

Figure 17. Michaelis-Menten and K_i analysis for matched and mismatched siRNAs reveal distinct contributions to binding and catalysis for the 5', central, and 3' regions of the siRNA. (a) siRNA was assembled into RISC under standard in vitro RNAi conditions, then diluted to achieve the desired RISC concentration. The initial rates of cleavage were determined for increasing concentrations of 5' ³²P-cap-radiolabeled target mRNA. Plot of initial velocity versus substrate concentration. K_M and V_{max} were determined by fitting the data to the Michaelis-Menten equation. See Table 1 for analysis. Representative initial rate determinations appear in Figure 20A, and the siRNA sequences used in this experiment are shown in Figures 21A and 21E. (b) K_i values were determined in competition assays using 2'-O-methyl oligonucleotides bearing 5', central, and 3' mismatches to the siRNA guide strand. Representative data are presented in Figure 20B, and a complete list of the 2'-O-methyl oligonucleotides used appears in Figure 21D.

Figure 18. A model for the cycle of RISC assembly, target recognition, catalysis, and recycling.

Figure 19. Exogenously programmed RISC is a bona fide enzyme siRNA was assembled into RISC for 1 hour in a standard in vitro RNAi reaction, then assembly was quenched with N-ethyl maleimide (NEM)21,29. The amount of RISC formed was determined by measuring 32P-radiolabeled siRNA retained on a tethered 5'-biotinylated, 31-nt, 2'-O-methyl oligonucleotide complementary to the guide strand of the siRNA. RISC binds essentially irreversibly to tethered 2'-O-methyl oligonucleotides, but cannot cleave these RNA-analogs (Hutvagner et al., 2004; Schwartz et al., 2003). In all experiments, target-cleaving activity was not detected in the supernatant, demonstrating that all the active RISC was retained on the beads. (a) Sequence of the siRNA used (guide strand in red, 32P-radiolabel marked with an asterisk). *Drosophila* let-7 is not expressed in 0–2 hour embryos (Hutvagner et al., 2001), so the only source of let-7 in the in vitro reactions was the exogenous let-7 siRNA. The 5' end of the guide strand of the let-7 siRNA is predicted to be thermodynamically more stable than the 5' end of the passenger strand, explaining why only a low concentrations of let-7-programmed RISC is formed (Schwartz et al., 2003, Khvorova et al., 2003). The maximum amount of RISC assembled varies widely with siRNA sequence. The siRNAs used in Figures 3–8 were designed to load =5-fold more guide strand-containing RISC (Hutvagner et al., 2001; Schwartz et al., 2003) (b) Representative gel confirming that the RISC was removed by the tethered 2'-O-methyl oligonucleotide. A reaction prior to incubation with the tethered 2'-O-methyl oligonucleotide (pre) was compared to the supernatant of a reaction incubated with beads alone (mock), and the supernatant of a reaction incubated with the complementary tethered 2'-O-methyl oligonucleotide (post). The buffer reaction contained no siRNA. (c) Analysis of the amount of RISC assembled at various siRNA concentrations. 5' 32P-radiolabeled siRNA was incubated with lysate for 1 hour, then reactions were quenched by treatment with NEM, and RISC concentration was measured using the tethered 2'-O-methyl oligonucleotide method.

Figure 20. Michaelis-Menton and Competitor Analysis of RISC (a) Representative data for the determination of initial velocities for the perfectly matched siRNA. Black, 1 nM target; red, 5 nM; blue, 20 nM; and green, 60 nM. (b) Three independent experiments for inhibition by a fully complementary 2'-O-methyl oligonucleotide competitor. ~1 nM RISC and 5 nM 32P-cap-radiolabeled target mRNA

were incubated with increasing concentration of competitor, and the initial velocities were calculated and plotted versus competitor concentration.

Figure 21. siRNAs, target sites, and 2'-O-methyl oligonucleotides used in this study. (a) List of siRNA and corresponding target RNA sequences used in the experiments described in Table 1 and Figures 14B-D, 15A, 15B, 17A, and 20A. (b) List of siRNA and corresponding target RNA sequences used in the experiment described in Figure 15C. (c) List of siRNA and corresponding target RNA sequences used in the experiment described in Figure 16C. (d) List of siRNAs and corresponding target 2'-O-methyl oligonucleotide sequences used in the experiments described in Figures 17B and 20B. (e) List of siRNA and corresponding target RNA sequences used in the experiments described in Figures 16A, 16D, 17A, and Table 1. (f) List of siRNA and corresponding target RNA sequences used in the experiment described in Figure 16B. .

Detailed Description of the Invention

A key step in RNA interference (RNAi) is the assembly of a catalytically active protein-RNA complex, the RNA-induced silencing complex (RISC), that mediates target RNA cleavage. Each RISC contains one of the two strands of the small interfering RNA (siRNA) duplex that triggers RNAi. The instant invention is based, at least in part, on the discovery that the two siRNA strands do not contribute equally to RISC assembly. Small changes in siRNA sequence were found to have profound and predictable effects on the extent to which the two strands of an siRNA duplex enter the RNAi pathway, a phenomenon termed siRNA functional "asymmetry". The discoveries described herein reveal that the strength of the base-pairing interactions made by the 5' end of each siRNA strand with the 3' region of strand to which it is paired determines which of the two strands participates in the RNAi pathway. RISC assembly appears to be governed by an enzyme that initiates unwinding of an siRNA duplex at the siRNA strand whose 5' end is less tightly paired to the complementary siRNA strand.

Remarkably, such highly asymmetric siRNA duplexes resemble proposed intermediates in the biogenesis pathway of microRNA (miRNA) (Hutvagner and Zamore, 2002; Reinhart et al., 2002; Lim et al., 2003b). miRNAs are endogenous, ~21-nt single-stranded RNAs processed by Dicer from stem-loop RNA precursors that

regulate gene expression in animals and plants. A striking feature of miRNA precursors is their lack of full complementarity in the stem region. The discoveries presented herein indicate an important role for the discontinuities in the stem region of miRNAs; it is likely that miRNAs are initially generated from their precursor RNAs as siRNA-like
5 duplexes, and that the structure of these duplexes predestines the miRNA strand to enter the RISC and the other strand to be destroyed. Thus, nature appears to have optimized the stem portion of miRNAs to follow a set of rules dictating which strand enters the RISC complex.

The discoveries made by the instant inventors provide rules according to which
10 siRNAs and shRNAs can be designed that are fully asymmetric, with only one of the two siRNA strands competent to enter the RISC complex. By applying these rules to the selection and design of a targeted RNAi agent, e.g., siRNAs and shRNAs, the antisense strand of the RNAi agent can be predictably directed to enter the RISC complex and mediate target RNA cleavage. Similarly, the sense strand can be discouraged from
15 entering the RISC complex, thereby reducing or eliminating undesired silencing of an inadvertent target by the sense strand.

Accordingly, the instant invention provides methods for improving the efficiency (or specificity) of an RNAi reaction comprising identifying an off target RNAi activity mediated by the sense strand of an RNAi agent, and modifying the RNAi agent such that
20 the base pair strength between the 5' end of the antisense strand and the 3' end of the sense strand is lessened relative to the base pair strength of the 5' end of the sense strand and the 3' end of the antisense strand (e.g., relative to the premodified RNAi agent), such that the sense strand is less effective at entering RISC (e.g., less effective than the premodified RNAi agent).

25 The instant invention also provides methods for improving the efficiency (or specificity) of an RNAi reaction comprising modifying (e.g., increasing) the asymmetry of the RNAi agent such that the ability of the sense or second strand to mediate RNAi (e.g., mediate cleavage of target RNA) is lessened. In preferred embodiments, the asymmetry is increased in favor of the 5' end of the first strand, e.g., lessening the bond
30 strength (e.g., the strength of the interaction) between the 5' end of the first strand and 3' end of the second strand relative to the bond strength (e.g., the strength of the interaction) between the 5' end of the second strand and the 3' end of the first strand. .

In other embodiments, the asymmetry is increased in favor of the 5' end of the first strand by increasing bond strength (*e.g.*, the strength of the interaction) between the 5' end of the second or sense strand and the 3' end of the first or antisense strand, relative to the bond strength (*e.g.*, the strength of the interaction) between the 5' end of the first and the 3' end of the second strand. In embodiments of the invention, the bond strength is increased, *e.g.*, the H bonding is increased between nucleotides or analogs at the 5' end, *e.g.*, within 5 nucleotides of the second or sense strand (numbered from the 5' end of the second strand) and complementary nucleotides of the first or antisense strand. It is understood that the asymmetry can be zero (*i.e.*, no asymmetry), for example, when the bonds or base pairs between the 5' and 3' terminal bases are of the same nature, strength or structure. More routinely, however, there exists some asymmetry due to the different nature, strength or structure of at least one nucleotide (often one or more nucleotides) between terminal nucleotides or nucleotide analogs.

Accordingly, in one aspect, the instant invention provides a method of enhancing the ability of a first strand of a RNAi agent to act as a guide strand in mediating RNAi, involving lessening the base pair strength between the 5' end of the first strand and the 3' end of a second strand of the duplex as compared to the base pair strength between the 3' end of the first strand and the 5' end of the second strand.

In a related aspect, the invention provides a method of enhancing the efficacy of a siRNA duplex, the siRNA duplex comprising a sense and an antisense strand, involving lessening the base pair strength between the antisense strand 5' end (AS 5') and the sense strand 3' end (S 3') as compared to the base pair strength between the antisense strand 3' end (AS 3') and the sense strand 5' end (S 5'), such that efficacy is enhanced.

In another aspect of the invention, a method is provided for promoting entry of a desired strand of an siRNA duplex into a RISC complex, comprising enhancing the asymmetry of the siRNA duplex, such that entry of the desired strand is promoted. In one embodiment of this aspect of the invention, the asymmetry is enhanced by lessening the base pair strength between the 5' end of the desired strand and the 3' end of a complementary strand of the duplex as compared to the base pair strength between the 3' end of the desired strand and the 5' end of the complementary strand.

In another aspect of the invention, a siRNA duplex is provided comprising a sense strand and an antisense strand, wherein the base pair strength between the antisense strand 5' end (AS 5') and the sense strand 3' end (S 3') is less than the base pair strength between the antisense strand 3' end (AS 3') and the sense strand 5' end (S 5'), such that the antisense strand preferentially guides cleavage of a target mRNA.

In one embodiment of these aspects of the invention, the base-pair strength is less due to fewer G:C base pairs between the 5' end of the first or antisense strand and the 3' end of the second or sense strand than between the 3' end of the first or antisense strand and the 5' end of the second or sense strand.

10 In another embodiment, the base pair strength is less due to at least one mismatched base pair between the 5' end of the first or antisense strand and the 3' end of the second or sense strand. Preferably, the mismatched base pair is selected from the group consisting of G:A, C:A, C:U, G:G, A:A, C:C and U:U.

In one embodiment, the base pair strength is less due to at least one wobble base pair, e.g., G:U, between the 5' end of the first or antisense strand and the 3' end of the second or sense strand.

In another embodiment, the base pair strength is less due to at least one base pair comprising a rare nucleotide, e.g., inosine (I). Preferably, the base pair is selected from the group consisting of an I:A, I:U and I:C.

20 In yet another embodiment, the base pair strength is less due to at least one base pair comprising a modified nucleotide. In preferred embodiments, the modified nucleotide is selected from the group consisting of 2-amino-G, 2-amino-A, 2,6-diamino-G, and 2,6-diamino-A.

In several embodiments of these aspects of the invention, the RNAi agent is a siRNA duplex or is derived from an engineered precursor, and can be chemically synthesized or enzymatically synthesized.

In another aspect of the instant invention, compositions are provided comprising a siRNA duplex of the invention formulated to facilitate entry of the siRNA duplex into a cell. Also provided are pharmaceutical composition comprising a siRNA duplex of the invention.

30 Further provided are an engineered pre-miRNA comprising a siRNA duplex as described above, as well as a vector encoding the pre-miRNA. In related aspects, the

invention provides a pri-miRNA comprising the pre-miRNA, as well as a vector encoding the pri-miRNA.

Also featured in the instant invention are small hairpin RNA (shRNA) comprising nucleotide sequence identical to the sense and antisense strand of the siRNA
5 duplexes as described above. In one embodiment, the nucleotide sequence identical to the sense strand is upstream of the nucleotide sequence identical to the antisense strand. In another embodiment, the nucleotide sequence identical to the antisense strand is upstream of the nucleotide sequence identical to the sense strand. Further provided are vectors and transgenes encoding the shRNAs of the invention.

10 In yet another aspect, the invention provides cells comprising the vectors featured in the instant invention. Preferably, the cell is a mammalian cell, e.g., a human cell.

In other aspects of the invention, methods of enhancing silencing of a target mRNA, comprising contacting a cell having an RNAi pathway with a RNAi agent as
15 described above under conditions such that silencing is enhanced.

Also provided are methods of enhancing silencing of a target mRNA in a subject, comprising administering to the subject a pharmaceutical composition comprising a RNAi agent as described above such that silencing is enhanced.

Further provided is a method of decreasing silencing of an inadvertant target
20 mRNA by a dsRNAi agent, the dsRNAi agent comprising a sense strand and an antisense strand involving the steps of: (a) detecting a significant degree of complementarity between the sense strand and the inadvertant target; and (b) enhancing the base pair strength between the 5' end of the sense strand and the 3' end of the antisense strand relative to the base pair strength between the 3' end of the sense strand
25 and the 5' end of the antisense strand; such that silencing of the inadvertant target mRNA is decreased. In a preferred embodiment, the silencing of the inadvertant target mRNA is decreased relative to silencing of a desired target mRNA.

The design rules described *supra* can also be applied to other classes of RNA silencing agents, for example, siRNA-like duplexes that include a miRNA strand. In
30 particular, the design rules can be applied to such siRNA-like duplexes to promote entry of the desired strand of the duplex (e.g., the miRNA strand) into a RISC complex. The degree of complementarity between the miRNA strand and its target determines whether the small RNA mediates mRNA cleavage or translational repression, as described *supra*.

Such duplexes are useful for effectively delivering miRNAs, for example, to cells or animals for therapeutic purposes.

Accordingly, in one aspect, the instant invention provides a method of enhancing the ability of a first strand of a siRNA-like duplex to act in mediating RNA silencing (e.g., RNAi or translational repression), involving lessening the base pair strength between the 5' end of the first strand and the 3' end of a second strand of the duplex as compared to the base pair strength between the 3' end of the first strand and the 5' end of the second strand.

In a related aspect, the invention provides a method of enhancing the efficacy of a siRNA-like duplex, the siRNA-like duplex comprising a miRNA and an miRNA* strand, involving lessening the base pair strength between the miRNA strand 5' end (miRNA 5') and the miRNA* strand 3' end (miRNA* 3') as compared to the base pair strength between the miRNA strand 3' end (miRNA 3') and the miRNA* strand 5' end (miRNA* 5'), such that efficacy is enhanced.

In another aspect of the invention, a method is provided for promoting entry of a desired strand of an siRNA-like duplex into a RISC complex, comprising enhancing the asymmetry of the siRNA duplex, such that entry of the desired strand is promoted. In one embodiment of this aspect of the invention, the asymmetry is enhanced by lessening the base pair strength between the 5' end of the desired strand and the 3' end of a complementary strand of the duplex as compared to the base pair strength between the 3' end of the desired strand and the 5' end of the complementary strand.

In another aspect of the invention, a siRNA-like duplex is provided comprising a miRNA strand and an miRNA* strand, wherein the base pair strength between the miRNA strand 5' end (miRNA 5') and the miRNA* strand 3' end (miRNA* 3') is less than the base pair strength between the miRNA strand 3' end (miRNA 3') and the miRNA* strand 5' end (miRNA* 5'), such that the miRNA strand preferentially silences (e.g., cleaves or represses) a target mRNA.

In one embodiment of these aspects of the invention, the base-pair strength is less due to fewer G:C base pairs between the 5' end of the first or miRNA strand and the 3' end of the second or miRNA* strand than between the 3' end of the first or miRNA strand and the 5' end of the second or miRNA* strand.

In another embodiment, the base pair strength is less due to at least one mismatched base pair between the 5' end of the first or miRNA strand and the 3' end of

the second or miRNA* strand. Preferably, the mismatched base pair is selected from the group consisting of G:A, C:A, C:U, G:G, A:A, C:C and U:U.

In one embodiment, the base pair strength is less due to at least one wobble base pair, e.g., G:U, between the 5' end of the first or miRNA strand and the 3' end of the
5 second or miRNA* strand.

In another embodiment, the base pair strength is less due to at least one base pair comprising a rare nucleotide, e.g., inosine (I). Preferably, the base pair is selected from the group consisting of an I:A, I:U and I:C.

In yet another embodiment, the base pair strength is less due to at least one base
10 pair comprising a modified nucleotide. In preferred embodiments, the modified nucleotide is selected from the group consisting of 2-amino-G, 2-amino-A, 2,6-diamino-G, and 2,6-diamino-A.

In several embodiments of these aspects of the invention, the RNA silencing agent is a siRNA-like duplex or is derived from an engineered precursor, and can be
15 chemically synthesized or enzymatically synthesized.

In another aspect of the instant invention, compositions are provided comprising a siRNA-like duplex of the invention formulated to facilitate entry of the siRNA-like duplex into a cell. Also provided are pharmaceutical composition comprising a siRNA-like duplex of the invention.

20 Further provided are an engineered pre-miRNA comprising the siRNA-like duplex as described above, as well as a vector encoding the pre-miRNA. In related aspects, the invention provides a pri-miRNA comprising the pre-miRNA, as well as a vector encoding the pri-miRNA.

Also featured in the instant invention are small hairpin RNA (shRNA)
25 comprising nucleotide sequence identical to the miRNA and miRNA* strand of the siRNA-like duplex as described above. In one embodiment, the nucleotide sequence identical to the miRNA strand is upstream of the nucleotide sequence identical to the miRNA* strand. In another embodiment, the nucleotide sequence identical to the miRNA* strand is upstream of the nucleotide sequence identical to the miRNA strand.

30 Further provided are vectors and transgenes encoding the shRNAs of the invention.

In yet another aspect, the invention provides cells comprising the vectors featured in the instant invention. Preferably, the cell is a mammalian cell, e.g., a human cell.

In other aspects of the invention, methods of enhancing silencing of a target mRNA, comprising contacting a cell having an RNA silencing pathway with the RNA silencing agent as described above under conditions such that silencing is enhanced.

Also provided are methods of enhancing silencing of a target mRNA in a subject,
5 comprising administering to the subject a pharmaceutical composition comprising the RNA silencing agent as described above such that silencing is enhanced.

Further provided is a method of decreasing silencing of an inadvertant target mRNA by a dsRNA silencing agent, the dsRNA silencing agent (*e.g.*, a siRNA-like duplex) comprising a miRNA strand and a miRNA* strand involving the steps of: (a)
10 detecting a significant degree of complementarity between the miRNA* strand and the inadvertant target; and (b) enhancing the base pair strength between the 5' end of the miRNA* strand and the 3' end of the miRNA strand relative to the base pair strength between the 3' end of the miRNA* strand and the 5' end of the miRNA strand; such that silencing of the inadvertant target mRNA is decreased. In a preferred embodiment, the
15 silencing of the inadvertant target mRNA is decreased relative to silencing of a desired target mRNA.

So that the invention may be more readily understood, certain terms are first defined.

20 The term "nucleoside" refers to a molecule having a purine or pyrimidine base covalently linked to a ribose or deoxyribose sugar. Exemplary nucleosides include adenosine, guanosine, cytidine, uridine and thymidine. The term "nucleotide" refers to a nucleoside having one or more phosphate groups joined in ester linkages to the sugar moiety. Exemplary nucleotides include nucleoside monophosphates, diphosphates and
25 triphosphates. The terms "polynucleotide" and "nucleic acid molecule" are used interchangeably herein and refer to a polymer of nucleotides joined together by a phosphodiester linkage between 5' and 3' carbon atoms.

The term "RNA" or "RNA molecule" or "ribonucleic acid molecule" refers to a polymer of ribonucleotides. The term "DNA" or "DNA molecule" or deoxyribonucleic
30 acid molecule" refers to a polymer of deoxyribonucleotides. DNA and RNA can be synthesized naturally (*e.g.*, by DNA replication or transcription of DNA, respectively). RNA can be post-transcriptionally modified. DNA and RNA can also be chemically

synthesized. DNA and RNA can be single-stranded (*i.e.*, ssRNA and ssDNA, respectively) or multi-stranded (*e.g.*, double stranded, *i.e.*, dsRNA and dsDNA, respectively). “mRNA” or “messenger RNA” is single-stranded RNA that specifies the amino acid sequence of one or more polypeptide chains. This information is translated
5 during protein synthesis when ribosomes bind to the mRNA.

As used herein, the term “RNA silencing agent” refers to an RNA which is capable of preventing complete processing (*e.g.*, the full translation and/or expression) of a mRNA molecule through a post-transcriptional silencing mechanism. RNA silencing agents include small (<50 b.p.), noncoding RNA molecules, for example RNA duplexes
10 comprising paired strands, as well as precursor RNAs from which such small non-coding RNAs can be generated. Exemplary RNA silencing agents include siRNAs, miRNAs, and siRNA-like duplexes, as well as precursors thereof.

As used herein, the term “small interfering RNA” (“siRNA”) (also referred to in the art as “short interfering RNAs”) refers to an RNA (or RNA analog) comprising
15 between about 10-50 nucleotides (or nucleotide analogs) which is capable of directing or mediating RNA interference. Preferably, an siRNA comprises between about 15-30 nucleotides or nucleotide analogs, more preferably between about 16-25 nucleotides (or nucleotide analogs), even more preferably between about 18-23 nucleotides (or nucleotide analogs), and even more preferably between about 19-22 nucleotides (or
20 nucleotide analogs) (*e.g.*, 19, 20, 21 or 22 nucleotides or nucleotide analogs).

As used herein, the term “microRNA” (“miRNA”), also referred to in the art as “small temporal RNAs” (“stRNAs”), refers to a small (10-50 nucleotide) RNA which is capable of directed or mediating RNA silencing. A “natural miRNA” refers to a microRNA that occurs naturally. An “miRNA disorder” shall refer to a disease or
25 disorder characterized by a aberrant expression or activity of a natural miRNA.

As used herein, the term “rare nucleotide” refers to a naturally occurring nucleotide that occurs infrequently, including naturally occurring deoxyribonucleotides or ribonucleotides that occur infrequently, *e.g.*, a naturally occurring ribonucleotide that is not guanosine, adenosine, cytosine, or uridine. Examples of rare nucleotides include,
30 but are not limited to, inosine, 1-methyl inosine, pseudouridine, 5,6-dihydrouridine, ribothymidine, ²N-methylguanosine and ^{2,2}N,N-dimethylguanosine.

The term “nucleotide analog” or “altered nucleotide” or “modified nucleotide” refers to a non-standard nucleotide, including non-naturally occurring ribonucleotides or deoxyribonucleotides. Preferred nucleotide analogs are modified at any position so as to alter certain chemical properties of the nucleotide yet retain the ability of the nucleotide analog to perform its intended function. Examples of preferred modified nucleotides include, but are not limited to, 2-amino-guanosine, 2-amino-adenosine, 2,6-diamino-guanosine and 2,6-diamino-adenosine. Examples of positions of the nucleotide which may be derivitized include the 5 position, *e.g.*, 5-(2-amino)propyl uridine, 5-bromo uridine, 5-propyne uridine, 5-propenyl uridine, etc.; the 6 position, *e.g.*, 6-(2-amino)propyl uridine; the 8-position for adenosine and/or guanosines, *e.g.*, 8-bromo guanosine, 8-chloro guanosine, 8-fluoroguanosine, etc. Nucleotide analogs also include deaza nucleotides, *e.g.*, 7-deaza-adenosine; O- and N-modified (*e.g.*, alkylated, *e.g.*, N6-methyl adenosine, or as otherwise known in the art) nucleotides; and other heterocyclically modified nucleotide analogs such as those described in Herdewijn, *Antisense Nucleic Acid Drug Dev.*, 2000 Aug. 10(4):297-310.

Nucleotide analogs may also comprise modifications to the sugar portion of the nucleotides. For example the 2' OH-group may be replaced by a group selected from H, OR, R, F, Cl, Br, I, SH, SR, NH₂, NHR, NR₂, COOR, or OR, wherein R is substituted or unsubstituted C₁–C₆ alkyl, alkenyl, alkynyl, aryl, etc. Other possible modifications include those described in U.S. Patent Nos. 5,858,988, and 6,291,438.

The phosphate group of the nucleotide may also be modified, *e.g.*, by substituting one or more of the oxygens of the phosphate group with sulfur (*e.g.*, phosphorothioates), or by making other substitutions which allow the nucleotide to perform its intended function such as described in, for example, Eckstein, *Antisense Nucleic Acid Drug Dev.* 2000 Apr. 10(2):117-21, Rusckowski *et al.* *Antisense Nucleic Acid Drug Dev.* 2000 Oct. 10(5):333-45, Stein, *Antisense Nucleic Acid Drug Dev.* 2001 Oct. 11(5): 317-25, Vorobjev *et al.* *Antisense Nucleic Acid Drug Dev.* 2001 Apr. 11(2):77-85, and U.S. Patent No. 5,684,143. Certain of the above-referenced modifications (*e.g.*, phosphate group modifications) preferably decrease the rate of hydrolysis of, for example, polynucleotides comprising said analogs *in vivo* or *in vitro*.

The term “oligonucleotide” refers to a short polymer of nucleotides and/or nucleotide analogs. The term “RNA analog” refers to an polynucleotide (*e.g.*, a

chemically synthesized polynucleotide) having at least one altered or modified nucleotide as compared to a corresponding unaltered or unmodified RNA but retaining the same or similar nature or function as the corresponding unaltered or unmodified RNA. As discussed above, the oligonucleotides may be linked with linkages which
5 result in a lower rate of hydrolysis of the RNA analog as compared to an RNA molecule with phosphodiester linkages. For example, the nucleotides of the analog may comprise methylenediol, ethylene diol, oxymethylthio, oxyethylthio, oxycarbonyloxy, phosphorodiamidate, phosphoroamidate, and/or phosphorothioate linkages. Preferred RNA analogues include sugar- and/or backbone-modified ribonucleotides and/or
10 deoxyribonucleotides. Such alterations or modifications can further include addition of non-nucleotide material, such as to the end(s) of the RNA or internally (at one or more nucleotides of the RNA). An RNA analog need only be sufficiently similar to natural RNA that it has the ability to mediate (mediates) RNA interference.

As used herein, the term “RNA interference” (“RNAi”) (also referred to in the
15 art as “gene silencing” and/or “target silencing”, *e.g.*, “target mRNA silencing”) refers to a selective intracellular degradation of RNA. RNAi occurs in cells naturally to remove foreign RNAs (*e.g.*, viral RNAs). Natural RNAi proceeds *via* fragments cleaved from free dsRNA which direct the degradative mechanism to other similar RNA sequences. As used herein, the term “translational repression” refers to a selective inhibition of
20 mRNA translation. Natural translational repression proceeds *via* miRNAs cleaved from shRNA precursors. Both RNAi and translational repression are mediated by RISC. Both RNAi and translational repression occur naturally or can be initiated by the hand of man, for example, to silence the expression of target genes.

As used herein, the term “antisense strand” of an siRNA or RNAi agent refers to
25 a strand that is substantially complementary to a section of about 10-50 nucleotides, *e.g.*, about 15-30, 16-25, 18-23 or 19-22 nucleotides of the mRNA of the gene targeted for silencing. The antisense strand or first strand has sequence sufficiently complementary to the desired target mRNA sequence to direct target-specific RNA interference (RNAi), *e.g.*, complementarity sufficient to trigger the destruction of the desired target mRNA by
30 the RNAi machinery or process (RNAi interference) or complementarity sufficient to trigger translational repression of the desired target mRNA. The term “sense strand” or “second strand” of an siRNA or RNAi agent refers to a strand that is complementary to

the antisense strand or first strand. Antisense and sense strands can also be referred to as first or second strands, the first or second strand having complementarity to the target sequence and the respective second or first strand having complementarity to said first or second strand. miRNA duplex intermediates or siRNA-like duplexes include a miRNA strand having sufficient complementarity to a section of about 10-50 nucleotides of the mRNA of the gene targeted for silencing and a miRNA* strand having sufficient complementarity to form a duplex with the miRNA strand.

As used herein, the term "guide strand" refers to a strand of an RNAi agent, e.g., an antisense strand of an siRNA duplex, that enters into the RISC complex and directs cleavage of the target mRNA.

A "target gene" is a gene whose expression is to be selectively inhibited or "silenced." This silencing can be achieved by cleaving the mRNA of the target gene or *via* translational repression of the target gene.

The term "engineered," as in an engineered RNA precursor, or an engineered nucleic acid molecule, indicates that the precursor or molecule is not found in nature, in that all or a portion of the nucleic acid sequence of the precursor or molecule is created or selected by man. Once created or selected, the sequence can be replicated, translated, transcribed, or otherwise processed by mechanisms within a cell. Thus, an RNA precursor produced within a cell from a transgene that includes an engineered nucleic acid molecule is an engineered RNA precursor.

As used herein, the term "asymmetry", as in the asymmetry of a RNA duplex (*e.g.* a siRNA duplex or siRNA-like or miRNA duplex), refers to an inequality of bond strength or base pairing strength between the duplex termini (*e.g.*, between terminal nucleotides on a first strand and terminal nucleotides on an opposing second strand), such that the 5' end of one strand of the duplex is more frequently in a transient unpaired, *e.g.*, single-stranded, state than the 5' end of the complementary strand. This structural difference determines that one strand of the duplex is preferentially incorporated into a RISC complex. The strand whose 5' end is less tightly paired to the complementary strand will preferentially be incorporated into RISC and mediate gene silencing (*e.g.* RNAi or translational repression).

As used herein, the term “bond strength” or “base pair strength” refers to the strength of the interaction between pairs of nucleotides (or nucleotide analogs) on opposing strands of an oligonucleotide duplex (e.g., an siRNA duplex), due primarily to H-bonding, Van der Waals interactions, and the like between said nucleotides (or
5 nucleotide analogs).

As used herein, the “5’ end”, as in the 5’ end of an antisense strand, refers to the 5’ terminal nucleotides, e.g., between one and about 5 nucleotides at the 5’ terminus of the antisense strand. As used herein, the “3’ end”, as in the 3’ end of a sense strand, refers to the region, e.g., a region of between one and about 5 nucleotides, that is
10 complementary to the nucleotides of the 5’ end of the complementary antisense strand.

As used herein, the term “isolated RNA” (e.g., “isolated shRNA”, “isolated siRNA”, “isolated siRNA-like duplex”, “isolated miRNA”, “isolated gene silencing agent”, or “isolated RNAi agent”) refers to RNA molecules which are substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically
15 synthesized.

As used herein, the term “transgene” refers to any nucleic acid molecule, which is inserted by artifice into a cell, and becomes part of the genome of the organism that develops from the cell. Such a transgene may include a gene that is partly or entirely
20 heterologous (i.e., foreign) to the transgenic organism, or may represent a gene homologous to an endogenous gene of the organism. The term “transgene” also means a nucleic acid molecule that includes one or more selected nucleic acid sequences, e.g., DNAs, that encode one or more engineered RNA precursors, to be expressed in a transgenic organism, e.g., animal, which is partly or entirely heterologous, i.e., foreign,
25 to the transgenic animal, or homologous to an endogenous gene of the transgenic animal, but which is designed to be inserted into the animal's genome at a location which differs from that of the natural gene. A transgene includes one or more promoters and any other DNA, such as introns, necessary for expression of the selected nucleic acid sequence, all operably linked to the selected sequence, and may include an enhancer sequence.

30 The term “*in vitro*” has its art recognized meaning, e.g., involving purified reagents or extracts, e.g., cell extracts. The term “*in vivo*” also has its art recognized

meaning, *e.g.*, involving living cells, *e.g.*, immortalized cells, primary cells, cell lines, and/or cells in an organism.

A gene “involved” in a disorder includes a gene, the normal or aberrant expression or function of which effects or causes a disease or disorder or at least one
5 symptom of said disease or disorder

Various methodologies of the instant invention include step that involves comparing a value, level, feature, characteristic, property, etc. to a “suitable control”, referred to interchangeably herein as an “appropriate control”. A “suitable control” or “appropriate control” is any control or standard familiar to one of ordinary skill in the art
10 useful for comparison purposes. In one embodiment, a “suitable control” or “appropriate control” is a value, level, feature, characteristic, property, etc. determined prior to performing an RNAi methodology, as described herein. For example, a transcription rate, mRNA level, translation rate, protein level, biological activity, cellular characteristic or property, genotype, phenotype, etc. can be determined prior to
15 introducing an RNAi agent of the invention into a cell or organism. In another embodiment, a “suitable control” or “appropriate control” is a value, level, feature, characteristic, property, etc. determined in a cell or organism, *e.g.*, a control or normal cell or organism, exhibiting, for example, normal traits. In yet another embodiment, a “suitable control” or “appropriate control” is a predefined value, level, feature,
20 characteristic, property, etc.

Various aspects of the invention are described in further detail in the following subsections.

25 I. RNA silencing agents

The present invention features RNA silencing agents (*e.g.* siRNAs or siRNA-like duplexes), in particular, RNA silencing agents having enhanced efficacy for gene silencing based on an asymmetry which promotes entry of a preferred strand into RISC.

30 1. Enhancement of RNA silencing agents

In certain aspects, the present invention features “small interfering RNA molecules” (“siRNA molecules” or “siRNA”), “miRNA” molecules, methods of making said siRNA and miRNA molecules, and methods (*e.g.*, research and/or therapeutic methods) for using said siRNA and miRNA molecules. An RNA silencing agent
5 (*e.g.* siRNA or miRNA molecule) of the invention is a duplex consisting of a sense strand and complementary antisense strand, the antisense strand having sufficient complementarity to a target mRNA to mediate RNAi or translational repression. Preferably, the strands are aligned such that there are at least 1, 2, or 3 bases at the end of the strands which do not align (*i.e.*, for which no complementary bases occur in the
10 opposing strand) such that an overhang of 1, 2 or 3 residues occurs at one or both ends of the duplex when strands are annealed. Preferably, the siRNA or miRNA molecule has a length from about 10-50 or more nucleotides, *i.e.*, each strand comprises 10-50 nucleotides (or nucleotide analogs). More preferably, the siRNA or miRNA molecule has a length from about 15-45 or 15-30 nucleotides. Even more preferably, the siRNA
15 or miRNA molecule has a length from about 16-25 or 18-23 nucleotides.

In other aspects, the invention features siRNA-like duplex molecules, as well as methods, for making and using such molecules in RNA silencing. siRNA-like duplexes (like siRNA duplexes) include a first or miRNA strand and a second or miRNA* strand and are structurally similar to siRNA duplexes. Modifications within the duplex are
20 permitted. Modifications that do not significantly affect RNA-silencing activity are particularly tolerated. siRNA-like duplexes mediate translational repression or RNAi depending on the degree of complementarity with target mRNA, as described *supra*.

RNA silencing agents (*e.g.* siRNAs or miRNAs) featured in the invention provide enhanced specificity and efficacy for mediating RISC-mediated cleavage or
25 translational repression of a desired target gene. In preferred aspect, the base pair strength between the antisense strand 5' end (AS 5') and the sense strand 3' end (S 3') of the siRNA, or miRNAs molecule is less than the bond strength or base pair strength between the antisense strand 3' end (AS 3') and the sense strand 5' end (S 5'), such that the antisense strand preferentially guides cleavage or translational repression of a target
30 mRNA.

In one embodiment, the bond strength or base-pair strength is less due to fewer G:C base pairs between the 5' end of the first or antisense strand and the 3' end of the

second or sense strand than between the 3' end of the first or antisense strand and the 5' end of the second or sense strand.

In another embodiment, the bond strength or base pair strength is less due to at least one mismatched base pair between the 5' end of the first or antisense strand and the 3' end of the second or sense strand. Preferably, the mismatched base pair is selected from the group consisting of G:A, C:A, C:U, G:G, A:A, C:C and U:U. In a related embodiment, the bond strength or base pair strength is less due to at least one wobble base pair, e.g., G:U, between the 5' end of the first or antisense strand and the 3' end of the second or sense strand.

10 In yet another embodiment, the bond strength or base pair strength is less due to at least one base pair comprising a rare nucleotide, e.g, inosine (I). Preferably, the base pair is selected from the group consisting of an I:A, I:U and I:C.

In yet another embodiment, the bond strength or base pair strength is less due to at least one base pair comprising a modified nucleotide. In preferred embodiments, the modified nucleotide is selected from the group consisting of 2-amino-G, 2-amino-A, 2,6-diamino-G, and 2,6-diamino-A.

2. Complementarity of RNA silencing agents with Target mRNA

20 The siRNA molecules of the invention further have a sequence that is "sufficiently complementary" to a target mRNA sequence to direct target-specific RNA interference (RNAi), as defined herein, *i.e.*, the siRNA has a sequence sufficient to trigger the destruction of the target mRNA by the RNAi machinery or process.

In general, siRNAcontaining nucleotide sequences sufficiently identical to a portion of the target gene to effect RISC-mediated cleavage of the target gene are preferred. 100% sequence identity between the siRNA and the target gene is not required to practice the present invention. The invention has the advantage of being able to tolerate certain sequence variations to enhance efficiency and specificity of RNAi. Moreover, siRNA sequences with insertions, deletions, and single point mutations relative to the target sequence can also be effective for inhibition. Alternatively, siRNA sequences with nucleotide analog substitutions or insertions can be effective for inhibition.

siRNA-like molecules of the invention have a sequence (i.e., have a strand having a sequence) that is “sufficiently complementary” to a target mRNA sequence to direct gene silencing either by RNAi or translational repression. As the degree of sequence identity between a miRNA sequence and the corresponding target gene
5 sequence is decreased, the tendency to mediate post-transcriptional gene silencing by translational repression rather than RNAi is increased. Therefore, in an alternative embodiment, where post-transcriptional gene silencing by translational repression of the target gene is desired, the miRNA sequence has partial complementarity with the target gene sequence. In certain embodiments, the miRNA sequence has partial
10 complementarity with one or more short sequences (complementarity sites) dispersed within the target mRNA (e.g. within the 3'-UTR of the target mRNA) (Hutvagner and Zamore, Science, 2002; Zeng et al., Mol. Cell, 2002; Zeng et al., RNA, 2003; Doench et al., Genes & Dev., 2003). Since the mechanism of translational repression is cooperative, multiple complementarity sites (e.g., 2, 3, 4, 5, or 6) may be targeted in
15 certain embodiments.

Sequence identity may determined by sequence comparison and alignment algorithms known in the art. To determine the percent identity of two nucleic acid sequences (or of two amino acid sequences), the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the first sequence or second
20 sequence for optimal alignment). The nucleotides (or amino acid residues) at corresponding nucleotide (or amino acid) positions are then compared. When a position in the first sequence is occupied by the same residue as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by
25 the sequences (*i.e.*, % homology = # of identical positions/total # of positions x 100), optionally penalizing the score for the number of gaps introduced and/or length of gaps introduced.

The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. In one embodiment,
30 the alignment generated over a certain portion of the sequence aligned having sufficient identity but not over portions having low degree of identity (*i.e.*, a local alignment). A preferred, non-limiting example of a local alignment algorithm utilized for the

comparison of sequences is the algorithm of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci. USA* 87:2264-68, modified as in Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-77. Such an algorithm is incorporated into the BLAST programs (version 2.0) of Altschul, *et al.* (1990) *J. Mol. Biol.* 215:403-10.

5 In another embodiment, the alignment is optimized by introducing appropriate gaps and percent identity is determined over the length of the aligned sequences (*i.e.*, a gapped alignment). To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, (1997) *Nucleic Acids Res.* 25(17):3389-3402. In another embodiment, the alignment is optimized by introducing
10 appropriate gaps and percent identity is determined over the entire length of the sequences aligned (*i.e.*, a global alignment). A preferred, non-limiting example of a mathematical algorithm utilized for the global comparison of sequences is the algorithm of Myers and Miller, *CABIOS* (1989). Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software
15 package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used.

Greater than 80% sequence identity, *e.g.*, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or even
20 100% sequence identity, between the siRNA antisense strand and the portion of the target gene is preferred. Conversely, miRNA sequences with less than 80% identity and the portion of the target gene (*i.e.* the site of complementarity) are preferred in order to mediate silencing by translational repression. Alternatively, the siRNA may be defined functionally as a nucleotide sequence (or oligonucleotide sequence) that is capable of
25 hybridizing with a portion of the target gene transcript (*e.g.*, 400 mM NaCl, 40 mM PIPES pH 6.4, 1 mM EDTA, 50°C or 70°C hybridization for 12-16 hours; followed by washing). Additional preferred hybridization conditions include hybridization at 70°C in 1xSSC or 50°C in 1xSSC, 50% formamide followed by washing at 70°C in 0.3xSSC or hybridization at 70°C in 4xSSC or 50°C in 4xSSC, 50% formamide followed by
30 washing at 67°C in 1xSSC. The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length should be 5-10°C less than the melting temperature (T_m) of the hybrid, where T_m is determined according to the following equations. For

hybrids less than 18 base pairs in length, $T_m(^{\circ}\text{C}) = 2(\# \text{ of A + T bases}) + 4(\# \text{ of G + C bases})$. For hybrids between 18 and 49 base pairs in length, $T_m(^{\circ}\text{C}) = 81.5 + 16.6(\log_{10}[\text{Na}^+]) + 0.41(\% \text{G+C}) - (600/\text{N})$, where N is the number of bases in the hybrid, and $[\text{Na}^+]$ is the concentration of sodium ions in the hybridization buffer ($[\text{Na}^+]$ for 1xSSC = 0.165 M). Additional examples of stringency conditions for polynucleotide hybridization are provided in Sambrook, J., E.F. Fritsch, and T. Maniatis, 1989, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, chapters 9 and 11, and *Current Protocols in Molecular Biology*, 1995, F.M. Ausubel et al., eds., John Wiley & Sons, Inc., sections 2.10 and 6.3-6.4, incorporated herein by reference. The length of the identical nucleotide sequences may be at least about 10, 12, 15, 17, 20, 22, 25, 27, 30, 32, 35, 37, 40, 42, 45, 47 or 50 bases.

The capacity of a siRNA-like duplex to mediate RNAi or translational repression may be predicted by the distribution of non-identical nucleotides between the target gene sequence and the nucleotide sequence of the silencing agent at the site of complementarity. In one embodiment, where gene silencing by translational repression is desired, at least one non-identical nucleotide is present in the central portion of the complementarity site so that duplex formed by the miRNA guide strand and the target mRNA contains a central "bulge" (Doench JG et al., *Genes & Dev.*, 2003). In another embodiment 2, 3, 4, 5, or 6 contiguous or non-contiguous non-identical nucleotides are introduced. The non-identical nucleotide may be selected such that it forms a wobble base pair (e.g., G:U) or a mismatched base pair (G:A, C:A, C:U, G:G, A:A, C:C, U:U). In a further preferred embodiment, the "bulge" is centered at nucleotide positions 12 and 13 from the 5' end of the miRNA molecule.

3. Modification of RNA silencing Agents

The RNA silencing molecules of the present invention can be modified to improve stability in serum or in growth medium for cell cultures. In order to enhance the stability, the 3'-residues may be stabilized against degradation, e.g., they may be selected such that they consist of purine nucleotides, particularly adenosine or guanosine nucleotides. Alternatively, substitution of pyrimidine nucleotides by modified

analogues, *e.g.*, substitution of uridine by 2'-deoxythymidine is tolerated and does not affect the efficiency of RNA interference.

In a preferred aspect, the invention features RNA silencing agents (*e.g.* small interfering RNAs (siRNAs) or siRNA-like molecules) that include first and second
5 strands wherein the second strand and/or first strand is modified by the substitution of internal nucleotides with modified nucleotides, such that *in vivo* stability is enhanced as compared to a corresponding unmodified agent (*e.g.* siRNA or siRNA-like duplex). As defined herein, an "internal" nucleotide is one occurring at any position other than the 5' end or 3' end of nucleic acid molecule, polynucleotide or oligonucleotide. An internal
10 nucleotide can be within a single-stranded molecule or within a strand of a duplex or double-stranded molecule. In one embodiment, the sense strand and/or antisense strand is modified by the substitution of at least one internal nucleotide. In another embodiment, the sense strand and/or antisense strand is modified by the substitution of at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or
15 more internal nucleotides. In another embodiment, the sense strand and/or antisense strand is modified by the substitution of at least 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or more of the internal nucleotides. In yet another embodiment, the sense strand and/or antisense strand is modified by the substitution of all of the internal nucleotides.

20 In a preferred embodiment of the present invention the RNA molecule may contain at least one modified nucleotide analogue. The nucleotide analogues may be located at positions where the target-specific silencing activity, *e.g.*, the RNAi mediating activity or translational repression activity is not substantially effected, *e.g.*, in a region at the 5'-end and/or the 3'-end of the RNA molecule. Particularly, the ends may be
25 stabilized by incorporating modified nucleotide analogues.

Preferred nucleotide analogues include sugar- and/or backbone-modified ribonucleotides (*i.e.*, include modifications to the phosphate-sugar backbone). For example, the phosphodiester linkages of natural RNA may be modified to include at least one of a nitrogen or sulfur heteroatom. In preferred backbone-modified
30 ribonucleotides the phosphoester group connecting to adjacent ribonucleotides is replaced by a modified group, *e.g.*, of phosphothioate group. In preferred sugar-modified ribonucleotides, the 2' OH-group is replaced by a group selected from H, OR,

R, halo, SH, SR, NH₂, NHR, NR₂ or ON, wherein R is C₁-C₆ alkyl, alkenyl or alkynyl and halo is F, Cl, Br or I.

Also preferred are nucleobase-modified ribonucleotides, *i.e.*, ribonucleotides, containing at least one non-naturally occurring nucleobase instead of a naturally occurring nucleobase. Bases may be modified to block the activity of adenosine deaminase. Exemplary modified nucleobases include, but are not limited to, uridine and/or cytidine modified at the 5-position, *e.g.*, 5-(2-amino)propyl uridine, 5-bromo uridine; adenosine and/or guanosines modified at the 8 position, *e.g.*, 8-bromo guanosine; deaza nucleotides, *e.g.*, 7-deaza-adenosine; O- and N-alkylated nucleotides, *e.g.*, N⁶-methyl adenosine are suitable. It should be noted that the above modifications may be combined.

4. Production of RNA silencing Agents

RNA may be produced enzymatically or by partial/total organic synthesis, any modified ribonucleotide can be introduced by *in vitro* enzymatic or organic synthesis. In one embodiment, a silencing agent (*e.g.* a RNAi agent or translational repression agent) is prepared chemically. Methods of synthesizing RNA molecules are known in the art, in particular, the chemical synthesis methods as described in Verma and Eckstein (1998) *Annul Rev. Biochem.* 67:99-134.

In one embodiment, a RNA silencing agent (*e.g.* siRNA or siRNA-like duplex) is prepared enzymatically. For example, a short duplex RNA (*e.g.* siRNA or siRNA-like duplex) can be prepared by enzymatic processing of a long ds RNA having sufficient complementarity to the desired target mRNA. Processing of long ds RNA can be accomplished *in vitro*, for example, using appropriate cellular lysates and duplex RNA (*e.g.* siRNA or siRNA-like duplex) can be subsequently purified by gel electrophoresis or gel filtration. Duplex RNA (*e.g.* siRNA or siRNA-like duplex) can then be denatured according to art-recognized methodologies. In an exemplary embodiment, RNA can be purified from a mixture by extraction with a solvent or resin, precipitation, electrophoresis, chromatography, or a combination thereof. Alternatively, the RNA may be used with no or a minimum of purification to avoid losses due to sample processing.

Alternatively, the RNA can also be prepared by enzymatic transcription from synthetic DNA templates or from DNA plasmids isolated from recombinant bacteria. Typically, phage RNA polymerases are used such as T7, T3 or SP6 RNA polymerase (Milligan and Uhlenbeck (1989) *Methods Enzymol.* 180:51-62). The RNA may be dried for storage or dissolved in an aqueous solution. The solution may contain buffers or salts to inhibit annealing, and/or promote stabilization of the single strands.

RNA silencing agents of the invention can also be prepared *in vivo* by enzymatic processing of a long dsRNA molecule (>30 b.p.) which has sufficient complementarity to the desired target mRNA. Preferably, *in vivo* processing of the long dsRNA molecule occurs in a non-mammalian cell or a mammalian cell which is deficient in the interferon-mediated inflammatory response to dsRNA. In one embodiment, the cell capable of dsRNA enzymatic processing may be present within an organism such that dsRNA processing can be induced *in vivo* to trigger gene silencing of a target gene within the organism. Alternatively, the cell (*i.e.* a host cell) containing endogenous machinery for dsRNA processing (*e.g.* DICER) or transformed with heterologous genes to enable dsRNA processing) be cultured and induced to process dsRNA *in vitro*. RNA silencing agents may then be purified from the host cell following dsRNA processing for administration to an organism containing the target gene to be silenced.

In another embodiment, RNA silencing agents (*e.g.* siRNAs or siRNA-like duplexes) are synthesized directly either *in vivo*, *in situ*, or *in vitro*. An endogenous RNA polymerase in the cell may mediate transcription of the RNA silencing agent *in vivo* or *in situ*, or a cloned RNA polymerase can be used for transcription of the RNA silencing agent *in vivo* or *in vitro*. For transcription from a transgene *in vivo* or an expression construct, a regulatory region (*e.g.*, promoter, enhancer, silencer, splice donor and acceptor, polyadenylation) may be used to transcribe the RNA silencing agent (*e.g.* siRNA or siRNA-like duplexes). Inhibition may be targeted by specific transcription in an organ, tissue, or cell type; stimulation of an environmental condition (*e.g.*, infection, stress, temperature, chemical inducers); and/or engineering transcription at a developmental stage or age. A transgenic organism that expresses a RNA silencing agent (*e.g.* siRNA or siRNA-like duplexes) from a recombinant construct may be produced by introducing the construct into a zygote, an embryonic stem cell, or another multipotent cell derived from the appropriate organism.

5. Targeted mRNAs of RNA silencing agents

In one embodiment, the target mRNA of the invention specifies the amino acid sequence of a cellular protein (*e.g.*, a nuclear, cytoplasmic, transmembrane, or membrane-associated protein). In another embodiment, the target mRNA of the invention specifies the amino acid sequence of an extracellular protein (*e.g.*, an extracellular matrix protein or secreted protein). As used herein, the phrase “specifies the amino acid sequence” of a protein means that the mRNA sequence is translated into the amino acid sequence according to the rules of the genetic code. The following classes of proteins are listed for illustrative purposes: developmental proteins (*e.g.*, adhesion molecules, cyclin kinase inhibitors, Wnt family members, Pax family members, Winged helix family members, Hox family members, cytokines/lymphokines and their receptors, growth/differentiation factors and their receptors, neurotransmitters and their receptors); oncogene-encoded proteins (*e.g.*, ABLI, BCLI, BCL2, BCL6, CBFA2, CBL, CSFIR, ERBA, ERBB, EBRB2, ETSI, ETV6, FGR, FOS, FYN, HCR, HRAS, JUN, KRAS, LCK, LYN, MDM2, MLL, MYB, MYC, MYCLI, MYCN, NRAS, PIM I, PML, RET, SRC, TALI, TCL3, and YES); tumor suppressor proteins (*e.g.*, APC, BRCA1, BRCA2, MADH4, MCC, NF I, NF2, RB I, TP53, and WTI); and enzymes (*e.g.*, ACC synthases and oxidases, ACP desaturases and hydroxylases, ADP-glucose pyrophorylases, ATPases, alcohol dehydrogenases, amylases, amyloglucosidases, catalases, cellulases, chalcone synthases, chitinases, cyclooxygenases, decarboxylases, dextriinases, DNA and RNA polymerases, galactosidases, glucanases, glucose oxidases, granule-bound starch synthases, GTPases, helicases, hemicellulases, integrases, inulinases, invertases, isomerases, kinases, lactases, lipases, lipxygenases, lysozymes, nopaline synthases, octopine synthases, pectinesterases, peroxidases, phosphatases, phospholipases, phosphorylases, phytases, plant growth regulator synthases, polygalacturonases, proteinases and peptidases, pullanases, recombinases, reverse transcriptases, RUBISCOs, topoisomerases, and xylanases).

In a preferred aspect of the invention, the target mRNA molecule of the invention specifies the amino acid sequence of a protein associated with a pathological condition. For example, the protein may be a pathogen-associated protein (*e.g.*, a viral

protein involved in immunosuppression of the host, replication of the pathogen, transmission of the pathogen, or maintenance of the infection), or a host protein which facilitates entry of the pathogen into the host, drug metabolism by the pathogen or host, replication or integration of the pathogen's genome, establishment or spread of infection
5 in the host, or assembly of the next generation of pathogen. Alternatively, the protein may be a tumor-associated protein or an autoimmune disease-associated protein.

In one embodiment, the target mRNA molecule of the invention specifies the amino acid sequence of an endogenous protein (*i.e.*, a protein present in the genome of a cell or organism). In another embodiment, the target mRNA molecule of the invention
10 specified the amino acid sequence of a heterologous protein expressed in a recombinant cell or a genetically altered organism. In another embodiment, the target mRNA molecule of the invention specified the amino acid sequence of a protein encoded by a transgene (*i.e.*, a gene construct inserted at an ectopic site in the genome of the cell). In yet another embodiment, the target mRNA molecule of the invention specifies the amino
15 acid sequence of a protein encoded by a pathogen genome which is capable of infecting a cell or an organism from which the cell is derived.

By inhibiting the expression of such proteins, valuable information regarding the function of said proteins and therapeutic benefits which may be obtained from said inhibition may be obtained.

20

II. Short hairpin RNAs (shRNAs)

In certain featured embodiments, the instant invention provides shRNAs having enhanced specificity or efficacy in mediating gene silencing (*e.g.* RNAi or translational
25 repression). In contrast to short RNA silencing duplexes (*e.g.* siRNA or siRNA-like duplexes), short hairpin RNAs (shRNAs) mimic the natural precursors of miRNAs and enter at the top of the gene silencing pathway. For this reason, shRNAs are believed to mediate gene silencing more efficiently by being fed through the entire natural gene silencing pathway.

30 A preferred shRNA of the invention is one that has been redesigned for increased specificity or enhancement relative to a previous shRNA. The new shRNA differs from

a previous shRNA in that a silencing duplex produced from the new shRNA has less base pair strength between the 5' end of the antisense strand or first strand and the 3' end of the sense strand or second strand than the base pair strength between the 3' end of the antisense strand or first strand and the 5' end of the sense strand or second strand.

5

1. Engineered RNA Precursors That Generate RNA silencing Agents

Naturally-occurring miRNA precursors (pre-miRNA) have a single strand that forms a duplex stem including two portions that are generally complementary, and a
 10 loop, that connects the two portions of the stem. In typical pre-miRNAs, the stem includes one or more bulges, e.g., extra nucleotides that create a single nucleotide "loop" in one portion of the stem, and/or one or more unpaired nucleotides that create a gap in the hybridization of the two portions of the stem to each other. Short hairpin RNAs, or engineered RNA precursors, of the invention are artificial constructs based on these
 15 naturally occurring pre-miRNAs, but which are engineered to deliver desired RNA silencing agents (*e.g.*, siRNAs or siRNA-like duplexes).

In shRNAs, or engineered precursor RNAs, of the instant invention, one portion of the duplex stem is a nucleic acid sequence that is complementary (or anti-sense) to the target mRNA. Thus, engineered RNA precursors include a duplex stem with two
 20 portions and a loop connecting the two stem portions. The two stem portions are about 18 or 19 to about 25, 30, 35, 37, 38, 39, or 40 or more nucleotides in length. When used in mammalian cells, the length of the stem portions should be less than about 30 nucleotides to avoid provoking non-specific responses like the interferon pathway. In non-mammalian cells, the stem can be longer than 30 nucleotides. In fact, the stem can
 25 include much larger sections complementary to the target mRNA (up to, and including the entire mRNA). The two portions of the duplex stem must be sufficiently complementary to hybridize to form the duplex stem. Thus, the two portions can be, but need not be, fully or perfectly complementary. In addition, the two stem portions can be the same length, or one portion can include an overhang of 1, 2, 3, or 4 nucleotides. The
 30 overhanging nucleotides can include, for example, uracils (Us), *e.g.*, all Us. The loop in the shRNAs or engineered RNA precursors may differ from natural pre-miRNA

sequences by modifying the loop sequence to increase or decrease the number of paired nucleotides, or replacing all or part of the loop sequence with a tetraloop or other loop sequences. Thus, the loop in the shRNAs or engineered RNA precursors can be 2, 3, 4, 5, 6, 7, 8, 9, or more, *e.g.*, 15 or 20, or more nucleotides in length.

5 In certain embodiments, shRNAs of the invention include the sequences of a desired RNA silencing agent (*e.g.* siRNA or siRNA-like duplex). The desired RNA silencing duplex (*e.g.* siRNA or siRNA-like duplex), and thus both of the two stem portions in the engineered RNA precursor, are selected by methods known in the art. These include, but are not limited to, selecting an 18, 19, 20, 21 nucleotide, or longer, sequence from the target gene mRNA sequence from a region 100 to 200 or 300
10 nucleotides on the 3' side of the start of translation. In general, the sequence can be selected from any portion of the mRNA from the target gene, such as the 5' UTR (untranslated region), coding sequence, or 3' UTR. This sequence can optionally follow immediately after a region of the target gene containing two adjacent AA nucleotides.
15 The last two nucleotides of the 21 or so nucleotide sequence can be selected to be UU (so that the anti-sense strand of the siRNA begins with UU). This 21 or so nucleotide sequence is used to create one portion of a duplex stem in the engineered RNA precursor. This sequence can replace a stem portion of a wild-type pre-stRNA sequence, *e.g.*, enzymatically, or is included in a complete sequence that is synthesized. For
20 example, one can synthesize DNA oligonucleotides that encode the entire stem-loop engineered RNA precursor, or that encode just the portion to be inserted into the duplex stem of the precursor, and using restriction enzymes to build the engineered RNA precursor construct, *e.g.*, from a wild-type pre-stRNA.

Engineered RNA precursors include in the duplex stem the 21-22 or so
25 nucleotide sequences of the siRNA or siRNA-like duplex desired to be produced *in vivo*. Thus, the stem portion of the engineered RNA precursor includes at least 18 or 19 nucleotide pairs corresponding to the sequence of an exonic portion of the gene whose expression is to be reduced or inhibited. The two 3' nucleotides flanking this region of the stem are chosen so as to maximize the production of the siRNA from the engineered
30 RNA precursor, and to maximize the efficacy of the resulting siRNA in targeting the corresponding mRNA for translational repression or destruction by RNAi *in vivo* and *in vitro*.

In certain embodiments, shRNAs of the invention include miRNA sequences, optionally end-modified miRNA sequences, to enhance entry into RISC. The miRNA sequence can be similar or identical to that of any naturally occurring miRNA (see *e.g.* The miRNA Registry; Griffiths-Jones S, *Nuc. Acids Res.*, 2004). Over one thousand
5 natural miRNAs have been identified to date and together they are thought to comprise ~1% of all predicted genes in the genome. Many natural miRNAs are clustered together in the introns of pre-mRNAs and can be identified *in silico* using homology-based searches (Pasquinelli et al., 2000; Lagos-Quintana et al., 2001; Lau et al., 2001; Lee and Ambros, 2001) or computer algorithms (*e.g.* MiRScan, MiRSeeker) that predict the
10 capability of a candidate miRNA gene to form the stem loop structure of a pri-mRNA (Grad et al., *Mol. Cell.*, 2003; Lim et al., *Genes Dev.*, 2003; Lim et al., *Science*, 2003; Lai EC et al., *Genome Bio.*, 2003). An online registry provides a searchable database of all published miRNA sequences (The miRNA Registry at the Sanger Institute website; Griffiths-Jones S, *Nuc. Acids Res.*, 2004). Exemplary, natural miRNAs include lin-4,
15 let-7, miR-10, mirR-15, miR-16, miR-168, miR-175, miR-196 and their homologs, as well as other natural miRNAs from humans and certain model organisms including *Drosophila melanogaster*, *Caenorhabditis elegans*, zebrafish, *Arabidopsis thaliana*, mouse, and rat as described in International PCT Publication No. WO 03/029459.

Naturally-occurring miRNAs are expressed by endogenous genes *in vivo* and are
20 processed from a hairpin or stem-loop precursor (pre-miRNA or pri-miRNAs) by Dicer or other RNases (Lagos-Quintana et al., *Science*, 2001; Lau et al., *Science*, 2001; Lee and Ambros, *Science*, 2001; Lagos-Quintana et al., *Curr. Biol.*, 2002; Mourelatos et al., *Genes Dev.*, 2002; Reinhart et al., *Science*, 2002; Ambros et al., *Curr. Biol.*, 2003; Brennecke et al., 2003; Lagos-Quintana et al., *RNA*, 2003; Lim et al., *Genes Dev.*, 2003;
25 Lim et al., *Science*, 2003). miRNAs can exist transiently *in vivo* as a double-stranded duplex but only one strand is taken up by the RISC complex to direct gene silencing. Certain miRNAs, *e.g.* plant miRNAs, have perfect or near-perfect complementarity to their target mRNAs and, hence, direct cleavage of the target mRNAs. Other miRNAs have less than perfect complementarity to their target mRNAs and, hence, direct
30 translational repression of the target mRNAs. The degree of complementarity between an miRNA and its target mRNA is believed to determine its mechanism of action. For example, perfect or near-perfect complementarity between a miRNA and its target

mRNA is predictive of a cleavage mechanism (Yekta *et al.*, *Science*, 2004), whereas less than perfect complementarity is predictive of a translational repression mechanism.

In particular embodiments, the miRNA sequence is that of a naturally-occurring miRNA sequence, the aberrant expression or activity of which is correlated with a
5 miRNA disorder.

Another defining feature of these engineered RNA precursors is that as a consequence of their length, sequence, and/or structure, they do not induce sequence non-specific responses, such as induction of the interferon response or apoptosis, or that they induce a lower level of such sequence non-specific responses than long, double-
10 stranded RNA (>150bp) that has been used to induce post-transcriptional gene silencing (*e.g.* RNAi). For example, the interferon response is triggered by dsRNA longer than 30 base pairs.

2. Transgenes Encoding Engineered RNA Precursors

15 The new engineered RNA precursors can be synthesized by standard methods known in the art, *e.g.*, by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). These synthetic, engineered RNA precursors can be used directly as described below or cloned into expression vectors by methods known in the field. The engineered RNA precursors
20 should be delivered to cells *in vitro* or *in vivo* in which it is desired to target a specific mRNA for silencing (*e.g.* destruction or repression). A number of methods have been developed for delivering DNA or RNA to cells. For example, for *in vivo* delivery, molecules can be injected directly into a tissue site or administered systemically. *In vitro* delivery includes methods known in the art such as electroporation and lipofection.

25 To achieve intracellular concentrations of the nucleic acid molecule sufficient to suppress expression of endogenous mRNAs, one can use, for example, a recombinant DNA construct in which the oligonucleotide is placed under the control of a strong *Pol* III (*e.g.*, U6 or *Pol* III H1-RNA promoter) or *Pol* II promoter. The use of such a construct to transfect target cells *in vitro* or *in vivo* will result in the transcription of sufficient
30 amounts of the engineered RNA precursor to lead to the production of a RNA duplex (*e.g.* siRNA or siRNA-like duplex) that can target a corresponding mRNA sequence for cleavage (*i.e.* RNAi) or translational repression to decrease the expression of the gene

encoding that mRNA. For example, a vector can be introduced *in vivo* such that it is taken up by a cell and directs the transcription of an engineered RNA precursor. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired stRNA precursor.

5 Such vectors can be constructed by recombinant DNA technology methods known in the art. Vectors can be plasmid, viral, or other vectors known in the art such as those described herein, used for replication and expression in mammalian cells or other targeted cell types. The nucleic acid sequences encoding the engineered RNA precursors can be prepared using known techniques. For example, two synthetic DNA
10 oligonucleotides can be synthesized to create a novel gene encoding the entire engineered RNA precursor. The DNA oligonucleotides, which will pair, leaving appropriate 'sticky ends' for cloning, can be inserted into a restriction site in a plasmid that contains a promoter sequence (e.g., a *Pol* II or a *Pol* III promoter) and appropriate terminator sequences 3' to the engineered RNA precursor sequences (e.g., a cleavage and
15 polyadenylation signal sequence from SV40 or a *Pol* III terminator sequence).

 The invention also encompasses genetically engineered host cells that contain any of the foregoing expression vectors and thereby express the nucleic acid molecules of the invention in the host cell. The host cells can be cultured using known techniques and methods (see, e.g., Culture of Animal Cells (R.I. Freshney, Alan R. Liss, Inc. 1987);
20 Molecular Cloning, Sambrook et al. (Cold Spring Harbor Laboratory Press, 1989)).

 Successful introduction of the vectors of the invention into host cells can be monitored using various known methods. For example, transient transfection can be signaled with a reporter, such as a fluorescent marker, such as Green Fluorescent Protein (GFP). Stable transfection can be indicated using markers that provide the transfected
25 cell with resistance to specific environmental factors (e.g., antibiotics and drugs), such as hygromycin B resistance, e.g., in insect cells and in mammalian cells.

3. Regulatory Sequences

 The expression of the engineered RNA precursors is driven by regulatory
30 sequences, and the vectors of the invention can include any regulatory sequences known in the art to act in mammalian cells, e.g., human or murine cells; in insect cells; in plant

cells; or other cells. The term regulatory sequence includes promoters, enhancers, and other expression control elements. It will be appreciated that the appropriate regulatory sequence depends on such factors as the future use of the cell or transgenic animal into which a sequence encoding an engineered RNA precursor is being introduced, and the level of expression of the desired RNA precursor. A person skilled in the art would be able to choose the appropriate regulatory sequence. For example, the transgenic animals described herein can be used to determine the role of a test polypeptide or the engineered RNA precursors in a particular cell type, *e.g.*, a hematopoietic cell. In this case, a regulatory sequence that drives expression of the transgene ubiquitously, or a hematopoietic-specific regulatory sequence that expresses the transgene only in hematopoietic cells, can be used. Expression of the engineered RNA precursors in a hematopoietic cell means that the cell is now susceptible to specific, targeted silencing (*e.g.* RNAi) of a particular gene. Examples of various regulatory sequences are described below.

The regulatory sequences can be inducible or constitutive. Suitable constitutive regulatory sequences include the regulatory sequence of a housekeeping gene such as the α -actin regulatory sequence, or may be of viral origin such as regulatory sequences derived from mouse mammary tumor virus (MMTV) or cytomegalovirus (CMV).

Alternatively, the regulatory sequence can direct transgene expression in specific organs or cell types (see, *e.g.*, Lasko et al., 1992, *Proc. Natl. Acad. Sci. USA* 89:6232). Several tissue-specific regulatory sequences are known in the art including the albumin regulatory sequence for liver (Pinkert et al., 1987, *Genes Dev.* 1:268276); the endothelin regulatory sequence for endothelial cells (Lee, 1990, *J. Biol. Chem.* 265:10446-50); the keratin regulatory sequence for epidermis; the myosin light chain-2 regulatory sequence for heart (Lee et al., 1992, *J. Biol. Chem.* 267:15875-85), and the insulin regulatory sequence for pancreas (Bucchini et al., 1986, *Proc. Natl. Acad. Sci. USA* 83:2511-2515), or the vav regulatory sequence for hematopoietic cells (Oligvy et al., 1999, *Proc. Natl. Acad. Sci. USA* 96:14943-14948). Another suitable regulatory sequence, which directs constitutive expression of transgenes in cells of hematopoietic origin, is the murine MHC class I regulatory sequence (Morello et al., 1986, *EMBO J.* 5:1877-1882). Since NMC expression is induced by cytokines, expression of a test gene operably linked to this regulatory sequence can be upregulated in the presence of cytokines.

In addition, expression of the transgene can be precisely regulated, for example, by using an inducible regulatory sequence and expression systems such as a regulatory sequence that is sensitive to certain physiological regulators, *e.g.*, circulating glucose levels, or hormones (Docherty *et al.*, 1994, *FASEB J.* 8:20-24). Such inducible
 5 expression systems, suitable for the control of transgene expression in cells or in mammals such as mice, include regulation by ecdysone, by estrogen, progesterone, tetracycline, chemical inducers of dimerization, and isopropyl-beta-D1 - thiogalactopyranoside (IPTG) (collectively referred to as "the regulatory molecule"). Each of these expression systems is well described in the literature and permits
 10 expression of the transgene throughout the animal in a manner controlled by the presence or absence of the regulatory molecule. For a review of inducible expression systems, see, *e.g.*, Mills, 2001, *Genes Devel.* 15:1461-1467, and references cited therein.

The regulatory elements referred to above include, but are not limited to, the cytomegalovirus hCMV immediate early gene, the early or late promoters of SV40
 15 adenovirus (Bernoist *et al.*, *Nature*, 290:304, 1981), the tet system, the lac system, the trp system, the TAC system, the TRC system; the major operator and promoter regions of phage A, the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase, the promoters of acid phosphatase, and the promoters of the yeast α -mating factors. Additional promoters include the promoter contained in the 3' long terminal
 20 repeat of Rous sarcoma virus (Yamamoto *et al.*, *Cell* 22:787-797, 1988); the herpes thymidine kinase promoter (Wagner *et al.*, *Proc. Natl. Acad. Sci. USA* 78:1441, 1981); or the regulatory sequences of the metallothionein gene (Brinster *et al.*, *Nature* 296:39, 1988).

25 4. Assay for Testing Engineered RNA Precursors

Drosophila embryo lysates can be used to determine if an engineered RNA precursor was, in fact, the direct precursor of a mature stRNA or siRNA. This lysate assay is described in Tuschl *et al.*, 1999, *supra*, Zamore *et al.*, 2000, *supra*, and Hutvagner *et al.* 2001, *supra*. These lysates recapitulate RNAi *in vitro*, thus permitting
 30 investigation into whether the proposed precursor RNA was cleaved into a mature stRNA or siRNA by an RNAi-like mechanism. Briefly, the precursor RNA is incubated with *Drosophila* embryo lysate for various times, then assayed for the production of the

mature siRNA or stRNA by primer extension or Northern hybridization. As in the *in vivo* setting, mature RNA accumulates in the cell-free reaction. Thus, an RNA corresponding to the proposed precursor can be shown to be converted into a mature stRNA or siRNA duplex in the *Drosophila* embryo lysate.

5 Furthermore, an engineered RNA precursor can be functionally tested in the *Drosophila* embryo lysates. In this case, the engineered RNA precursor is incubated in the lysate in the presence of a 5' radiolabeled target mRNA in a standard *in vitro* RNAi reaction for various lengths of time. The target mRNA can be 5' radiolabeled using guanylyl transferase (as described in Tuschl *et al.*, 1999, *supra* and references therein) or
10 other suitable methods. The products of the *in vitro* reaction are then isolated and analyzed on a denaturing acrylamide or agarose gel to determine if the target mRNA has been cleaved in response to the presence of the engineered RNA precursor in the reaction. The extent and position of such cleavage of the mRNA target will indicate if the engineering of the precursor created a pre-siRNA capable of mediating sequence-
15 specific RNAi.

IV. Methods of Introducing RNAs, Vectors, and Host Cells

Physical methods of introducing nucleic acids include injection of a solution containing the RNA, bombardment by particles covered by the RNA, soaking the cell or
20 organism in a solution of the RNA, or electroporation of cell membranes in the presence of the RNA. A viral construct packaged into a viral particle would accomplish both efficient introduction of an expression construct into the cell and transcription of RNA encoded by the expression construct. Other methods known in the art for introducing nucleic acids to cells may be used, such as lipid-mediated carrier transport, chemical-
25 mediated transport, such as calcium phosphate, and the like. Thus the RNA may be introduced along with components that perform one or more of the following activities: enhance RNA uptake by the cell, inhibit annealing of single strands, stabilize the single strands, or other-wise increase inhibition of the target gene.

RNA may be directly introduced into the cell (*i.e.*, intracellularly); or introduced
30 extracellularly into a cavity, interstitial space, into the circulation of an organism, introduced orally, or may be introduced by bathing a cell or organism in a solution

containing the RNA. Vascular or extravascular circulation, the blood or lymph system, and the cerebrospinal fluid are sites where the RNA may be introduced.

The cell with the target gene may be derived from or contained in any organism. The organism may a plant, animal, protozoan, bacterium, virus, or fungus. The plant
 5 may be a monocot, dicot or gymnosperm; the animal may be a vertebrate or invertebrate. Preferred microbes are those used in agriculture or by industry, and those that are pathogenic for plants or animals. Fungi include organisms in both the mold and yeast morphologies. Plants include arabidopsis; field crops (*e.g.*, alfalfa, barley, bean, com, cotton, flax, pea, rape, nice, rye, safflower, sorghum, soybean, sunflower, tobacco, and
 10 wheat); vegetable crops (*e.g.*, asparagus, beet, broccoli, cabbage, carrot, cauliflower, celery, cucumber, eggplant, lettuce, onion, pepper, potato, pumpkin, radish, spinach, squash, taro, tomato, and zucchini); fruit and nut crops (*e.g.*, almond, apple, apricot, banana, black- berry, blueberry, cacao, cherry, coconut, cranberry, date, faJoa, filbert, grape, grapefruit, guava, kiwi, lemon, lime, mango, melon, nectarine, orange, papaya,
 15 passion fruit, peach, peanut, pear, pineapple, pistachio, plum, raspberry, strawberry, tangerine, walnut, and watermelon); and ornamentals (*e.g.*, alder, ash, aspen, azalea, birch, boxwood, camellia, carnation, chrysanthemum, elm, fir, ivy, jasmine, juniper, oak, palm, poplar, pine, redwood, rhododendron, rose, and rubber). Examples of vertebrate animals include fish, mammal, cattle, goat, pig, sheep, rodent, hamster, mouse, rat,
 20 primate, and human; invertebrate animals include nematodes, other worms, drosophila, and other insects.

The skilled artisan will appreciate that the enumerated organisms are also useful for practicing other aspects of the invention, *e.g.*, making transgenic organisms as described *infra*.

25 The cell having the target gene may be from the germ line or somatic, totipotent or pluripotent, dividing or non-dividing, parenchyma or epithelium, immortalized or transformed, or the like. The cell may be a stem cell or a differentiated cell. Cell types that are differentiated include adipocytes, fibroblasts, myocytes, cardiomyocytes, endothelium, neurons, glia, blood cells, megakaryocytes, lymphocytes, macrophages,
 30 neutrophils, eosinophils, basophils, mast cells, leukocytes, granulocytes, keratinocytes, chondrocytes, osteoblasts, osteoclasts, hepatocytes, and cells of the endocrine or exocrine glands.

Depending on the particular target gene and the dose of double stranded RNA material delivered, this process may provide partial or complete loss of function for the target gene. A reduction or loss of gene expression in at least 50%, 60%, 70%, 80%, 90%, 95% or 99% or more of targeted cells is exemplary. Inhibition of gene expression
5 refers to the absence (or observable decrease) in the level of protein and/or mRNA product from a target gene. Specificity refers to the ability to inhibit the target gene without manifest effects on other genes of the cell. The consequences of inhibition can be confirmed by examination of the outward properties of the cell or organism (as presented below in the examples) or by biochemical techniques such as RNA solution
10 hybridization, nuclease protection, Northern hybridization, reverse transcription, gene expression monitoring with a microarray, antibody binding, enzyme linked immunosorbent assay (ELISA), Western blotting, radioimmunoassay (RIA), other immunoassays, and fluorescence activated cell analysis (FACS).

For RNA-mediated inhibition in a cell line or whole organism, gene expression is
15 conveniently assayed by use of a reporter or drug resistance gene whose protein product is easily assayed. Such reporter genes include acetohydroxyacid synthase (AHAS), alkaline phosphatase (AP), beta galactosidase (LacZ), beta glucuronidase (GUS), chloramphenicol acetyltransferase (CAT), green fluorescent protein (GFP), horseradish peroxidase (HRP), luciferase (Luc), nopaline synthase (NOS), octopine synthase (OCS),
20 and derivatives thereof. Multiple selectable markers are available that confer resistance to ampicillin, bleomycin, chloramphenicol, gentamycin, hygromycin, kanamycin, lincomycin, methotrexate, phosphinothricin, puromycin, and tetracyclin. Depending on the assay, quantitation of the amount of gene expression allows one to determine a degree of inhibition which is greater than 10%, 33%, 50%, 90%, 95% or 99% as
25 compared to a cell not treated according to the present invention. Lower doses of injected material and longer times after administration of a RNA silencing agent (*e.g.*, an RNAi agent) may result in inhibition in a smaller fraction of cells (*e.g.*, at least 10%, 20%, 50%, 75%, 90%, or 95% of targeted cells). Quantitation of gene expression in a cell may show similar amounts of inhibition at the level of accumulation of target
30 mRNA or translation of target protein. As an example, the efficiency of inhibition may be determined by assessing the amount of gene product in the cell; mRNA may be detected with a hybridization probe having a nucleotide sequence outside the region

used for the inhibitory double-stranded RNA, or translated polypeptide may be detected with an antibody raised against the polypeptide sequence of that region.

The RNA may be introduced in an amount which allows delivery of at least one copy per cell. Higher doses (*e.g.*, at least 5, 10, 100, 500 or 1000 copies per cell) of material may yield more effective inhibition; lower doses may also be useful for specific applications.

V. Methods of Treatment:

The present invention provides for both prophylactic and therapeutic methods of treating a subject at risk of (or susceptible to) a disease or disorder or having a disease or disorder associated with aberrant or unwanted target gene expression or activity. The invention further provides methods of treating cell or subject having (or susceptible to) an miRNA disorder.

"Treatment", or "treating" as used herein, is defined as the application or administration of a therapeutic agent (*e.g.*, a RNAi agent or agent mediating translational control, or vector or transgene encoding same) to a patient, or application or administration of a therapeutic agent to an isolated tissue or cell line from a patient, who has a disease or disorder, a symptom of disease or disorder or a predisposition toward a disease or disorder, with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve or affect the disease or disorder, the symptoms of the disease or disorder, or the predisposition toward disease.

With regards to both prophylactic and therapeutic methods of treatment, such treatments may be specifically tailored or modified, based on knowledge obtained from the field of pharmacogenomics. "Pharmacogenomics", as used herein, refers to the application of genomics technologies such as gene sequencing, statistical genetics, and gene expression analysis to drugs in clinical development and on the market. More specifically, the term refers the study of how a patient's genes determine his or her response to a drug (*e.g.*, a patient's "drug response phenotype", or "drug response genotype"). Thus, another aspect of the invention provides methods for tailoring an individual's prophylactic or therapeutic treatment with either the target gene molecules of the present invention or target gene modulators according to that individual's drug

response genotype. Pharmacogenomics allows a clinician or physician to target prophylactic or therapeutic treatments to patients who will most benefit from the treatment and to avoid treatment of patients who will experience toxic drug-related side effects.

5 1. Prophylactic Methods

In one aspect, the invention provides a method for preventing in a subject, a disease or condition associated with an aberrant or unwanted target gene expression or activity, by administering to the subject a therapeutic agent (*e.g.*, a RNAi agent or agent mediating translational control, or vector or transgene encoding same). Subjects at risk
10 for a disease which is caused or contributed to by aberrant or unwanted target gene expression or activity can be identified by, for example, any or a combination of diagnostic or prognostic assays as described herein. Administration of a prophylactic agent can occur prior to the manifestation of symptoms characteristic of the target gene aberrancy, such that a disease or disorder is prevented or, alternatively, delayed in its
15 progression. Depending on the type of target gene aberrancy, for example, a target gene, target gene agonist or target gene antagonist agent can be used for treating the subject. The appropriate agent can be determined based on screening assays described herein.

2. Therapeutic Methods

Another aspect of the invention pertains to methods of modulating target gene
20 expression, protein expression or activity for therapeutic purposes. Accordingly, in an exemplary embodiment, the modulatory method of the invention involves contacting a cell capable of expressing target gene with a therapeutic agent (*e.g.*, a RNAi agent or agent mediating translational control, or vector or transgene encoding same) that is specific for the target gene or protein (*e.g.*, is specific for the mRNA encoded by said
25 gene or specifying the amino acid sequence of said protein) such that expression or one or more of the activities of target protein is modulated. These modulatory methods can be performed *in vitro* (*e.g.*, by culturing the cell with the agent) or, alternatively, *in vivo* (*e.g.*, by administering the agent to a subject). As such, the present invention provides methods of treating an individual afflicted with a disease or disorder characterized by
30 aberrant or unwanted expression or activity of a target gene polypeptide or nucleic acid molecule. Inhibition of target gene activity is desirable in situations in which target gene

is abnormally unregulated and/or in which decreased target gene activity is likely to have a beneficial effect.

3. miRNA Therapy

5 Another aspect of the invention pertains to methods of treating a cell or subject having or susceptible to an miRNA disorder. MiRNA disorders can result, for example, when a naturally occurring miRNA is not present in a cell or subject in an amount sufficient to regulate cellular processes controlled by the miRNA. In an exemplary embodiment, the method of the invention involves administering to a subject having or
10 susceptible to an miRNA disorder an effective amount of a RNA silencing agent of the invention (*e.g.*, a siRNA-like duplex) or vector or transgene encoding same, the silencing agent delivering at least one strand (*e.g.*, a miRNA strand of the siRNA-like agent) to the subject (*e.g.*, to the RISC complex of said subject). In another exemplary embodiment, the method of the invention involves contacting a cell from a subject
15 having or susceptible to an miRNA disorder with an effective amount of a RNA silencing agent of the invention (*e.g.*, a siRNA-like duplex) or vector or transgene encoding same, the silencing agent delivering at least one strand (*e.g.*, a miRNA strand of the siRNA-like agent) to the cell (*e.g.*, to the RISC complex of said cell). An effective amount of a siRNA-like agent is, for example, an amount sufficient to silence
20 the natural target of the miRNA strand of the siRNA-like duplex.

miRNAs have been shown to regulate a diverse set of biological processes in metazoans (*e.g.* invertebrates, vertebrates, and plants) including cell proliferation and cell death (McManus MT *et al.*, *Semin. Cancer Biol.*, 2003; Brenneke *et al.*, *Cell*, 2003; Baehrecke EH. *Curr. Biol.*, 2003), developmental timing (Pasquinelli AE and Ruvkun G,
25 *Annu Rev. Cell. Dev.*, 2002), embryonic and post-embryonic development (Lee RC and Ambros V, *Science*, 2001; Boutla A *et al.*, *Nuc. Acids. Res.*, 2003; Carrington JC and Ambros V, *Science*, 2003; Houbaviy HB *et al.*, *Dev. Cell*, 2003; Aravin AA *et al.*, *Dev. Cell*, 2003; Yekta *et al.*, *Science*, 2004; Vaucheret *et al.*, *Genes & Dev.*, 2004), neuronal development (Johnston RJ and Hobert O, *Nature*, 2003; Kawasaki H and Taira K,
30 *Nature*, 2003; Krichevsky AM *et al.*, *RNA*, 2003; Kim J *et al.*, *PNAS*, 2004; Sempere LF *et al.*, *Genome Biology*, 2004), fat metabolism (Xu *et al.*, *Curr. Biol.*, 2003), haematopoietic cell differentiation (Chen CZ *et al.*, *Science*, 2004) and flower and leaf

development (Aukerman MJ and Sakai H, Plant Cell, 2003; Bartel B and Bartel DP, Plant Physiol., 2003; Chen X, Science, 2003; Emery JF et al., Curr. Biol., 2003; Hake et al., Curr. Biol., 2003; Hunter C and Poethig R, Curr. Opin. Genet. Dev., 2003; Kidner CA and Martienssen RA, Trends Genet., 2003; Palatnik JF, Nature, 2003; Achard P et al., Development, 2004; Zhong R et al., Plant Cell Physiol., 2004; Kidner CA et al., Nature, 2004; Mallory AC et al., Curr. Biol., 2004). miRNA disorders can result in severe disruption of these processes. miRNA disorders have been implicated in a wide variety of plant and animal diseases including cancer, neurological disorders (*e.g.* neurodegeneration), and developmental defects. For example, miRNA disorders have been correlated with nonsmall cell lung cancer (Takamizawa *et al.*, Cancer Research, 2004), colorectal cancer (Michael MZ et al., Mol. Cancer. Res., 2003), chronic lymphocytic leukemia (Calin GA et al., PNAS, 2002; Calin GA et al., PNAS, 2004), Fragile X Syndrome (Caudy A et al, Genes & Dev., 2002), spinal muscular atrophy and Waisman Syndrome (Dostie et al., RNA, 2003). Additional miRNA-silenced target gene linked with a particular miRNA disorder can be identified *in silico* using computational algorithms described in the art (Rhoades et al. Cell, 2002; Enright et al. Genome Biol., 2003; Lewis et al., Cell, 2003; Stark et al., PLOS Biol., 2003).

The choice of which strand of the siRNA-like duplex is preferentially loaded in the RISC complex may depend on the nature of the miRNA disorder. In one scenario, an miRNA disorders can result from a deficiency in the expression of a particular natural miRNA due to a mutation or chromosomal lesion (*e.g.* translocation, inversion, or deletion) in the miRNA gene or a regulatory sequence controlling its expression (*e.g.* an miRNA promoter). In another scenario, an miRNA disorders can result from an acquisition of a mutation in a target mRNA. For example, a mutation in a miRNA-silenced target gene can render the target mRNA resistant to silencing, leading to overexpression of a target mRNA that is normally silenced. Alternatively, a mutation in the miRNA-silenced target gene can render the target mRNA susceptible to miRNA-guided cleavage, leading to underexpression of a target mRNA that is normally constitutively expressed.

In another scenario, an miRNA disorder can result from the acquisition of a mutation in a natural miRNA. A mutation in the 5' end of an individual strand of an miRNA duplex can result in the loading the wrong strand in RISC. For example, when

the miRNA* strand is inappropriately loaded in RISC, the target mRNA may no longer be silenced, leading to overexpression of a target mRNA that is normally silenced and/or underexpression of a new, inappropriate target mRNA that is normally not silenced (“off-target silencing”). In yet another scenario, an miRNA disorder can result from a
 5 mutation in a gene encoding a component of the RISC complex such that miRNA-mediated gene silencing is abrogated.

Use of the design rules described *supra* can be applied to the RNA silencing agents to correct for an miRNA disorder. In particular, miRNA duplex intermediates or siRNA-like duplexes that include a miRNA strand and/or a miRNA* strand can be
 10 altered with the design rules to promote entry of the desired strand of the duplex (e.g., the miRNA strand) into a RISC complex. In one aspect of the invention, entry of the miRNA strand into the RISC complex is desired in order to treat the disorder. In another embodiment of the invention, entry of the miRNA* strand is desired in order to treat the disorder. Entry of the miRNA strand may be promoted by either lessening the base pair
 15 strength between the 5' end of the miRNA strand and the 3' end of the miRNA* strand (miRNA* 3') as compared to the base pair strength between the 3' end of the miRNA strand and the 5' end of the miRNA* strand or enhancing the base pair strength between the 3' end of the miRNA strand and the 5' end of the miRNA* strand as compared to the base pair strength between the 5' end of miRNA strand and the 3' end of the miRNA*
 20 strand. Alternatively, entry of the sense strand may be promoted by lessening the base pair strength between the 5' end of the miRNA* strand (miRNA* 5') and the 3' end of the miRNA strand (miRNA 3') as compared to the base pair strength between the 3' end of the miRNA* strand (miRNA* 3') and the 5' end of the miRNA strand (miRNA 5') or enhancing the base pair strength between the 3' end of the miRNA* strand (miRNA 3')
 25 and the 5' end of the miRNA strand (miRNA 5') as compared to the base pair strength between the 5' end of miRNA* strand (miRNA* 5') and the 3' end of the miRNA strand (miRNA 3').

4. Pharmacogenomics

30 The therapeutic agents (e.g., a RNAi agent or vector or transgene encoding same) of the invention can be administered to individuals to treat (prophylactically or therapeutically) disorders associated with aberrant or unwanted target gene activity. In

conjunction with such treatment, pharmacogenomics (*i.e.*, the study of the relationship between an individual's genotype and that individual's response to a foreign compound or drug) may be considered. Differences in metabolism of therapeutics can lead to severe toxicity or therapeutic failure by altering the relation between dose and blood concentration of the pharmacologically active drug. Thus, a physician or clinician may consider applying knowledge obtained in relevant pharmacogenomics studies in determining whether to administer a therapeutic agent as well as tailoring the dosage and/or therapeutic regimen of treatment with a therapeutic agent.

Pharmacogenomics deals with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See, for example, Eichelbaum, M. *et al.* (1996) *Clin. Exp. Pharmacol. Physiol.* 23(10-11): 983-985 and Linder, M.W. *et al.* (1997) *Clin. Chem.* 43(2):254-266. In general, two types of pharmacogenetic conditions can be differentiated. Genetic conditions transmitted as a single factor altering the way drugs act on the body (altered drug action) or genetic conditions transmitted as single factors altering the way the body acts on drugs (altered drug metabolism). These pharmacogenetic conditions can occur either as rare genetic defects or as naturally-occurring polymorphisms. For example, glucose-6-phosphate dehydrogenase deficiency (G6PD) is a common inherited enzymopathy in which the main clinical complication is haemolysis after ingestion of oxidant drugs (anti-malarials, sulfonamides, analgesics, nitrofurans) and consumption of fava beans.

One pharmacogenomics approach to identifying genes that predict drug response, known as "a genome-wide association", relies primarily on a high-resolution map of the human genome consisting of already known gene-related markers (*e.g.*, a "bi-allelic" gene marker map which consists of 60,000-100,000 polymorphic or variable sites on the human genome, each of which has two variants.) Such a high-resolution genetic map can be compared to a map of the genome of each of a statistically significant number of patients taking part in a Phase II/III drug trial to identify markers associated with a particular observed drug response or side effect. Alternatively, such a high resolution map can be generated from a combination of some ten-million known single nucleotide polymorphisms (SNPs) in the human genome. As used herein, a "SNP" is a common alteration that occurs in a single nucleotide base in a stretch of

DNA. For example, a SNP may occur once per every 1000 bases of DNA. A SNP may be involved in a disease process, however, the vast majority may not be disease-associated. Given a genetic map based on the occurrence of such SNPs, individuals can be grouped into genetic categories depending on a particular pattern of SNPs in their individual genome. In such a manner, treatment regimens can be tailored to groups of genetically similar individuals, taking into account traits that may be common among such genetically similar individuals.

Alternatively, a method termed the "candidate gene approach", can be utilized to identify genes that predict drug response. According to this method, if a gene that encodes a drug's target is known (*e.g.*, a target gene polypeptide of the present invention), all common variants of that gene can be fairly easily identified in the population and it can be determined if having one version of the gene versus another is associated with a particular drug response.

As an illustrative embodiment, the activity of drug metabolizing enzymes is a major determinant of both the intensity and duration of drug action. The discovery of genetic polymorphisms of drug metabolizing enzymes (*e.g.*, N-acetyltransferase 2 (NAT 2) and cytochrome P450 enzymes CYP2D6 and CYP2C19) has provided an explanation as to why some patients do not obtain the expected drug effects or show exaggerated drug response and serious toxicity after taking the standard and safe dose of a drug. These polymorphisms are expressed in two phenotypes in the population, the extensive metabolizer (EM) and poor metabolizer (PM). The prevalence of PM is different among different populations. For example, the gene coding for CYP2D6 is highly polymorphic and several mutations have been identified in PM, which all lead to the absence of functional CYP2D6. Poor metabolizers of CYP2D6 and CYP2C19 quite frequently experience exaggerated drug response and side effects when they receive standard doses. If a metabolite is the active therapeutic moiety, PM show no therapeutic response, as demonstrated for the analgesic effect of codeine mediated by its CYP2D6-formed metabolite morphine. The other extreme are the so called ultra-rapid metabolizers who do not respond to standard doses. Recently, the molecular basis of ultra-rapid metabolism has been identified to be due to CYP2D6 gene amplification.

Alternatively, a method termed the "gene expression profiling", can be utilized to identify genes that predict drug response. For example, the gene expression of an

animal dosed with a therapeutic agent of the present invention can give an indication whether gene pathways related to toxicity have been turned on.

Information generated from more than one of the above pharmacogenomics approaches can be used to determine appropriate dosage and treatment regimens for prophylactic or therapeutic treatment an individual. This knowledge, when applied to dosing or drug selection, can avoid adverse reactions or therapeutic failure and thus enhance therapeutic or prophylactic efficiency when treating a subject with a therapeutic agent, as described herein.

Therapeutic agents can be tested in an appropriate animal model. For example, an RNAi agent (or expression vector or transgene encoding same) as described herein can be used in an animal model to determine the efficacy, toxicity, or side effects of treatment with said agent. Alternatively, a therapeutic agent can be used in an animal model to determine the mechanism of action of such an agent. For example, an agent can be used in an animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent can be used in an animal model to determine the mechanism of action of such an agent.

VI. Pharmaceutical Compositions

The invention pertains to uses of the above-described agents for therapeutic treatments as described infra. Accordingly, the modulators of the present invention can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, antibody, or modulatory compound and a pharmaceutically acceptable carrier. As used herein the language "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, intraperitoneal, intramuscular, oral (*e.g.*, inhalation), transdermal (topical), and transmucosal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be

brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, 5 dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying 10 which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and 15 used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the 20 following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, 25 methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For 30 transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and

fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

- 5 The compounds can also be prepared in the form of suppositories (*e.g.*, with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

 In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled
10 release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova
15 Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

 It is especially advantageous to formulate oral or parenteral compositions in
20 dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are
25 dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

 Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for
30 determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio

LD50/ED50. Compounds that exhibit large therapeutic indices are preferred. Although compounds that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

5 The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the
10 method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the EC50 (*i.e.*, the concentration of the test compound which achieves a half-maximal response) as determined in cell culture. Such information can be used to more accurately determine useful doses in
15 humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

 The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

VI. Knockout and/or Knockdown Cells or Organisms

20 A further preferred use for the RNAi agents of the present invention (or vectors or transgenes encoding same) is a functional analysis to be carried out in eukaryotic cells, or eukaryotic non-human organisms, preferably mammalian cells or organisms and most preferably human cells, *e.g.* cell lines such as HeLa or 293 or rodents, *e.g.* rats and mice. By administering a suitable RNAi agent which is sufficiently complementary to a
25 target mRNA sequence to direct target-specific RNA interference, a specific knockout or knockdown phenotype can be obtained in a target cell, *e.g.* in cell culture or in a target organism.

 Thus, a further subject matter of the invention is a eukaryotic cell or a eukaryotic non-human organism exhibiting a target gene-specific knockout or knockdown
30 phenotype comprising a fully or at least partially deficient expression of at least one endogeneous target gene wherein said cell or organism is transfected with at least one

vector comprising DNA encoding an RNAi agent capable of inhibiting the expression of the target gene. It should be noted that the present invention allows a target-specific knockout or knockdown of several different endogeneous genes due to the specificity of the RNAi agent.

- 5 Gene-specific knockout or knockdown phenotypes of cells or non-human organisms, particularly of human cells or non-human mammals may be used in analytic to procedures, *e.g.* in the functional and/or phenotypical analysis of complex physiological processes such as analysis of gene expression profiles and/or proteomes. Preferably the analysis is carried out by high throughput methods using oligonucleotide
10 based chips.

- Using RNAi based knockout or knockdown technologies, the expression of an endogeneous target gene may be inhibited in a target cell or a target organism. The endogeneous gene may be complemented by an exogenous target nucleic acid coding for the target protein or a variant or mutated form of the target protein, *e.g.* a gene or a
15 DNA, which may optionally be fused to a further nucleic acid sequence encoding a detectable peptide or polypeptide, *e.g.* an affinity tag, particularly a multiple affinity tag.

- Variants or mutated forms of the target gene differ from the endogeneous target gene in that they encode a gene product which differs from the endogeneous gene product on the amino acid level by substitutions, insertions and/or deletions of single or
20 multiple amino acids. The variants or mutated forms may have the same biological activity as the endogeneous target gene. On the other hand, the variant or mutated target gene may also have a biological activity, which differs from the biological activity of the endogeneous target gene, *e.g.* a partially deleted activity, a completely deleted activity, an enhanced activity etc. The complementation may be accomplished by compressing
25 the polypeptide encoded by the endogeneous nucleic acid, *e.g.* a fusion protein comprising the target protein and the affinity tag and the double stranded RNA molecule for knocking out the endogeneous gene in the target cell. This compression may be accomplished by using a suitable expression vector expressing both the polypeptide encoded by the endogenous nucleic acid, *e.g.* the tag-modified target protein and the
30 double stranded RNA molecule or alternatively by using a combination of expression vectors. Proteins and protein complexes which are synthesized *de novo* in the target cell will contain the exogenous gene product, *e.g.*, the modified fusion protein. In order to

avoid suppression of the exogenous gene product by the RNAi agent, the nucleotide sequence encoding the exogenous nucleic acid may be altered at the DNA level (with or without causing mutations on the amino acid level) in the part of the sequence which is homologous to the RNAi agent. Alternatively, the endogeneous target gene may be
5 complemented by corresponding nucleotide sequences from other species, *e.g.* from mouse.

VII. Transgenic Organisms

Engineered RNA precursors of the invention can be expressed in transgenic animals. These animals represent a model system for the study of disorders that are
10 caused by, or exacerbated by, overexpression or underexpression (as compared to wildtype or normal) of nucleic acids (and their encoded polypeptides) targeted for destruction by the RNAi agents, *e.g.*, siRNAs and shRNAs, and for the development of therapeutic agents that modulate the expression or activity of nucleic acids or polypeptides targeted for destruction.

15 Transgenic animals can be farm animals (pigs, goats, sheep, cows, horses, rabbits, and the like), rodents (such as rats, guinea pigs, and mice), non-human primates (for example, baboons, monkeys, and chimpanzees), and domestic animals (for example, dogs and cats). Invertebrates such as *Caenorhabditis elegans* or *Drosophila* can be used as well as non-mammalian vertebrates such as fish (*e.g.*, zebrafish) or birds (*e.g.*,
20 chickens).

Engineered RNA precursors with stems of 18 to 30 nucleotides in length are preferred for use in mammals, such as mice. A transgenic founder animal can be identified based upon the presence of a transgene that encodes the new RNA precursors in its genome, and/or expression of the transgene in tissues or cells of the animals, for
25 example, using PCR or Northern analysis. Expression is confirmed by a decrease in the expression (RNA or protein) of the target sequence.

A transgenic founder animal can be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene encoding the RNA precursors can further be bred to other transgenic animals carrying other transgenes. In
30 addition, cells obtained from the transgenic founder animal or its offspring can be cultured to establish primary, secondary, or immortal cell lines containing the transgene.

1. Procedures for Making Transgenic, Non-Human Animals

A number of methods have been used to obtain transgenic, non-human animals, which are animals that have gained an additional gene by the introduction of a transgene into their cells (e.g., both the somatic and germ cells), or into an ancestor's germ line. In some cases, transgenic animals can be generated by commercial facilities (e.g., The Transgenic Drosophila Facility at Michigan State University, The Transgenic Zebrafish Core Facility at the Medical College of Georgia (Augusta, Georgia), and Xenogen Biosciences (St. Louis, MO). In general, the construct containing the transgene is supplied to the facility for generating a transgenic animal.

Methods for generating transgenic animals include introducing the transgene into the germ line of the animal. One method is by microinjection of a gene construct into the pronucleus of an early stage embryo (e.g., before the four-cell stage; Wagner et al., 1981, Proc. Natl. Acad. Sci. USA 78:5016; Brinster et al., 1985, Proc. Natl. Acad. Sci. USA 82:4438). Alternatively, the transgene can be introduced into the pronucleus by retroviral infection. A detailed procedure for producing such transgenic mice has been described (see e.g., Hogan et al., *Manipulating the Mouse Embryo*. Cold Spring Harbour Laboratory, Cold Spring Harbour, NY (1986); U.S. Patent No. 5,175,383 (1992)). This procedure has also been adapted for other animal species (e.g., Hammer et al., 1985, *Nature* 315:680; Murray et al., 1989, *Reprod. Fert. Devl.* 1:147; Pursel et al., 1987, *Vet. Immunol. Histopath.* 17:303; Rexroad et al., 1990, *J. Reprod. Fert.* 41 (suppl): 1-19; Rexroad et al., 1989, *Molec. Reprod. Devl.* 1:164; Simons et al., 1988, *BioTechnology* 6:179; Vize et al., 1988, *J. Cell. Sci.* 90:295; and Wagner, 1989, *J. Cell. Biochem.* 13B (suppl): 164).

In brief, the procedure involves introducing the transgene into an animal by microinjecting the construct into the pronuclei of the fertilized mammalian egg(s) to cause one or more copies of the transgene to be retained in the cells of the developing mammal(s). Following introduction of the transgene construct into the fertilized egg, the egg may be incubated in vitro for varying amounts of time, or reimplanted in a surrogate host, or both. One common method is to incubate the embryos in vitro for about 1-7 days, depending on the species, and then reimplant them into the surrogate host. The presence of the transgene in the progeny of the transgenically manipulated embryos can be tested by Southern blot analysis of a segment of tissue.

Another method for producing germ-line transgenic animals is through the use of embryonic stem (ES) cells. The gene construct can be introduced into embryonic stem cells by homologous recombination (Thomas et al., 1987, Cell 51:503; Capecchi, Science 1989, 244:1288; Joyner et al., 1989, Nature 338:153) in a transcriptionally
5 active region of the genome. A suitable construct can also be introduced into embryonic stem cells by DNA-mediated transfection, such as by electroporation (Ausubel et al., Current Protocols in Molecular Biology, John Wiley & Sons, 1987). Detailed procedures for culturing embryonic stem cells (e.g., ES-D3@ ATCC# CCL-1934, ES-E14TG2a, ATCC# CCL-1821, American Type Culture Collection, Rockville, AM) and
10 methods of making transgenic animals from embryonic stem cells can be found in Teratocarcinomas and Embryonic Stem Cells, A Practical Approach, ed. E.J. Robertson (IRL Press, 1987). In brief, the ES cells are obtained from pre-implantation embryos cultured in vitro (Evans et al., 1981, Nature 292:154-156). Transgenes can be efficiently introduced into ES cells by DNA transfection or by retrovirus-mediated transduction.
15 The resulting transformed ES cells can thereafter be combined with blastocysts from a non-human animal. The ES cells colonize the embryo and contribute to the germ line of the resulting chimeric animal.

In the above methods, the transgene can be introduced as a linear construct, a circular plasmid, or a viral vector, which can be incorporated and inherited as a
20 transgene integrated into the host genome. The transgene can also be constructed to permit it to be inherited as an extrachromosomal plasmid (Gassmann et al., 1995, Proc. Natl. Acad. Sci. USA 92:1292). A plasmid is a DNA molecule that can replicate autonomously in a host.

The transgenic, non-human animals can also be obtained by infecting or
25 transfecting cells either in vivo (e.g., direct injection), ex vivo (e.g., infecting the cells outside the host and later reimplanting), or in vitro (e.g., infecting the cells outside host), for example, with a recombinant viral vector carrying a gene encoding the engineered RNA precursors. Examples of suitable viral vectors include recombinant retroviral vectors (Valerio et al., 1989, Gene 84:419; Scharfinan et al., 1991, Proc. Natl. Acad. Sci.
30 USA 88:462; Miller and Buttimore, 1986, Mol. Cell. Biol. 6:2895), recombinant adenoviral vectors (Freidman et al., 1986, Mol. Cell. Biol. 6:3791; Levrero et al., 1991, Gene 101: 195), and recombinant Herpes simplex viral vectors (Fink et al., 1992,

Human Gene Therapy 3:11). Such methods are also useful for introducing constructs into cells for uses other than generation of transgenic animals.

Other approaches include insertion of transgenes encoding the new engineered RNA precursors into viral vectors including recombinant adenovirus, adenoassociated virus, and herpes simplex virus-1, or recombinant bacterial or eukaryotic plasmids. Viral vectors transfect cells directly. Other approaches include delivering the transgenes, in the form of plasmid DNA, with the help of, for example, cationic liposomes (lipofectin) or derivatized (e.g. antibody conjugated) polylysine conjugates, gramicidin S, artificial viral envelopes, or other such intracellular carriers, as well as direct injection of the transgene construct or CaPO_4 precipitation carried out *in vivo*. Such methods can also be used *in vitro* to introduce constructs into cells for uses other than generation of transgenic animals.

Retrovirus vectors and adeno-associated virus vectors can be used as a recombinant gene delivery system for the transfer of exogenous genes *in vivo* or *in vitro*. These vectors provide efficient delivery of genes into cells, and the transferred nucleic acids are stably integrated into the chromosomal DNA of the host. The development of specialized cell lines (termed "packaging cells") which produce only replication-defective retroviruses has increased the utility of retroviruses for gene therapy, and defective retroviruses are characterized for use in gene transfer for gene therapy purposes (for a review see Miller, 1990, Blood 76:271). A replication defective retrovirus can be packaged into virions which can be used to infect a target cell through the use of a helper virus by standard techniques. Protocols for producing recombinant retroviruses and for infecting cells *in vitro* or *in vivo* with such viruses can be found in Current Protocols in Molecular Biology, Ausubel, F.M. et al., (eds.) Greene Publishing Associates, (1989), Sections 9.9.14 and other standard laboratory manuals.

Examples of suitable retroviruses include pLJ, pZIP, pWE and pEM which are known to those skilled in the art. Examples of suitable packaging virus lines for preparing both ecotropic and amphotropic retroviral systems include Psi-Crip, PsiCre, Psi-2 and Psi-Am. Retroviruses have been used to introduce a variety of genes into many different cell types, including epithelial cells, *in vitro* and/or *in vivo* (see for example Egli et al., 1985, Science 230:1395-1398; Danos and Mulligan, 1988, Proc. Natl. Acad. Sci. USA 85:6460-6464; Wilson et al., 1988, Proc. Natl. Acad. Sci. USA

85:3014-3018; Armentano et al., 1990, Proc. Natl. Acad. Sci. USA 87:6141-6145; Huber et al., 1991, Proc. Natl. Acad. Sci. USA 88:8039-8043; Ferry et al., 1991, Proc. Natl. Acad. Sci. USA 88:8377-8381; Chowdhury et al., 1991, Science 254:1802-1805; van Beusechem. et al., 1992, Proc. Natl. Acad. Sci. USA 89:7640-19 ; Kay et al., 1992, 5 Human Gene Therapy 3:641-647; Dai et al., 1992, Proc. Natl. Acad. Sci. USA 89:10892-10895; Hwu et al., 1993, J. Immunol. 150:4104-4115; U.S. Patent No. 4,868,116; U.S. Patent No. 4,980,286; PCT Application WO 89/07136; PCT Application WO 89/02468; PCT Application WO 89/05345; and PCT Application WO 92/07573).

In another example, recombinant retroviral vectors capable of transducing and 10 expressing genes inserted into the genome of a cell can be produced by transfecting the recombinant retroviral genome into suitable packaging cell lines such as PA317 and Psi-CRIP (Comette et al., 1991, Human Gene Therapy 2:5-10; Cone et al., 1984, Proc. Natl. Acad. Sci. USA 81:6349). Recombinant adenoviral vectors can be used to infect a wide variety of cells and tissues in susceptible hosts (e.g., rat, hamster, dog, and chimpanzee) 15 (Hsu et al., 1992, J. Infectious Disease, 166:769), and also have the advantage of not requiring mitotically active cells for infection. Another viral gene delivery system useful in the present invention also utilizes adenovirus-derived vectors. The genome of an adenovirus can be manipulated such that it encodes and expresses a gene product of interest but is inactivated in terms of its ability to replicate in a normal lytic viral life 20 cycle. See, for example, Berkner et al. (1988, BioTechniques 6:616), Rosenfeld et al. (1991, Science 252:431-434), and Rosenfeld et al. (1992, Cell 68:143-155). Suitable adenoviral vectors derived from the adenovirus strain Ad type 5 d1324 or other strains of adenovirus (e.g., Ad2, AO, Ad7 etc.) are known to those skilled in the art. Recombinant adenoviruses can be advantageous in certain circumstances in that they are 25 not capable of infecting nondividing cells and can be used to infect a wide variety of cell types, including epithelial cells (Rosenfeld et al., 1992, cited supra). Furthermore, the virus particle is relatively stable and amenable to purification and concentration, and as above, can be modified to affect the spectrum of infectivity. Additionally, introduced adenoviral DNA (and foreign DNA contained therein) is not integrated into the genome 30 of a host cell but remains episomal, thereby avoiding potential problems that can occur as a result of insertional mutagenesis *in situ* where introduced DNA becomes integrated into the host genome (e.g., retroviral DNA). Moreover, the carrying capacity of the

adenoviral genome for foreign DNA is large (up to 8 kilobases) relative to other gene delivery vectors (Berkner et al. cited supra; Haj-Ahmand and Graham, 1986, J. Virol. 57:267).

Yet another viral vector system useful for delivery of the subject transgenes is the adeno-associated virus (AAV). Adeno-associated virus is a naturally occurring defective virus that requires another virus, such as an adenovirus or a herpes virus, as a helper virus for efficient replication and a productive life cycle. For a review, see Muzyczka et al. (1992, Curr. Topics in Micro. and Immunol. 158:97-129). It is also one of the few viruses that may integrate its DNA into non-dividing cells, and exhibits a high frequency of stable integration (see for example Flotte et al. (1992, Am. J. Respir. Cell. Mol. Biol. 7:349-356; Samulski et al., 1989, J. Virol. 63:3822-3828; and McLaughlin et al. (1989, J. Virol. 62:1963-1973). Vectors containing as little as 300 base pairs of AAV can be packaged and can integrate. Space for exogenous DNA is limited to about 4.5 kb. An AAV vector such as that described in Tratschin et al. (1985) Mol. Cell. Biol. 5:3251-3260 can be used to introduce DNA into cells. A variety of nucleic acids have been introduced into different cell types using AAV vectors (see for example Hennonat et al. (1984) Proc. Nat. Acad. Sci. USA 81:6466-6470; Tratschin et al. (1985) Mol. Cell. Biol. 4:2072-2081; Wondisford et al. (1988) Mol. Endocrinol. 2:32-39; Tratschin et al. (1984) J. Virol. 51:611-619; and Flotte et al. (1993) J. Biol. Chem. 268:3781-3790).

In addition to viral transfer methods, such as those illustrated above, non-viral methods can also be employed to cause expression of an shRNA or engineered RNA precursor of the invention in the tissue of an animal. Most non-viral methods of gene transfer rely on non-viral mechanisms used by mammalian cells for the uptake and intracellular transport of macromolecules. In preferred embodiments, non-viral gene delivery systems of the present invention rely on endocytic pathways for the uptake of the subject gene of the invention by the targeted cell. Exemplary gene delivery systems of this type include liposomal derived systems, poly-lysine conjugates, and artificial viral envelopes. Other embodiments include plasmid injection systems such as are described in Meuli et al., (2001) J. Invest. Dermatol., 116(1):131-135; Cohen et al., (2000) Gene Ther., 7(22):1896-905; and Tam et al., (2000) Gene Ther., 7(21):1867-74.

In a representative embodiment, a gene encoding an shRNA or engineered RNA precursor of the invention can be entrapped in liposomes bearing positive charges on

their surface (e.g., lipofectins) and (optionally) which are tagged with antibodies against cell surface antigens of the target tissue (Mizuno et al., (1992) No Shinkei Geka, 20:547-55 1; PCT publication WO91/06309; Japanese patent application 10473 8 1; and European patent publication EP-A-43 075).

5 Animals harboring the transgene can be identified by detecting the presence of the transgene in genomic DNA (e.g., using Southern analysis). In addition, expression of the shRNA or engineered RNA precursor can be detected directly (e.g., by Northern analysis). Expression of the transgene can also be confirmed by detecting a decrease in the amount of protein corresponding to the targeted sequence. When the transgene is
10 under the control of an inducible or developmentally regulated promoter, expression of the target protein is decreased when the transgene is induced or at the developmental stage when the transgene is expressed, respectively.

2. Clones of Transgenic Animals

15 Clones of the non-human transgenic animals described herein can be produced according to the methods described in Wilmut et al. ((1997) Nature, 385:810-813) and PCT publication Nos. WO 97/07668 and WO 97/07669. In brief, a cell, e.g., a somatic cell from the transgenic animal, can be isolated and induced to exit the growth cycle and enter the G0 phase to become quiescent. The quiescent cell can then be fused, e.g.,
20 through the use of electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops into a morula or blastocyte and is then transferred to a pseudopregnant female foster animal. Offspring borne of this female foster animal will be clones of the animal from which the cell, e.g., the somatic cell, was isolated.

25 Once the transgenic animal is produced, cells of the transgenic animal and cells from a control animal are screened to determine the presence of an RNA precursor nucleic acid sequence, e.g., using polymerase chain reaction (PCR). Alternatively, the cells can be screened to determine if the RNA precursor is expressed (e.g., by standard procedures such as Northern blot analysis or reverse transcriptase-polymerase chain
30 reaction (RT-PCR); Sambrook et al., Molecular Cloning - A Laboratory Manual, (Cold Spring Harbor Laboratory, 1989)).

The transgenic animals of the present invention can be homozygous or heterozygous, and one of the benefits of the invention is that the target mRNA is effectively degraded even in heterozygotes. The present invention provides for transgenic animals that carry a transgene of the invention in all their cells, as well as
5 animals that carry a transgene in some, but not all of their cells. That is, the invention provides for mosaic animals. The transgene can be integrated as a single transgene or in concatamers, e.g., head-to-head tandems or head-to-tail tandems.

For a review of techniques that can be used to generate and assess transgenic animals, skilled artisans can consult Gordon (IwL Rev. Cytol 11 5:171-229, 1989), and
10 may obtain additional guidance from, for example: Hogan et al. "Manipulating the Mouse Embryo" (Cold Spring Harbor Press, Cold Spring Harbor, NY, 1986; Krimpenfort et al., BiolTechnology 9:86, 1991; Palmiter et al., Cell 41:343, 1985; Kraemer et al., "Genetic Manipulation of the Early Mammalian Embryo," Cold Spring Harbor Press, Cold Spring Harbor, NY, 1985; Hammer et al., Nature 315:680, 1985;
15 Purcel et al., Science, 244:1281, 1986; Wagner et al., U.S. Patent No. 5,175,385; and Krimpenfort et al., U.S. Patent No. 5,175,384.

3. Transgenic Plants

Among the eukaryotic organisms featured in the invention are plants containing an exogenous nucleic acid that encodes an engineered RNA precursor of the invention.

20 Accordingly, a method according to the invention comprises making a plant having a nucleic acid molecule or construct, e.g., a transgene, described herein. Techniques for introducing exogenous nucleic acids into monocotyledonous and dicotyledonous plants are known in the art, and include, without limitation, Agrobacterium-mediated transformation, viral vector-mediated transformation,
25 electroporation and particle gun transformation, see, e.g., U.S. Patents Nos. 5,204,253 and 6,013,863. If a cell or tissue culture is used as the recipient tissue for transformation, plants can be regenerated from transformed cultures by techniques known to those skilled in the art. Transgenic plants can be entered into a breeding program, e.g., to introduce a nucleic acid encoding a polypeptide into other lines, to transfer the nucleic
30 acid to other species or for further selection of other desirable traits. Alternatively, transgenic plants can be propagated vegetatively for those species amenable to such techniques. Progeny includes descendants of a particular plant or plant line. Progeny of

a plant include seeds formed on F1, F2, F3, and subsequent generation plants, or seeds formed on BQ, BC2, BC3, and subsequent generation plants. Seeds produced by a transgenic plant can be grown and then selfed (or outcrossed and selfed) to obtain seeds homozygous for the nucleic acid encoding a novel polypeptide.

5 A suitable group of plants with which to practice the invention include dicots, such as safflower, alfalfa, soybean, rapeseed (high erucic acid and canola), or sunflower. Also suitable are monocots such as corn, wheat, rye, barley, oat, rice, millet, amaranth or sorghum. Also suitable are vegetable crops or root crops such as potato, broccoli, peas, sweet corn, popcorn, tomato, beans (including kidney beans, lima beans, dry beans,
10 green beans) and the like. Also suitable are fruit crops such as peach, pear, apple, cherry, orange, lemon, grapefruit, plum, mango and palm. Thus, the invention has use over a broad range of plants, including species from the genera *Anacardium*, *Arachis*, *Asparagus*, *Atropa*, *Avena*, *Brassica*, *Citrus*, *Citrullus*, *Capsicum*, *Carthamus*, *Cocos*, *Coffea*, *Cucumis*, *Cucurbita*, *Daucus*, *Elaeis*, *Fragaria*, *Glycine*, *Gossypium*, *Helianthus*,
15 *Heterocallis*, *Hordeum*, *Hyoscyamus*, *Lactuca*, *Linum*, *Lolium*, *Lupinus*, *Lycopersicon*, *Malus*, *Manihot*, *Majorana*, *Medicago*, *Nicotiana*, *Olea*, *Oryza*, *Panicum*, *Pannisetum*, *Persea*, *Phaseolus*, *Pistachia*, *Pisum*, *Pyrus*, *Prunus*, *Raphanus*, *Ricinus*, *Secale*, *Senecio*, *Sinapis*, *Solanum*, *Sorghum*, *Theobromus*, *Trigonella*, *Triticum*, *Vicia*, *Vitis*, *Vigna* and *Zea*.

20 The skilled artisan will appreciate that the enumerated organisms are also useful for practicing other aspects of the invention, e.g., as host cells, as described *supra*.

 The nucleic acid molecules of the invention can be expressed in plants in a cell- or tissue-specific manner according to the regulatory elements chosen to include in a particular nucleic acid construct present in the plant. Suitable cells, tissues, and organs
25 in which to express a chimeric polypeptide of the invention include, without limitation, egg cell, central cell, synergid cell, zygote, ovule primordia, nucellus, integuments, endothelium, female gametophyte cells, embryo, axis, cotyledons, suspensor, endosperm, seed coat, ground meristem, vascular bundle, cambium, phloem, cortex, shoot or root apical meristems, lateral shoot or root meristems, floral meristem, leaf
30 primordia, leaf mesophyll cells, and leaf epidermal cells, e.g., epidermal cells involved in forming the cuticular layer. Also suitable are cells and tissues grown in liquid media or on semi-solid media.

4. Transgenic Fungi

Other eukaryotic organisms featured in the invention are fungi containing an exogenous nucleic acid molecule that encodes an engineered RNA precursor of the invention. Accordingly, a method according to the invention comprises introducing a
5 nucleic acid molecule or construct as described herein into a fungus. Techniques for introducing exogenous nucleic acids into many fungi are known in the art, see, e.g., U.S. Patents Nos. 5,252,726 and 5,070,020. Transformed fungi can be cultured by techniques known to those skilled in the art. Such fungi can be used to introduce a nucleic acid encoding a polypeptide into other fungal strains, to transfer the nucleic acid to other
10 species or for further selection of other desirable traits.

A suitable group of fungi with which to practice the invention include fission yeast and budding yeast, such as *Saccharomyces cerevisiae*, *S. pombe*, *S. carlsbergensis* and *Candida albicans*. Filamentous fungi such as *Aspergillus* spp. and *Penicillium* spp. are also useful.

15

VIII. Functional Genomics and/or Proteomics

Preferred applications for the cell or organism of the invention is the analysis of gene expression profiles and/or proteomes. In an especially preferred embodiment an analysis of a variant or mutant form of one or several target proteins is carried out,
20 wherein said variant or mutant forms are reintroduced into the cell or organism by an exogenous target nucleic acid as described above. The combination of knockout of an endogenous gene by application of a RNA silencing agent of the invention and rescue of the knockout by using mutated, e.g. partially deleted exogenous target has advantages compared to the use of a knockout cell. Further, this method is particularly suitable for
25 identifying functional domains of the targeted protein. In a further preferred embodiment a comparison, e.g. of gene expression profiles and/or proteomes and/or phenotypic characteristics of at least two cells or organisms is carried out. These organisms are selected from: (i) a control cell or control organism without target gene inhibition, (ii) a cell or organism with target gene inhibition and (iii) a cell or organism
30 with target gene inhibition plus target gene complementation by an exogenous target nucleic acid.

Furthermore, the RNA knockout complementation method may be used for is preparative purposes, *e.g.* for the affinity purification of proteins or protein complexes from eukaryotic cells, particularly mammalian cells and more particularly human cells. In this embodiment of the invention, the exogenous target nucleic acid preferably codes
5 for a target protein which is fused to an affinity tag. This method is suitable for functional proteome analysis in mammalian cells, particularly human cells.

Another utility of the present invention could be a method of identifying gene function in a cell or organism comprising the use of an RNA silencing agent of the invention (*e.g.* RNAi agent or agent causing translational repression) to inhibit the
10 activity of a target gene of previously unknown function. Instead of the time consuming and laborious isolation of mutants by traditional genetic screening, functional genomics would envision determining the function of uncharacterized genes by employing the RNA silencing agents of the invention to reduce the amount and/or alter the timing of target gene activity.

15 The invention could be used in determining potential pharmaceuticals, potential targets for pharmaceuticals, understanding normal and pathological events associated with development, determining signaling pathways responsible for postnatal development/aging, and the like. The increasing speed of acquiring nucleotide sequence information from genomic and expressed gene sources, including total genome
20 sequences for model organisms (*e.g.* rat, mouse, zebrafish, yeast, *Arabidopsis thaliana*, *D. melanogaster*, and *C. elegans*), can be coupled with the invention to determine gene function in that organisms, a related organism (*e.g.*, a parasitic fly or nematode), or an organism containing a homolog of the gene of interest (*e.g.* human). The preference of different organisms to use particular codons, searching sequence databases for related
25 gene products, correlating the linkage map of genetic traits with the physical map from which the nucleotide sequences are derived, and artificial intelligence methods may be used to define putative open reading frames from the nucleotide sequences acquired in such sequencing projects. A simple assay would be to inhibit gene expression according to the partial sequence available from an expressed sequence tag (EST). Functional
30 alterations in growth, development, metabolism, disease resistance, or other biological processes would be indicative of the normal role of the EST's gene product.

The ease with which RNA can be introduced into an intact cell or model organism containing the target gene allows the RNA silencing agents of the present invention to be used in high throughput screening (HTS). Solutions containing RNA silencing agents (e.g. RNAi agents or translational repression agents) that are capable of inhibiting the different expressed genes can be placed into individual wells positioned on a microtiter plate as an ordered array, and intact cells organisms in each well can be assayed for any changes or modifications in behavior or development due to inhibition of target gene activity. The amplified RNA can be fed directly to, injected into, the cell/organism containing the target gene. Alternatively, the RNA silencing agent (e.g. RNAi agent or translational repression agent) can be produced from a vector, as described herein. Vectors can be injected into the cell/organism containing the target gene. The function of the target gene can be assayed from the effects it has on the cell/organism when gene activity is inhibited. This screening could be amenable to small subjects that can be processed in large number, for example: arabidopsis, bacteria, drosophila, fungi, nematodes, viruses, zebrafish, and tissue culture cells derived from mammals. A nematode or other organism that produces a colorimetric, fluorogenic, or luminescent signal in response to a regulated promoter (e.g., transfected with a reporter gene construct) can be assayed in an HTS format.

The present invention may be useful in allowing the inhibition of essential genes. Such genes may be required for cell or organism viability at only particular stages of development or cellular compartments. The functional equivalent of conditional mutations may be produced by inhibiting activity of the target gene when or where it is not required for viability. The invention allows addition of the RNA silencing agent (e.g. RNAi or translational repression agent) at specific times of development and locations in the organism without introducing permanent mutations into the target genome.

siRNA-like silencing agents of the present invention can be used in method for identifying the natural target (e.g., the natural mRNA target) of a miRNA in a cell or organism. An exemplary method comprises contacting a plurality or panel of candidate target genes (e.g., target mRNA sequences), for example, several hundred natural mRNAs represented on a microwell plate). mRNAs can be separated into individual wells positioned on a microtiter plate as an ordered array. The miRNA can then be

assessed for its ability to bind to, direct cleavage of, recruit RISC components, or any other indicator of RNA silencing (*e.g.*, by RNAi or translational repression) using methods described herein or routine in the art. In other embodiments, siRNA-like duplexes or miRNAs can be arrayed on plates for testing with a specific target gene or mRNA. In preferred embodiments, the miRNAs are used at low concentrations (*e.g.*, picomolar concentration) to prevent non-specific silencing of the target.

When used to analyse gene expression profiles and/or proteomes, to identify or inhibit gene function, or to validate pharmaceutic targets, the RNA silencing agents of the invention offer improvements over existing RNA silencing agents. The RNA silencing agents of the invention have several-fold improvement in potency (*e.g.*, can be used at picomolar concentrations vs. the nanomolar or micromolar concentrations required when using un-modified, *e.g.*, symmetrical silencing agents) so that lower and more economical amounts of these reagents can be used for the methods described *supra*. Alternatively, the use of the RNA silencing agents of the invention may obviate the need for costly chemical modifications.

IX. Screening Assays

The methods of the invention are also suitable for use in methods to identify and/or characterize potential pharmacological agents, *e.g.* identifying new pharmacological agents from a collection of test substances and/or characterizing mechanisms of action and/or side effects of known pharmacological agents. In one exemplary embodiment, a library of pharmaceutical agents can be screened to identify a pharmaceutical agent which suppresses or “rescues” the gene silencing phenotype in a cell or organism caused by previous administration of the RNA silencing agent.

Thus, the present invention also relates to a system for identifying and/or characterizing pharmacological agents acting on at least one target protein comprising: (a) a eukaryotic cell or a eukaryotic non-human organism capable of expressing at least one endogenous target gene coding for said so target protein, (b) at least one silencing agent (*e.g.* RNAi agent) molecule capable of inhibiting the expression of said at least one endogenous target gene, and (c) a test substance or a collection of test substances wherein pharmacological properties of said test substance or said collection are to be

identified and/or characterized. Further, the system as described above preferably comprises: (d) at least one exogenous target nucleic acid coding for the target protein or a variant or mutated form of the target protein wherein said exogenous target nucleic acid differs from the endogeneous target gene on the nucleic acid level such that the expression of the exogenous target nucleic acid is substantially less inhibited by the silencing agent (*e.g.* RNAi agent) than the expression of the endogeneous target gene.

The test compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the 'one-bead one-compound' library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam, K.S. (1997) *Anticancer Drug Des.* 12:145).

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt *et al.* (1993) *Proc. Natl. Acad. Sci. U.S.A.* 90:6909; Erb *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:11422; Zuckermann *et al.* (1994). *J. Med. Chem.* 37:2678; Cho *et al.* (1993) *Science* 261:1303; Carrell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2059; Carell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2061; and in Gallop *et al.* (1994) *J. Med. Chem.* 37:1233.

Libraries of compounds may be presented in solution (*e.g.*, Houghten (1992) *Biotechniques* 13:412-421), or on beads (Lam (1991) *Nature* 354:82-84), chips (Fodor (1993) *Nature* 364:555-556), bacteria (Ladner USP 5,223,409), spores (Ladner USP '409), plasmids (Cull *et al.* (1992) *Proc Natl Acad Sci USA* 89:1865-1869) or on phage (Scott and Smith (1990) *Science* 249:386-390); (Devlin (1990) *Science* 249:404-406); (Cwirla *et al.* (1990) *Proc. Natl. Acad. Sci.* 87:6378-6382); (Felici (1991) *J. Mol. Biol.* 222:301-310); (Ladner *supra.*)).

In a preferred embodiment, the library is a natural product library, *e.g.*, a library produced by a bacterial, fungal, or yeast culture. In another preferred embodiment, the library is a synthetic compound library.

When used in the screening assays described herein, the RNA silencing agents of the invention offer improvements in potency over existing RNA silencing agents. For example, the RNA silencing agents of the invention have several-fold improvement in potency (picomolar vs. micromolar) such that lower and more economical amounts of these reagents can be used for the methods described *supra*. Alternatively, the use of the RNA silencing agents of the invention may obviate the need for costly chemical modifications.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are incorporated herein by reference.

EXAMPLES

Example I: Functionally asymmetric siRNA duplexes

To assess quantitatively if the two strands of an siRNA duplex are equally competent to direct RNAi, the individual rates of sense and anti-sense target cleavage for an siRNA duplex directed against the firefly luciferase mRNA were examined (Figure 1A). The relevant portions of the sense and anti-sense target RNA sequences are shown in Figure 1A and the siRNA sequence in Figure 1B. This siRNA duplex effectively silences firefly luciferase expression in culture human HeLa cells. Using a *Drosophila* embryo-derived *in vitro* RNAi reaction, a significant difference in the rate of target cleavage for the two siRNA strands was found; the anti-sense siRNA strand directed more efficient RNAi against a sense RNA target than the sense siRNA strand for an anti-sense target (Figure 1B). (Anti-sense siRNA strands and sense target RNAs are always shown in black, and sense siRNAs and anti-sense targets, in grey). Control experiments showed that using siRNA duplexes with 5' phosphates did not alter this result (data not shown), indicating that different rates of phosphorylation for the two strands is not the cause for the observed asymmetry. Surprisingly, the two strands of the luciferase duplex siRNA duplex, used individually as 5' phosphorylated single strands, had identical rates of target cleavage (Figure 1C). RNAi directed by single-stranded siRNA is roughly 10-fold less efficient than that triggered by siRNA duplexes, reflecting the ~ 100-fold lower stability of single-stranded siRNAs *in vitro* and *in vivo* (Schwarz et al., 2002). The difference in the rate of cleavage directed by the sense and anti-sense strands when the reaction was programmed with an siRNA duplex is unlikely to reflect a difference in the inherent susceptibility of the two targets to RNAi. Instead, the observation that the same two siRNA strands are equally effective as single-strands, but show dramatically different activities when paired with each other, indicates that the asymmetry in their function is established at a step in the RNAi pathway prior to the encounter of the programmed RISC with its corresponding RNA target.

Example II: Differential RISC assembly accounts for siRNA strand functional asymmetry

To identify the source of asymmetry in the function of this siRNA duplex, the unwinding of the two siRNA strands when the duplex was incubated in a standard *in vitro* RNAi reaction was measured. This assay was shown previously to determine accurately the fraction of siRNA that is unwound in an ATP-dependent step in the RNAi pathway; no functional RISC is assembled in the absence of ATP (Nykänen et al., 2001). Previous studies show that siRNA unwinding correlates with capacity of an siRNA to function in target cleavage (Nykänen et al., 2001; Martinez et al., 2002), demonstrating that siRNA duplex unwinding is required to assemble a RISC competent to base pair with its target RNA. Here, the accumulation of single stranded siRNA from the luciferase siRNA duplex after 1 hour incubation in an *in vitro* RNAi reaction in the absence of target RNA was measured. After one hour of incubation with *Drosophila* embryo lysate in a standard RNAi reaction, 22% of the anti-sense strand of the luciferase siRNA was converted to single-strand (Figure 1D; 'siRNA B' solid black bar). Remarkably, a corresponding amount of single-stranded sense siRNA was not detected. Instead, only 3% of the sense strand accumulated as single-stranded siRNA (Figure 1D; 'siRNA B' solid grey bar). In control experiments, no single-stranded RNA was detected without incubation in lysate (not shown), demonstrating that the siRNA was entirely double-stranded at the beginning of the reaction. Since the production of single-stranded anti-sense siRNA must be accompanied by an equal amount of single-stranded sense siRNA, the missing sense-strand must have been destroyed after unwinding.

To establish that the observed asymmetry in the accumulation of the two single-strands was not an artifact of our unwinding assay, an independent method for measuring the fraction of siRNA present as single-strands in protein-RNA complexes was. In this assay, double-stranded siRNA was incubated with *Drosophila* embryo lysate in a standard RNAi reaction for 1 h, then a 31 nt 2'-O-methyl RNA oligonucleotide containing a 21 nt sequence complementary to the radiolabeled siRNA strand was added. 2'-O-methyl oligonucleotides are not cleaved by the RNAi machinery, but can bind stably to complementary siRNA within the RISC (Martin Simard, GH, Craig Mello, and PDZ, manuscript in preparation). To allow recovery of RISC, the 2'-O-methyl oligonucleotide was tethered to a magnetic bead via a biotin-streptavidin linkage.

After washing away unbound RNA and protein, the amount of radioactive siRNA bound to the bead was measured. The assay was performed with separate siRNA duplexes in which either the sense or the anti-sense strand was 5'-³²P-radiolabeled. Capture of ³²P-siRNA was observed when the 2'-*O*-methyl oligonucleotide contained a 21-nt region
5 complementary to the radiolabeled siRNA strand, but not when an unrelated oligonucleotide was used. The assay captures all RISC activity directed by the siRNA strand complementary to the tethered oligonucleotide, demonstrating that it measures siRNA present in the lysate as single-strand complexed with RISC proteins. This assay recapitulates the results of the unwinding assay described above: for the siRNA in Figure
10 1D; 'si RNA B' open bars, nearly ten-fold more anti-sense siRNA was detected than sense strand. An explanation for these results is that the two strands of this siRNA duplex are differentially loaded into the RISC, and that single-stranded siRNA not assembled into RISC is degraded. Functional asymmetry occurred only when the trigger siRNA was double-stranded, not when the two siRNA strands were tested individually
15 (Fig. 1B and 1C). Thus, asymmetric assembly of RISC was a feature of the siRNA duplex, rather than of either the sequences of the individual siRNA strands or the accessibility of the targeted sites to cleavage.

Example III: Base-pairing at the 5' end of the siRNA strand gates RISC assembly

20 The finding that the two siRNA strands can have different capacities to form RISC when paired in a duplex indicates that some feature of the 19 base-pairs of the duplex determines functional asymmetry. These base-pairs must be disrupted to produce RISC (Nykanen et al., 2001), which contains single-stranded siRNA (Martinez et al., 2002). The siRNAs used in Figure 1B were examined for base-pairing features that
25 might distinguish the two siRNA strands. For the siRNA in Figure 1B, the 5' end of the anti-sense siRNA strand begins with U and is thus paired to the sense siRNA strand by an A:U base pair (2 hydrogen bonds). In contrast, the 5' nucleotide of the sense siRNA strand is linked to the anti-sense strand by a C:G base pair (3 hydrogen bonds). The sense siRNA strand forms 8-10-fold less RISC and guides cleavage of its RNA target at
30 a correspondingly slower rate than the anti-sense strand. A working hypothesis to explain the observed functional asymmetry is that the siRNA strand whose 5' end is more weakly bound to the complementary strand more readily incorporates into RISC.

In this view, the relative base-pairing strengths of the 5' ends of the two siRNA strands would determine their relative extents of RISC formation.

As an initial test of this idea, the 5' nucleotide of the siRNA sense strand was changed from C to U (Figure 1E). This changed the base pair formed between the 5' most nucleotide of the sense strand and position 19 of the anti-sense strand from the Watson-Crick base pair C:G to the weaker, less stable wobble pair U:G, while leaving the anti-sense strand of the siRNA unaltered. Remarkably, the change of this single nucleotide not only enhanced the rate of cleavage directed by the sense strand, but virtually eliminated the ability of the anti-sense strand to direct RNAi (Figure 1E).

To determine the basis for the reversed functional asymmetry for the siRNA in Figure 1E, the amount of each strand that was single stranded after incubation of the siRNA duplex in *Drosophila* embryo lysates was determined. After 1h, nearly 30% of the sense siRNA strand was converted to single stranded, but no single-stranded anti-sense strand was detected (Figure 1D; 'siRNA E'). Therefore, the simplest explanation for the asymmetric function of this siRNA is that the sense strand, but not the anti-sense, of this siRNA duplex was incorporated into RISC. Thus, a single nucleotide mutation in the sense siRNA strand of the siRNA in Figure 1B completely reversed the relative abilities of the two strands to assemble in the enzyme complex that directs RNAi.

The stability of the initial five base pairs of the siRNA strands was calculated in Figure 1 using the nearest-neighbor method and the mfold algorithm (D.H. Mathews, 1999; Zuker, 2003). The 5' end of the sense siRNA strand in Figure 1E, but not that in 1B, is predicted to exist as an equilibrium of two conformers of nearly equal energy (Figure 10). In one conformer, the 5' nucleotide of the sense strand is bound to the anti-sense strand by a U:G wobble pair, whereas in the other conformer the 5' end of this siRNA strand is unpaired. The analysis suggests that RISC assembly favors the siRNA strand whose 5' end has a greater propensity to fray.

To test this hypothesis further, the strand-specific rates of cleavage of sense and anti-sense human *Cu, Zn superoxide dismutase-1 (sod1)* RNA targets (Figure 3A) triggered by the siRNA duplex shown in Figure 3B were examined. Given that the 5' ends of both siRNA strands of this duplex are in G:C base pairs, it was anticipated that this duplex would not display pronounced target cleavage asymmetry. As shown in Figure 3B, the two strands are similar in their rates of target cleavage, although the rate

of anti-sense cleavage directed by the sense-strand is clearly faster than the rate of sense-target cleavage guided by the anti-sense strand. This small difference in rate is likely explained by the sense-strand forming 20 base pairs with its target RNA, whereas the anti-sense strand can form only 19, consistent with previous reports that the penultimate position of an siRNA makes a small contribution to its efficacy (Elbashir et al., 2001b). Next, the C at position 19 of the sense strand was changed to A, causing the anti-sense strand to begin with an unpaired nucleotide. This change, which was made to the sense-strand of the siRNA, caused the rate of target cleavage guided by the *anti-sense* siRNA strand to be dramatically enhanced and the sense strand rate to be suppressed (Figure 3C). Because the enhancement of sense target cleavage was caused by a mutation in the sense siRNA strand, which does not participate in the recognition of the sense target, the effect of the mutation must be on a step in the RNAi pathway that is spatially or temporally coupled to siRNA unwinding. However, the suppression of anti-sense target cleavage clearly might have resulted from the single-nucleotide mismatch between the sense strand and its target RNA generated by the C-to-U substitution.

To test if the suppression of the rate of anti-sense target cleavage was a consequence of the position 19 mismatch, a different strategy was used to unpair the 5' end of the anti-sense strand. Figure 3D shows an siRNA in which the sense-strand is identical to that in Figure 3B, but the first nucleotide of the anti-sense strand has been changed from G to U, creating a U-C mismatch at its 5' end, in place of the G-A of Figure 3C. Nonetheless, this siRNA duplex showed pronounced asymmetry, with the anti-sense strand guiding target cleavage to the nearly complete exclusion of the sense strand (Figure 3D). Thus, the suppression of the cleavage rate of the sense-strand in Figure 3C was not a consequence of the position 19 mismatch. This finding is consistent with previous studies that suggest that mismatches with the target RNA are well tolerated if they occur near the 3' end of the siRNA guide strand (Amarzguioui et al., 2003). The finding that the siRNAs in Figures 3C and 3D display profound asymmetry demonstrates that both the enhancement of the target cleavage rate of the anti-sense strand and the suppression of the function of the sense strand is a consequence of their relative abilities to enter the RNAi pathway, not their intrinsic capacity to direct target cleavage.

Finally, the sense strand of Figure 3C was paired with the anti-sense strand of Figure 3D to create the siRNA duplex shown in Figure 3E. The sense strand of this siRNA, like that in Figure 3C, contains a mismatch with the anti-sense target at position 19. Like the anti-sense siRNA strand in Figure 3D, the anti-sense strand contains a mismatch with the sense target at position 1. This siRNA duplex directs target anti-sense cleavage significantly better than the siRNA in Figure 3C, despite the fact that the two siRNAs contain the same sense strand (Figure 3E).

Figures 3F, G, and H show a similar analysis in which the 5' end of the sense strand or position 19 of the anti-sense strand of the siRNA in Figure 3B was altered to produce siRNA duplexes in which the 5' end of the sense strand was either fully unpaired (Figures 3F and G) or paired in an A:U base pair (Figure 3H). Again, unpairing the 5' end of an siRNA strand—the sense strand, in this case—caused that strand to function to the exclusion of the other strand. When the sense strand 5' end was present in an A:U base pair and the anti-sense strand 5' end was in a G:C pair, the sense strand dominated the reaction (Figure 3H), although now the anti-sense strand showed activity similar to that seen for the original siRNA (Figure 3B) in which both strands were in G:C pairs at their 5' ends. Converting the unpaired 5' end of the siRNAs in Figure 3 to an A:U pair reduced the functional asymmetry of the two strands by enhancing the efficacy of the sense strand (Figure 3E) or the anti-sense strand (Figure 3H). The relative ease with which the 5' ends of the two siRNAs can be liberated from the duplex determines the degree of asymmetry. Additional data supporting this idea is shown in Figure 8, using a different siRNA. Figure 8B shows an siRNA that cleaved the two *sod1* target RNAs (Figure 8A), with modest functional asymmetry that reflects the collective base pairing strength of the first four or five nucleotides of each siRNA strand (Figure 8E; see below). Asymmetry was dramatically increased when a G:U wobble was introduced at the 5' end of the anti-sense strand of the siRNA (Figure 8C), but no asymmetry was seen when the individual single-strands strands were used to trigger RNAi (Figure 8D), demonstrating that differential RISC assembly, not target accessibility, explains the functional asymmetry of the siRNA duplex.

Together, the data in Figures 1, 2, and 8 indicates that the symmetry of RISC assembly is determined by a competition between the fraying of the 5' ends of the two siRNAs in the duplex. Such fraying may initiate a directional process of unwinding in

which the strand at which unwinding is initiated preferentially enters RISC. Such a model requires that either that RISC assembly factors or RISC components themselves are loaded onto one of the two siRNA strands before unwinding is completed, or that information about the siRNA strands prior state of pairing is retained, perhaps by a
5 protein such as the helicase remaining bound to a strand.

Example IV: A single hydrogen bond can determine which strand of an siRNA duplex directs RNAi

To explore this hypothesis further, additional changes were made to the *sod1*-
10 specific siRNA in Figure 3. These modifications alter the function of the two strands of the siRNA, but do not change the site cleaved on the two target RNA's. In Figure 3A, the anti-sense strand of Figure 3B was paired with a sense strand identical to that in Figure 3B except the 5' G was replaced with inosine (I). Like G, I pairs with C, but makes two instead of three hydrogen bonds. In this respect, an I:C pair is similar in
15 energy to an A:U pair. The resulting siRNA was functionally asymmetric, when the sense-strand began with an I, it directed target cleavage more efficiently than antisense-strand (Figure 4A). The asymmetry reflects an enhancement in efficacy of the sense siRNA strand, with little loss in the function of the anti-sense strand. An inosine at the 5' end of the anti-sense strand had the opposite effect. When the G at position 1 of the
20 anti-sense strand was substituted with inosine and the sense strand is that of Figure 3B, the anti-sense strand was enhanced relative to the sense strand (Figure 4B). Thus, the strand whose 5' end is in the weaker base pair was more effective at target cleavage.

Remarkably, when the 5' nucleotides of both siRNA strands engage in I:C base pairs (Figure 4C), the relative efficacy of the two siRNA strands is restored to that reported in
25 Figure 3B. The slightly faster rate for anti-sense target cleavage than for sense target cleavage is also seen for RNAi triggered with the individual, inosine-containing single strands, indicating that it reflects a difference in the intrinsic capacity of the two strands to guide cleavage, rather than a difference in RISC assembly. Although the relative rates of cleavage of the two strands are comparable for the siRNAs in Figure 3B and 4C, the
30 absolute rates are faster for the siRNA in Figure 4C. These data indicate that production of RISC from an individual strand is governed both by the relative propensity of the siRNA 5' end to fray compared to that of its complementary strand and by the absolute

propensity of the siRNA 5' end to fray. This latter finding is particularly unexpected, in that it shows that a difference of a single hydrogen bond has a marked effect on the rate of RISC assembly. siRNA end fraying provides an entry site for an ATP-dependent RNA helicase that unwinds siRNA duplexes (Figure 4). The helicase makes many abortive attempts to dissociate the two siRNA strands before succeeding to load one strand into RISC. The involvement of a helicase in RISC assembly is supported by previous observations: (1) both siRNA unwinding and production of functional RISC require ATP *in vitro* (Nykänen et al., 2001) and (2) several proteins with sequence homology to ATP-dependent RNA helicases have been implicated in RNA silencing (Wu-Scharf et al., 2000; Dalmay et al., 2001; Hutvágner and Zamore, 2002; Ishizuka et al., 2002; Kennerdell et al., 2002; Tabara et al., 2002; Tijsterman et al., 2002).

The effect of single-nucleotide mismatches in this region of the siRNA, using a series of siRNAs containing a mismatch at the second, third, or fourth position of each siRNA strand was further tested. The siRNAs bearing G:U wobble pairs at the second, third, or both second and third positions (Figure 11) was also analyzed. The results of this series demonstrate that mismatches, but not G:U wobbles, at positions 2-4 of an siRNA strand alter the relative loading of the two siRNA strands into RISC. Mismatches at position five, have very modest effects on the relative loading of the siRNA strands into RISC (data not shown). In contrast, the effects of internal mismatches at positions 6-15 cannot be explained by their influencing the symmetry of RISC assembly (data not shown). In sum, these data are consistent with the action of a non-processive helicase that can bind about four nucleotides of RNA.

Example V: Implications of siRNA asymmetry in miRNA biogenesis

One implication of the findings presented herein is that although siRNAs are predominantly present as duplexes at steady state *in vitro* (Nykänen et al., 2001) and perhaps *in vivo* (Hamilton and Baulcombe, 1999; Djikeng et al., 2001), both strands of an siRNA are unlikely to be present equally in RISC. That is, the strength of the base pairs at the 5' ends of the two siRNA strands can influence their accumulation as single-strands. When the 5' end of one strand is unpaired, this asymmetry can be nearly absolute. This observation suggested that asymmetric incorporation into RISC, as a consequence of directional unwinding from a frayed end of an siRNA duplex, might also

explain why miRNAs accumulate as single strands. Animal miRNAs are derived from the double-stranded stem of ~ 70 nt stem-loop precursor RNAs (Lee et al., 1993; Pasquinelli et al., 2000; Reinhart et al., 2000; Lagos-Quintana et al., 2001; Lau et al., 2001; Lee and Ambros, 2001; Lagos-Quintana et al., 2002). pre-miRNAs stems are only
5 partially double-stranded; the typical pre-miRNA contains mismatches, internal loops, and G•U base pairs predicted to distort an A-form RNA helix. miRNAs are generated from pre-miRNAs by the double-stranded RNA-specific endonuclease Dicer (Hutvagner et al., 2001; Grishok et al., 2001; Ketting et al., 2001). It was previously proposed by the instant inventors that miRNAs are single-stranded because helical discontinuities
10 constrain Dicer to break only two, rather than four, phosphodiester bonds, yielding a single-stranded miRNA, rather than an siRNA-like duplex (Hutvagner et al., 2001). Such a mechanism has precedent, because *E. coli* RNase III can be constrained by helical distortions to make only one or two breaks in an RNA chain (Chelladurai et al., 1993).

15 An alternative hypothesis is that the Dicer cleaves four phosphodiester bonds in all of its substrates, both long dsRNA and pre-miRNAs, and always generates a product with the essential siRNA duplex (Hutvagner and Zamore, 2002; Reinhart et al., 2002; Lim et al., 2003b). This mechanism for miRNA production was originally suggested by Bartel and colleagues. Using a small RNA cloning strategy to identify mature miRNAs
20 in *C. elegans*, they recovered small RNAs corresponding to the non-miRNA side of the precursor's stem (Lim et al., 2003b). Although these 'miRNA*' sequences were recovered at about 100 times lower frequency than the miRNAs themselves, they could always be paired with the corresponding miRNA to give 'miRNA duplexes' with 2 nt overhanging 3' ends (Lim et al., 2003b). Their data suggest that miRNAs are born as
25 duplexes, but accumulate as single-strands because some subsequent process stabilizes the miRNA, destabilizes the miRNA*, or both.

The incorporation of miRNA into RISC is this process. Our results with siRNA suggest that preferential assembly of a miRNA into the RISC would be accompanied by destruction of the miRNA. If the rate asymmetric RISC assembly was
30 faster than the production of the miRNA duplexes, only single-stranded miRNAs would be observed at steady-state (Figure 4). The accumulation of single-strands and not duplexes for miRNAs would simply be a consequence of Dicer being significantly less

efficient in cleaving pre-miRNAs compared to long dsRNA (Hutvagner et al., 2001). The rate of asymmetric RISC assembly might be faster than the production of miRNA duplexes, so only single-stranded miRNAs would be observed at steady-state. Two key predictions of this hypothesis are that (1) purified Dicer should cleave pre-miRNAs into equal amounts of miRNA and miRNA* products and (2) pre-miRNA structures should be processed by Dicer into duplexes with the 5' end of the miRNA strand frayed or weakly hydrogen bonded and the 5' end of the miRNA* strand more securely base paired.

10 **A. Dicer cleaves pre-*let-7* symmetrically**

To begin to test the idea that pre-miRNA are cleaved by Dicer to generate a product with an essential structure of an siRNA a duplex, we incubated the *Drosophila* pre-miRNA, pre-*let-7*, with purified, recombinant Dicer and analyzed the products by Northern hybridization using probes specific for either the 5' side of the precursor stem that encodes mature *let-7* or for products derived from the 3' side of the precursor stem (*let-7** products). As a control, the *let-7* precursor RNA was incubated in *Drosophila* embryo lysate, which recapitulates both pre-*let-7* maturation and RNAi *in vitro*. As previously reported, incubation of pre-*let-7* RNA in the lysate produced a single band corresponding to authentic *let-7*, but no *let-7** products (Hutvagner et al., 2001; Figure 5A and 5B). In contrast, incubation of pre-*let-7* with Dicer yielded approximately equal amounts of *let-7* and *let-7** products. At least three distinct RNAs were generated from each side of the stem, rather than the single band corresponding to mature *let-7* observed in the embryo lysate. Thus, the absence of *let-7** *in vivo* and in the embryo lysate reaction cannot be explained by the influence of pre-*let-7* structure on Dicer.

25

B. Asymmetric RISC assembly explains why miRNAs are single-stranded

If Dicer cleaves both sides of the pre-*let-7* stem, then some step downstream from Dicer action selects mature *let-7* from an siRNA-like duplex in which *let-7* is paired with *let-7**. A good candidate for such a step would be the asymmetric incorporation of *let-7* into RISC, accompanied by the degradation of *let-7**. To test this idea, the siRNA that might be formed if pre-*let-7* were cleaved by Dicer into an siRNA

duplex-like structure was deduced. The sequence of this 'pre-*let-7* siRNA,' generated by 'conceptual dicing,' is shown in Figure 6A (see below). Notably, the 5' end of *let-7* is unpaired in this duplex, whereas the 5' end of the *let-7** strand is in an A:U base pair. The results presented in Figures 2, 3, and 4 suggest that this structure should cause the *let-7* strand to enter the RISC to the near exclusion of the *let-7** strand, which would consequently be degraded.

C. miRNA versus miRNA* selection in *Drosophila*

This analysis was next extended to the other published *Drosophila* miRNA genes (Lagos-Quintana et al., 2001). For each precursor structure, the double-strand predicted to be produced by Dicer. These conceptually diced duplexes are shown in Figure 6A. For 23 of the 27 duplexes generated by this analysis (including pre-*let-7*), the difference in the base pairing of first five nucleotides of the miRNA versus miRNA* strands accurately predicted the miRNA, and not the miRNA*, accumulates *in vivo*. The analysis succeeded irrespective of which side of the pre-miRNA stem encoded the mature miRNA. This analysis, previous observations that single mismatches in the first four nucleotides of an siRNA strand, an initial G:U wobble pair, but not internal G:U wobbles, directed the asymmetric incorporation of an siRNA strand into RISC (Figures 1, 2, 3, 8, 9, and 11). However, no difference was discerned in the propensity to fray of the 5' ends of the miRNA and * strands for miR-4, miR-5, the three miR-6-2 paralogs, and miR-10. Therefore, it could not be explained why a particular strand would accumulate as the mature miRNA for these three miRNA precursors. miR-5 and miR-10, like other *Drosophila* miRNAs, were identified by the cloning and sequencing of small RNAs from embryos (Lagos-Quintana et al., 2001). Determinants other than end fraying appear to function in the selection of miR-4 and miR-6; these unknown determinants may also play a role in the assembly of an siRNA strand into RISC. However, miR-5 and miR-10 were cloned only once, raising the possibility that miR-5* or miR-10* is present in embryos, but not represented among the library of small RNAs from which the miRNAs were cloned. Similarly, miR-6 is encoded by three paralogous genes, only one of which we predict to produce detectable amounts of the miR*, so this * strand might have also gone undetected. To test if both the miRNA and * strands might accumulate for some or all of these three genes, Northern hybridization was used

to examined the relative abundance of miR-10 and miR-10* in adult *Drosophila* males and females, and in syncytial blastoderm embryos. The results detected both miR-10* and mi-R10 *in vivo* (Figure 6C). In fact, the results indicated that more miR-10* was detected than miR-10 in adult males. This finding strengthens the proposal that miRNA genes (*i.e.*, premiRNA's) uniquely specify on which side of the stem the miRNA residues by generating siRNA-like duplexes from which only one of the two strands of the duplex is assembled into RISC. When these double-stranded intermediates do not contain structural features enforcing asymmetric RISC assembly, both strands accumulate *in vivo*. It is possible that pre-miRNAs such as pre-miR-10, which generates roughly equal amounts of small RNA products from both sides of the precursor stem, simultaneously regulate target RNAs with partial complementary to both small RNA products.

Example VI: Increased rate of siRNA efficiency through the use of dTdT

tails

Art-recognized protocols for designing siRNA duplexes teach the inclusion of dTdT tails (*i.e.*, 2-nucleotide overhangs consisting of dTs). Two duplexes were created to test whether the addition of 3' overhanging dTdT tails increases the rate of siRNA targeting efficiency of the *Cu, Zn superoxide-dismutase-1 (Sod1)* mRNA. The first duplex contained sense and antisense strands, each including 21 nucleotides with 19 complementary bases plus 2-nucleotide overhangs (the overhangs consisting of bases in common with the target sequence). The second duplex contained sense and antisense strands, each including 19 complementary nucleotides (in common with the *Sod1* target), plus 2-nucleotide dTdT tails at the 5' end of the strand (not matching the *Sod1* target). Results demonstrate that the rate of siRNA efficiency improved ~ 8 fold ~ when using the duplex having mismatched dTdT tails (Figure 12).

Discussion of Examples I-VI: Implications for RNA silencing

The observations described herein provide rules for siRNA design. Clearly, siRNA structure can profoundly influence the entry of the anti-sense siRNA strand into the RNAi pathway. Thus, the sequence of the siRNA, rather than that of the target site, may explain at least some previous reports of ineffective siRNAs duplexes. Such

inactive duplexes may be coaxed back to life by modifying the sense strand of the siRNA to reduce the strength of the base pair at the 5' end of the anti-sense strand. An example of this *in vitro* is shown in Figure 9, for an ineffective siRNA directed against the *huntingtin* (*htt*) mRNA (Figure 9A). Changing the G:C (Figure 9B) to an A:U pair (Figure 9C) or a G-A mismatch (Figure 9D) dramatically improved its target cleavage rate *in vitro* and its efficacy *in vivo* (Eftim Milkani, NA, and PDZ, unpublished observations). In fact, Khvorova and colleagues have found that a low base-pairing stability at the 5' end of the antisense strand, but not the sense strand, is a prerequisite for siRNA function in cultured mammalian cells (Anastasia Khvorova, Angela Reynolds, and Sumedha D. Jayasena, manuscript submitted).

siRNAs designed to function asymmetrically may also be used to enhance RNAi specificity. Recently, expression profiling studies have shown that the sense-strand of an siRNA can direct off-target gene silencing (A.L. Jackson, et al. (2003) *Nature Biotechnology*, May 18; Smizarov et al., PNAS, 2003; Chi et al., PNAS, 2003). The data presented herein provide a strategy for eliminating such sequence-specific but undesirable effects: redesigning the siRNA so that only the anti-sense strand enters the RNAi pathway.

The observations described herein provide new design rules for the construction of short hairpin RNAs (shRNAs), which produce siRNAs transcriptionally in cultured cells or *in vivo* (Brummelkamp et al., 2002; McManus et al., 2002; Paddison et al., 2002; Paul et al., 2002; Sui et al., 2002; Yu et al., 2002). shRNA strategies typically employ a *Pol* III promoter to drive transcription, so the shRNA must begin with several G residues. As a consequence, the 5' end of the siRNA may be sequestered in a G:C base pair, significantly reducing entry of the anti-sense strand into the RNAi pathway. To avoid this problem, the anti-sense strand of the desired siRNA can be placed on the 3' side of the loop, so as to ensure that its 5' end is in an A:U, rather than the G:C pair typically encoded. Alternatively, the hairpin can be designed to place the 5' end of the anti-sense siRNA strand in a mismatch or G•U base pair, in which case it can be placed on either side of the stem. Moreover, a recent report suggests that some shRNAs may induce the interferon response (Bridge et al., 2003). The data suggest that mismatches and G:U pairs could be designed into these shRNAs simultaneously to promote entry of the correct siRNA strand into the RNAi pathway and to diminish the

capacity of the shRNA stem to trigger non-sequence specific responses to double-stranded RNA.

Finally, the data identify an unanticipated step in the RNAi pathway: the direct coupling of siRNA unwinding to RISC assembly. This finding suggests that the helicase
5 responsible for unwinding siRNA duplexes will be intimately linked to other components of the RNAi machinery. Identifying the helicase and the proteins with which it functions to assemble the RISC is clearly an important challenge for the future.

Example VII: The siRNA-programmed RISC is an enzyme

10 RISC programmed with small RNA in vivo catalyzes the destruction of target RNA in vitro without consuming its small RNA guide (Tang et al., 2003) (Hutvagner et al., 2002). To begin a kinetic analysis of RISC, the RISC programmed in vitro with siRNA is likewise a multiple-turnover enzyme was first confirmed. To engineer an RNAi reaction that contained a high substrate concentration relative to RISC, an siRNA
15 was used in which the guide strand is identical to the let-7 miRNA, but unlike the miRNA, the let-7 siRNA is paired to an RNA strand anti-sense to let-7 (Hutvagner et al., 2002). The let-7 strand of this siRNA has a high intrinsic cleaving activity, but a reduced efficiency of incorporation into RISC (Figure 19A).

After incubating the let-7 siRNA with *Drosophila* embryo lysate in the presence
20 of ATP, RISC assembly was inactivated by treatment with N-ethyl maleimide (NEM), and the amount of RISC generated was measured using the previously described tethered 2'-O-methyl oligonucleotide assay (Hutvagner et al., 2004; Schwartz et al., 2003) (Figure 19 B,C). The amount of let-7 programmed RISC increased with increasing siRNA concentration, until the assembly reaction began to saturate at ~50
25 nM, reaching an asymptote between 3 and 4 nM RISC. Using 0.6 nM RISC, >50 cycles of target recognition and cleavage per enzyme complex (data not shown) was observed, confirming that siRNA-programmed RISC is a multiple-turnover enzyme.

Example VIII: Multiple-turnover is limited by product release

30 The evaluation of the kinetics of siRNA-directed target cleavage in the presence or absence of ATP was further performed (Figure 13). RISC was assembled in the presence of ATP, then the energy regenerating enzyme, creatine kinase, was inactivated with NEM, and ATP depleted by adding hexokinase and glucose (-ATP conditions). For

+ATP measurements creatine kinase was added to the reaction after NEM-treatment, and the hexokinase treatment was omitted. A faster rate of cleavage in the presence than in the absence of ATP was observed. This difference was only apparent late in the reaction time course, indicating that the ATP-dependent rate of cleavage was faster than the ATP-independent rate only at steady state (Figure 13 A). The analysis was repeated in more detail (Figure 13 B). In the absence of ATP, a burst of cleaved product early in the reaction, followed by a ~4-fold slower rate of target cleavage was observed. No burst was observed in the presence of ATP (Figure 13A). If the burst corresponds to a single-turnover of enzyme, then extrapolation of the slower steady state rate back to the y-axis should give the amount of active enzyme in the reaction. The y-intercept at the start of the reaction for the steady-state rate was 4.9 nM, in good agreement with the amount of RISC estimated using the tethered 2'-O-methyl oligonucleotide assay (~4 nM; Figure 13 B).

In principle, ATP could enhance target recognition by RISC, promote a rearrangement of the RISC/target complex to an active form, facilitate cleavage itself, promote the release of the cleavage products from the siRNA guide strand, or help restore RISC to a catalytically competent state after product release. All of these steps, except product release and restoration to catalytic competence, should affect the rate of both multiple and single-turnover reactions. Therefore, the rate of reaction in the presence and in the absence of ATP under conditions in which RISC was in excess over the RNA target was analyzed. At early times under these conditions, the reaction rate should reflect only single-turnover cleavage events, in which events after cleavage do not determine the rate of reaction. Using single-turnover reaction conditions, identical rates of RISC-mediated cleavage in the presence or absence of ATP was observed (Figure 13 C). Thus, ATP must enhance a step that occurs only when each RISC catalyzes multiple cycles of target cleavage.

If product release is rate-determining for multiple-turnover catalysis by RISC in the absence, but not the presence, of ATP, then modifications that weaken the strength of pairing to the target RNA might enhance product release, but would not be expected to accelerate the return of the RISC to a catalytically competent state. Mismatches between the siRNA and its RNA target at the 3' end of the siRNA guide strand was incorporated and designed the siRNAs to be functionally asymmetric, ensuring efficient and

predictable incorporation of the let-7 strand into RISC (Figure 14 A). The reaction velocity under conditions of substrate excess in the presence and in the absence of ATP for siRNAs with zero to four mismatches between the guide strand 3' end and the RNA target were compared. Cleavage was measured from 100 and 540 s, when > 90% of the target remained uncleaved, ensuring that the multiple-turnover reaction was at steady state. Even a single 3' mismatch between the siRNA and its target increased the -ATP rate, relative to the +ATP rate, and siRNAs with two or more mismatches showed no significant difference in rate between the presence and absence of ATP (Figure 14B). The results indicated that in the absence of ATP, product release is the rate-determining step for siRNAs fully matched to their RNA targets.

Example IX: siRNA:target complementarity and RISC function

Mismatches between the siRNA and its target facilitate product release, but not without cost: the rate of reaction, irrespective of ATP concentration, decreases with each additional 3' mismatch: When the concentration of RISC was ~16–80-fold greater than the target RNA concentration, each additional mismatch between the 3' end of the siRNA guide strand and the RNA target further slowed the reaction (Figure 14 C,D). Under conditions of substrate excess, the effect of mismatches between the 3' end of the siRNA guide strand and its RNA target was more striking (Figure 15A): the rate of cleavage slowed ~20% for each additional mismatch. To test the limits of the tolerance of RISC for 3' mismatches, cleavage under modest (8-fold, Figure 15 B) and vast (~80-fold, Figure 15 C and 16 C) enzyme excess over target RNA was analyzed. Remarkably, cleavage was detected for siRNAs with as many as nine 3' mismatches to the RNA target (Figure 15 C and 16 C), but only after 24 hour incubation. No cleavage was detected for an siRNA with ten 3' mismatches to the RNA target (Figure 15 C).

Linsley and colleagues have proposed siRNA-directed down-regulation of an mRNA with as few as eleven contiguous bases complementary to the siRNA guide strand (Jackson et al., 2003). In that study, the mRNA target paired with both nts 2–5 and nts 7–17 of the siRNA guide strand, but mismatched at nts 1 and 6 of the siRNA. Results indicated that up to five mismatched bases are tolerated between the 5' end of the siRNA and its RNA target (Figure 16 A,B). No cleavage was detected for siRNAs with six, seven, or eight 5' mismatches to the target, even after 24 hour incubation. The siRNA bearing eight mismatches between its 5' end and the let-7 complementary target

was fully active when eight compensatory mutations were introduced into the let-7 binding site (Figure 15C and 16B), demonstrating that mutation of the siRNA was not the cause for its inactivity against the mismatched target. Similarly, when eight mismatches with the 3' or 5' end of the siRNA were created by changing the sequence of the RNA target, target RNA cleavage when the target contained eight mismatches with the siRNA 3' end, but not with the 5' end was detected (Figure 16 B,C).

To begin to estimate the minimal number of base pairs between the siRNA and its target that permit detectable cleavage by RISC at 24 hour incubation, seven, eight, or nine 3' mismatches with increasing numbers of 5' mismatches were combined (Figure 16 C). Cleavage was detected for as many as nine 3' mismatches. However, no detectable cleavage occurred when seven, eight, or nine 3' mismatches were combined with two or more 5' mismatches. In contrast, a single 5' mismatch (p1) enhanced target cleavage directed by all three 3' mismatched siRNAs. Only 6% of the target RNA was cleaved after 24 hours when the siRNA contained nine contiguous 3' mismatches with the target RNA, but 10% was cleaved when the siRNA contained both nine 3' mismatches and a single (p1) 5' mismatch. Cleavage was similarly enhanced by the addition of a p1 mismatch to seven 3' mismatches (49% cleavage versus 75% cleavage at 24 hours) or to eight 3' mismatches (21% versus 42% cleavage at 24 hours). The finding that unpairing of the first base of the siRNA guide strand potentiated cleavage under single-turnover conditions indicated that a conformational change occurs in RISC during which the paired p1 base becomes unpaired prior to cleavage. Intriguingly, p1 is often predicted to be unpaired for miRNAs bound to their targets (Lewis et al., 2003; Rhoades et al., 2002; Stark et al., 2003).

For siRNAs that pair fully with their RNA targets, the scissile phosphate always lies between the target nucleotides that pair with siRNA bases 10 and 11 (Elbashir et al., 2001; Elbashir et al., 2001). Analysis at single nucleotide resolution of the 5' cleavage products generated by siRNAs with three, four, or five 5' mismatches (Figure 16 D) or six 3' terminal mismatches (data not shown) revealed that the scissile phosphate on the target RNA remained the same, even when five 5' nts of the siRNA guide strand were mismatched with the target RNA (Figure 16 D). As discussed below, this result indicates that the identity of the scissile phosphate is a consequence of the structure of RISC, rather than being measured from the 5' end of the helix formed between the siRNA and its RNA target.

Example X: Kinetic analysis of RISC catalysis

The role of nucleotides in the terminal regions of the siRNA guide strand in directing RISC activity was next studied. Reduced pairing between an siRNA and its target might disrupt the binding of RISC to its target. Alternatively, mismatches might disrupt the structure, but not the affinity, of the siRNA/target interaction. Fully matched siRNAs are thought to form a 21 base-pair, A-form helix with the target RNA (Chiu et al., 2003; Shiu et al., 2002), but do all parts of this helix contribute equally to target binding or do some regions provide only a catalytically permissive geometry? To distinguish between these possibilities, the Michaelis-Menten kinetics of siRNA-directed target-RNA cleavage for a perfectly matched siRNA and for three siRNAs mismatched at their termini was analyzed. siRNAs were assembled into RISC, then diluted with reaction buffer to the desired RISC concentration and mixed with target RNA. For each siRNA, the initial velocity of reaction was determined at multiple substrate concentrations (Figure 21 A), and K_M and k_{cat} determined from a non-linear least squares fit of substrate concentration versus initial velocity (Figure 17 A). By this assay, the K_M of the let-7 siRNA with complete complementarity to its target was ~ 8.4 nM was estimated (Table 1). A significant difference in K_M , within error, between the fully paired siRNA and siRNA variants bearing three to five mismatches at their 3' end or three mismatches at their 5' end was not detected (Figure 17A and Table 1). For the mismatched siRNAs a higher than optimal enzyme concentration in order to detect cleavage was used. Therefore, the K_M measurements for the mismatched siRNAs represent an upper bound for the actual K_M values.

While the K_M was unaltered for the let-7 siRNA containing several terminal mismatches, the turnover number, k_{cat} , was decreased by terminal mismatches (Table 1). Three mismatches at either the 3' or the 5' end of the siRNA halved the k_{cat} . The introduction of five, 3' mismatches also had no significant effect on K_M , yet decreased k_{cat} nearly 17-fold (Table 1).

Table 1 Summarizes the kinetic data from the analysis in Figure 17A. For comparison, the K_M and k_{cat} values of four well studied protein enzymes are provided. K_M and $k_{cat} \pm$ error of fit are reported.

Example XI: KM reflects the binding strength of RISC

To estimate the contribution of binding to K_M , a competition assay that measures the ability of 2'-O-methyl oligonucleotides to inhibit target cleavage by RISC was used (Figure 17 B,C). Such a strategy was used previously to analyze the mechanism of target destruction by antisense oligonucleotides that recruit RNase H (Lima et al., 1997). The anticipation was that 2'-O-methyl oligonucleotides would act as competitive inhibitors of RISC, because they bind to RISC containing complementary siRNA but not to RISC containing unrelated siRNA (Hutvagner et al., 2004; Meister et al., 2004). Thirty-one nt long, 2'-O-methyl oligonucleotides were designed as described previously (Hutvagner et al., 2004), taking care to exclude sequences predicted to form stable internal structures. 2'-O-methyl oligonucleotides were chosen because of their marked stability in *Drosophila* lysate and because they can be added to the reaction at high μM concentration.

Competition by 2'-O-methyl oligonucleotides and bona fide RNA targets was quantitatively similar. The reaction velocities of siRNA-directed cleavage of a 32P-radiolabeled target in the presence of increasing concentrations of unlabeled capped RNA target or a 31-nt 2'-O-methyl oligonucleotide corresponding to the region of the target containing the siRNA binding site was analyzed (Figure 17 B). Lineweaver-Burk analysis of the data confirm that 2'-O-methyl oligonucleotides act as competitive inhibitors of RISC (data not shown). These data were used to calculate K_i values for the perfectly matched RNA and 2'-O-methyl competitors. For the capped RNA competitor, the K_i was $\sim 7.7 \pm 4$ nM (Figure 17 B), nearly identical to the K_M for this siRNA, 8.4 nM (Table 1). The K_i for the perfectly matched 2'-O-methyl competitor oligonucleotide was 3.2 ± 1 nM (Figure 17 B), essentially the same, within error, as that of the all-RNA competitor. The results indicated that 2'-O-methyl oligonucleotides are good models for 5'-capped RNA targets and that the K_M for target cleavage by RISC is largely determined by the affinity (K_D) of RISC for its target RNA.

Although targets with more than five contiguous mismatches to either end of the siRNA are poor substrates for cleavage, they might nonetheless bind RISC and compete with the 32P-radiolabeled target RNA. The 2'-O-methyl oligonucleotide competition assay to determine the K_i values for oligonucleotides containing as many as eight mismatches to the siRNA guide strand was used (Figure 17 B). 2'-O-methyl

oligonucleotides with 3' terminal mismatches to the siRNA were good competitors: a four nucleotide mismatch with the 3' end of the siRNA increased the K_i by only ~3-fold (9.0 ± 0.9 nM) and an eight nucleotide mismatch with the 3' end of the siRNA increased the K_i by ~10-fold (34.8 ± 7 nM). In contrast, mismatches with the 5' end of the siRNA had a dramatic effect on binding. A four nucleotide mismatch to the 5' end of the siRNA increased the K_i ~12-fold (36.4 ± 9.2 nM) and an eight nucleotide mismatch to the 5' end of the siRNA increased the K_i 53-fold (173 ± 16 nM). The differential effect on binding between 5' and 3' mismatches was maintained even at the center of the siRNA: a 2'-O-methyl oligonucleotide bearing four mismatches with siRNA nucleotides 11, 12, 13, and 14 (4 nt 3' central mismatch, Figure 17 B) bound more tightly to RISC (*i.e.*, had a lower K_i) than an oligonucleotide with four mismatches to siRNA positions 7, 8, 9, and 10 (4 nt 5' central mismatch, Figure 17 B).

Discussion of Examples VII-XI:

RISC programmed with exogenous siRNA is an enzyme, capable of multiple rounds of target cleavage. Previous studies showed that cleavage of a target RNA by RISC does not require ATP (Nykänen et al., 2001; Tomari et al., 2004). The more detailed kinetic analysis presented herein indicates that there are no ATP-assisted steps in either target recognition or cleavage by *Drosophila* RISC; no difference in rate in the presence or absence of ATP for RNAi reactions analyzed under conditions of substrate excess at early time points (pre-steady state) or under conditions of enzyme excess where the reaction was essentially single-turnover was detected. In contrast, the steady-state rate of cleavage under multiple turnover conditions was enhanced four-fold by ATP. The results indicate that release of the products of the RISC endonuclease is rate determining under these conditions in the absence of ATP, but not in the presence of ATP. The most straightforward explanation for this finding is that an ATP-dependent RNA helicase facilitates the dissociation of the products of target cleavage from the RISC-bound siRNA. The involvement of such an ATP-dependent helicase in RNAi *in vivo* may explain why siRNAs can be active within a broad range of GC content (Reynolds et al., 2004).

In the presence of ATP, siRNA-programmed *Drosophila* RISC is a classical Michaelis-Menten enzyme. The guide strand of the siRNA studied here has the sequence of let-7, an endogenous miRNA. *In vivo*, let-7 is not thought to direct mRNA cleavage,

but rather is believed to repress productive translation of its mRNA targets. Nonetheless, the let-7 siRNA is among the most potent of the siRNAs we have studied *in vitro* and provides a good model for effective siRNA in general. With a k_{cat} of $\sim 7 \times 10^{-3} \text{ s}^{-1}$, the let-7 siRNA-programmed RISC was slow compared to enzymes with small molecule
5 substrates (Table 1). The K_M for this RISC was $\sim 8 \text{ nM}$. Enzymes typically have K_M values between 1- and 100-fold greater than the physiological concentrations of their substrates (Stryer et al., 1981). The results indicate that RISC is no exception: individual abundant mRNA species are present in eukaryotic cells at high pM or low nM concentration. The K_M of RISC is likely determined primarily by the strength of its
10 interaction with the target RNA, because the K_M is nearly identical to the K_i of a non-cleavable 2'-O-methyl oligonucleotide inhibitor.

Recently, a study of the kinetic parameters of target RNA cleavage by human RISC was described (Martinez et al., 2004). In that study, the minimal active human RISC was highly purified; in this study, *Drosophila* RISC activity was measured for the
15 unpurified, intact holo-RISC, believed to be an 80S multi-protein complex (Pham et al., 2004). Different siRNAs were used in the two studies. Nonetheless, the K_M and k_{cat} values reported here and for the minimal human RISC are remarkably similar: the K_M was 2.7–8.4 nM and the k_{cat} was $7.1 \times 10^{-3} \text{ sec}^{-1}$ for the let-7 siRNA-programmed *Drosophila* holo-RISC versus a K_M of 1.1–2.3 nM and a k_{cat} of $1.7 \times 10^{-2} \text{ sec}^{-1}$ for a
20 different siRNA in minimal human RISC. As in this study, a pre-steady-state burst was observed in the absence of ATP, consistent with the idea that product release is ATP-assisted *in vivo*.

The ratio of k_{cat} to K_M is a classical measure of enzyme efficiency and corresponds to the second order rate constant for the reaction when the concentration of
25 substrate is much less than the K_M . For the let-7 programmed RISC, k_{cat}/K_M equals $\sim 8.4 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$ ($\sim 8.4 \times 10^{-4} \text{ nM}^{-1} \text{ s}^{-1}$), a value far slower than the expected rate of collision of RISC with mRNA, $\sim 10^7 \text{ M}^{-1} \text{ s}$. It is possible that the rate of catalysis by RISC is constrained by the rate of conformational changes required for formation of the enzyme-substrate complex or by subsequent conformational rearrangements required for
30 catalysis. It is possible that siRNAs can be designed that significantly improve either the k_{cat} or K_M of RISC without compromising specificity.

Although siRNAs are typically envisioned to bind their target RNAs through 19 to 21 complementary base pairs, we find that the 5', central, and 3' regions of the siRNA

make distinct contributions to binding and catalysis (Figure 18). Measurements of K_M and K_i suggest that the 5' nucleotides of the siRNA contribute more to target binding than do the 3' nucleotides. At least for the siRNA examined here, the first three and the last five nucleotides of a 21 nt siRNA contribute little to binding. If the K_D of RISC bound to its target RNA is essentially its K_M , ~ 8 nM, then the free energy ($\Delta G^\circ = -RT \ln K_D$) of the let-7-programmed RISC:target interaction is approximately -11 kcal mol $^{-1}$, considerably less than the -35 kcal mol $^{-1}$ ($K_D \sim 10$ – 29) predicted³² for the let-7 RNA bound to a fully complementary RNA in 100 mM K $^+$ and 1.2 mM Mg $^{2+}$ at 25°C. It is possible that RISC discards potential binding energy by binding less tightly to its target, an siRNA in RISC gains the ability to discriminate between well matched and poorly match targets, but only for bases in the 5' region of the siRNA guide strand.

Mismatches between the central and 3' regions of an siRNA and its target RNA reduce k_{cat} far more than mismatches at the 5' end of the siRNA. These results fit well with recent findings by Doench and Sharp that translational repression by siRNA, designed to act like animal miRNA, is dramatically disrupted by mismatches with the 5' end of the siRNA, but not with similar mismatches at the 3' end¹⁸. These authors propose that miRNA binding is mediated primarily by nucleotides at the 5' end of the small RNA. In fact, complementarity between the 5' end of miRNAs and their targets has been required by all computational approaches for predicting animal miRNA targets (Rajewsky et al., 2004; Lewis et al., 2003; Stark et al., 2003; Enright et al., 2003). The instant discovery that central and 3' siRNA sequences must pair with the target sequence for effective target cleavage but not for target binding reinforces this view; both central and 3' miRNA sequences are usually mismatched with their binding sites in their natural targets (Lee et al., 1993; Reinhart et al., 2000; Brennecke et al., 2003; Abrahante et al., 2003; Vella et al., 2004; Xu et al., 2003; Johnston et al., 2003).

Formation of a contiguous A-form helix surrounding the scissile phosphate of the target mRNA has been proposed to be a quality control step for RISC-mediated target cleavage (Chiu et al., 2003). The instant invention discovers that RISC can direct cleavage when the siRNA is paired with the target RNA only at nts 2–12 of the guide strand, corresponding to one complete turn of an RNA:RNA helix. This region of the siRNA includes nts 2–8, which appear to be critical for miRNA recognition of mRNAs targeted for translational repression, plus two nts flanking either side of the scissile phosphate. The instant invention further discovers unpairing the first nt of the guide

strand enhances the activity of siRNAs with seven, eight or nine 3' mismatches to the RNA target is striking, since many miRNAs do not pair with their targets at this position. Furthermore, such pairing resembles that reported by Linsley and colleagues for siRNA-directed off-target effects in cultured mammalian cells (Jackson et al., 2003).

5 The requirement for a full turn of a helix may reflect a mechanism of 'quality control' by RISC. Since RISC can apparently assemble on any siRNA sequence, it must use the structure of the siRNA paired to its target to determine whether or not to cleave. Despite the apparent surveillance of the structure of the siRNA/target pair, the identity of the scissile phosphate is unaltered by extensive mismatch between the 5' end of the
10 siRNA and its target. Yet the scissile phosphate is determined by its distance from the 5' end of the siRNA guide strand (Elbashir et al., 2001; Elbashir et al., 2001). The simplest explanation for the instant discovery is that the scissile phosphate is identified by a protein loaded onto the siRNA during RISC assembly, *i.e.*, before the encounter of the RISC with its target RNA.

15 The remarkable tolerance of RISC for mismatches between the siRNA and its targets—up to nine contiguous 3' nucleotides—implies that a large number off-target genes should be expected for many siRNA sequences when RISC is present in excess over its RNA targets. However, RISC with extensive mismatches between the siRNA and target are quite slow to cleave, so off-target effects may be minimized by keeping
20 the amount of RISC as low as possible. These understandings of the molecular basis of siRNA-directed gene silencing assist the skilled artisan in creating siRNAs designed to balance the competing demands of siRNA efficacy and specificity.

Experimental Procedures

25 A. General methods

Drosophila embryo lysate preparation, *in vitro* RNAi reactions, and cap-labeling of target RNAs using Guanylyl transferase were carried out as previously described (Tuschl et al., 1999; Zamore et al., 2000). Target RNAs were used at ~ 5 nM
30 concentration to ensure that reactions occurred under single-turnover conditions. Target cleavage under these conditions was proportionate to siRNA concentrations. Cleavage products of RNAi reactions were analyzed by electrophoresis on 5% or 8% denaturing acrylamide gels. 5' end labeling and determination of siRNA unwinding status were

according to Nykänen et al. (Nykänen et al., 2001) except that unlabeled competitor RNA was used at 100-fold molar excess. Gels were dried, then exposed to image plates (Fuji), which were scanned with a Fuji FLA-5000 phosphorimager. Images were analyzed using Image Reader FLA-5000 version 1.0 (Fuji) and Image Gauge version 5
 3.45 or 4.1 (Fuji). Data analysis was performed using Excel (Microsoft) and IgorPro 5.0 (Wavemetrics).

B. *Drosophila* embryo lysate, siRNA labeling with polynucleotide kinase (New England Biolabs), target RNA preparation and labeling with guanylyl transferase were carried out as described (Hutvagner et al., 2002, Haley et al., 2003) and the forward primer sequence for 379 nt target mRNA was 5'-CGC TAA TAC GAC TCA CTA TAG CAG TTG GCG CCG CGA ACG A-3', and 5'-GCG TAA TAC GAC TCA CTA TAG TCA CAT CTC ATC TAC CTC C-3 for the 182 nt target. Reverse primers used to generate fully matched and mismatched target RNAs were: 5'-CCC ATT TAG GTG
 15 ACA CTA TAG ATT TAC ATC GCG TTG AGT GTA GAA CGG TTG TAT AAA AGG TTG AGG TAG TAG GTT GTA TAG TGA AGA GAG GAG TTC ATG ATC AGT G-3' (perfect match to let-7); 5'-CCC ATT TAG GTG ACA CTA TAG ATT TAC ATC GCG TTG AGT GTA GAA CGG TTG TAT AAA AGG TTG AGG TAG TAG GTT CAT GCA GGA AGA GAG GAG TTC ATG ATC AGT G-3' (7 nt 3' mismatch);
 20 5'-CCC ATT TAG GTG ACA CTA TAG ATT TAC ATC GCG TTG AGT GTA GAA CGG TTG TAT AAA AGG TTG AGG TAG TAG GTA CAU GCA GGA AGA GAG GAG TTC ATG ATC AGT G-3' (8 nt 3' mismatch); 5'-CCC ATT TAG GTG ACA CTA TAG ATT TAC ATC GCG TTG AGT GTA GAA CGG TTG TAT AAA AGG TTG AGG TAG TAG GAA CAT GCA GGA AGA GAG GAG TTC ATG ATC AGT
 25 G-3' (9 nt 3' mismatch); 5'-CCC ATT TAG GTG ACA CTA TAG ATT TAC ATC GCG TTG AGT GTA GAA CGG TTG TAT AAA AGG TAC TCC ATC TAG GTT GTA TAG TGA AGA GAG GAG TTC ATG ATC AGT G-3' (8 nt 5' mismatch); 5'-CCC ATT TAG GTG ACA CTA TAG ATT TAC ATC GCG TTG AGT GTA GAA CGG TTG TAT AAA AGG TAC TCG TAG TAG GTT GTA TAG TGA AGA GAG
 30 GAG TTC ATG ATC AGT G-3' (4 nt 5' mismatch). In Figures 13, 14, 15, 17A, 19 and 20A, the target sequence was 613 nt long; 379 nt in Figures 16A-C, 17B and 20B; and 182 nt in Figure 16D. All siRNAs were deprotected according to the manufacturer's

protocol (Dharmacon), 5'-radiolabeled where appropriate, then gel purified on a 15% denaturing polyacrylamide gel. 2'-O-methyl oligonucleotides were from Dharmacon. siRNA strands were annealed at high concentrations and serially diluted into lysis buffer (30 nM HEPES pH 7.4, 100 mM KOAc, and 2 mM MgCl₂). Gels were dried and
 5 imaged as described (Schwartz et al., 2003). Images were analyzed using Image Gauge 4.1 (Fuji). Initial rates were determined by linear regression using Excel X (Microsoft) or IgorPro 5.01 (Wavemetrics). Kaleidagraph 3.6.2 (Synergy Software) was used to determine K_M and K_i by global fitting to the equations: $V = (V_{max} \times S)/(K_M + S)$ and $V = (V_{max} \times K_i(app))/(K_i(app) + I)$, where V is velocity, S is target RNA
 10 concentration, and I is the concentration of 2'-O-methyl oligonucleotide competitor. K_i was calculated by correcting K_i(app) by the K_M and substrate concentration, $K_i = K_i(app)/(1 + (S/K_M))$.

C. siRNA preparation

15 Synthetic RNAs (Dharmacon) were deprotected according to the manufacturer's protocol. siRNA strands were annealed (Elbashir et al., 2001a) and used at 50 nM final concentration unless otherwise noted. siRNA single strands were phosphorylated with polynucleotide kinase (New England Biolabs) and 1 mM ATP according to the manufacturer's directions and used at 500 nM final concentration.

20

D. Target RNA preparation

Target RNAs were transcribed with recombinant, histidine-tagged, T7 RNA Polymerase from PCR products as described (Nykänen et al., 2001; Hutvágner and Zamore, 2002), except for sense *sod1* mRNA, which was transcribed from a plasmid
 25 template (Crow et al., 1997) linearized with Bam HI. PCR templates for *htt* sense and anti-sense and *sod1* anti-sense target RNAs were generated by amplifying 0.1 ng/ml (final concentration) plasmid template encoding *htt* or *sod1* cDNA using the following primer pairs: *htt* sense target, 5'-GCG TAA TAC GAC TCA CTA TAG GAA CAG TAT GTC TCA GAC ATC-3' and 5'-UUCG AAG UAU UCC GCG UAC GU-3'; *htt*
 30 anti-sense target, 5'-GCG TAA TAC GAC TCA CTA TAG GAC AAG CCT AAT TAG TGATGC-3' and 5'-GAA CAG TAT GTC TCA GAC ATC-3'; *sod1* anti-sense target,

5'-GCG TAA TAC GAC TCA CTA TAG GGC TTT GTT AGC AGC CGG AT-3' and
5'-GGG AGA CCA CAA CGG TTT CCC-3'.

Immobilized 2'-O-methyl oligonucleotide capture of RISC

5 The 5' end of the siRNA strand to be measured was 32 P-radiolabeled with PNK.
10 pmol biotinylated 2'-O-Methyl RNA was immobilized on Dynabeads M280 (Dyna) by incubation in 10 ml lysis buffer containing 2mM DTT for 1 h on ice with the equivalent of 50 ml of the suspension of beads provided by the manufacturer. The beads were then washed to remove unbound oligonucleotide. 50 nM siRNA was pre-incubated
10 in a standard 50 ml in vitro RNAi reaction for 15 min at 25°C. Then, all of the immobilized 2'-O-Methyl oligonucleotide was added to the reaction and the incubation continued for 1 h at 25°C. After incubation, the beads were rapidly washed three times with lysis buffer containing 0.1% (w/v) NP-40 and 2 mM DTT followed by a wash with the same buffer without NP-40. Input and bound radioactivity were determined by
15 scintillation counting (Beckman). The 5'-biotin moiety was linked via a six-carbon spacer arm. 2'-O-methyl oligonucleotides (IDT) were: 5'-biotin-ACA UUU CGA AGU AUU CCG CGU ACG UGA UGU U-3' (to capture the siRNA sense strand) 5'-biotin-CAU CAC GUA CGC GGA AUA CUU CGA AAU GUC C-3' (to capture the anti-sense strand).

20

mfold Analysis

To model the end of an siRNA, the following 16 nt RNA sequence were submitted to mfold 3.1: (37°C, 1 M NaCl): CGU ACU UUU GUA CGU G, UGU ACU UUU GUA CGU G, and UCG AAU UU UUC GAA A.

25

Pre-let-7 Processing

Pre-let-7 RNA was incubated with N-terminal histadine-tagged, human Dicer according to the manufacturer's directions (Gene therapy Systems) or in a standard
30 Drosophila embryo in vitro RNAi reaction as described previously (Hutvagner et al., 2001; Hutvagner and Zamore, 2002).

Northern Hybridization

Northern hybridization was essentially as described (Hutvagner et al., 2001). 50 mg total RNA was loaded per lane. 5' ³²P-radiolabeled synthetic RNA probes (Dharmacon) were: 5'-ACA AAU UCG GAU CUA CAG GGU-3' (to detect miR-10) and 5'-AAA CCU CUC UAG AAC CGA AUU U-3' (to detect miR-10*). The amount of miR-10 or miR-10* detected was normalized to the non-specific hybridization of the probe to 5S rRNA. Normalizing to hybridization of the probe to a known amount of a miR-10 or miR-10* synthetic RNA control yielded essentially the same result.

ATP-depletion and N-ethyl maleimide (NEM) Inhibition

RNAi reactions using *Drosophila* embryo lysate were as described (Haley et al., 2003). To compare 'minus' and 'plus' ATP conditions, samples were treated with 10 mM NEM (Pierce) for 10 min at 4°C, then the NEM was quenched with 11 mM dithiothreitol (DTT). For ATP depletion (−ATP), 1 unit of hexokinase and 20 mM (final concentration) glucose were added and the incubation continued for 30 min at 25°C. For 'plus' ATP reactions, 0.05 mg ml^{−1} (final concentration) creatine kinase and one-tenth volume H₂O substituted for hexokinase and glucose. The addition of fresh creatine kinase after NEM treatment did not rescue the defect in RISC assembly, but did restore ATP to high levels (Nykänen et al., 2001). ATP levels were measured using an ATP assay kit (Sigma) and a PhL luminometer (Mediators Diagnostika).

REFERENCES

- Abrahante, J. E., Daul, A. L., Li, M., Volk, M. L., Tennessen, J. M., Miller, E. A., and Rougvie, A. E. (2003). The *Caenorhabditis elegans* hunchback-like Gene *lin-57/hbl-1* Controls Developmental Time and Is Regulated by MicroRNAs. *Dev Cell* 4, 625-637.
- Achard, P., Herr, A., Baulcombe, D.C., and Harberd, N.P. (2004). Modulation of floral development by a gibberellin-regulated microRNA. *Development*. 131(14): 3357-65.
- Amarzguioui, M., Holen, T., Babaie, E., and Prydz, H. (2003). Tolerance for mutations and chemical modifications in a siRNA. *Nucleic Acids Res* 31, 589-595.
- Ambros, V. (2003). MicroRNA pathways in flies and worms: growth, death, fat,

- stress, and timing. *Cell*. 113(6): 673-6.
- Ambros, V., Bartel, B., Bartel, D. P., Burge, C. B., Carrington, J. C., Chen, X., Dreyfuss, G., Eddy, S. R., Griffiths-Jones, S., Marshall, M., *et al.* (2003). A uniform system for microRNA annotation. *RNA* 9, 277-279.
- 5 Amarzguioui, M., Holen, T., Babaie, E., and Prydz, H. (2003). Tolerance for mutations and chemical modifications in a siRNA. *Nucleic Acids Res* 31, 589-595.
- Ambros, V., Lee, R.C., Lavanway, A., Williams, P.T., and Jewell, D. (2003). MicroRNAs and other tiny endogenous RNAs in *C. elegans*. *Current Biology*. 13: 807-818.
- 10 Aravin, A.A., Lagos-Quintana, M., Yalcin, A., Zavolan, M., Marks, D., Snyder, B., Gassterland, T., Meyer, J., and Tuschl, T. (2003). The small RNA profile during *Drosophila melanogaster* development. *Developmental Cell*. 5(2): 337-50.
- Aukerman, M.J., and Sakai, H. (2003). Regulation of flowering time and floral organ identity by a MicroRNA and APETALA2-like target genes. *Plant Cell*. 10: 10.
- 15 Baehrecke, E.H. (2003). miRNAs: micromanagers of programmed cell death. *Current Biology*. 13(12): R473-5.
- Bartel, D.P. MicroRNAs: Genomics, biogenesis, mechanism, and function. *Cell* 116, 281-297 (2004).
- Bartel, D.P., and Chen, C.Z. (2004). Micromanagers of gene expression: the potentially
- 20 widespread influence of metazoan microRNAs. *Nature Reviews Genetics*. 396-400.
- Bartel, B. and Bartel, D.P. (2003). MicroRNAs: at the root of plant development? *Plant Physiology*. 132(2): 709-17.
- Bernstein, E., Caudy, A. A., Hammond, S. M., and Hannon, G. J. (2001). Role for a
- 25 bidentate ribonuclease in the initiation step of RNA interference. *Nature* 409, 363-366.
- Billy, E., Brondani, V., Zhang, H., Muller, U., and Filipowicz, W. (2001). Specific interference with gene expression induced by long, double-stranded RNA in mouse embryonal teratocarcinoma cell lines. *Proc Natl Acad Sci U S A* 98,
- 30 14428-14433.
- Boutla, A., Delidakis, C., and Tabler, M. (2003). Developmental defects by antisense-

- mediated inactivation of micro-RNAs 2 and 13 in *Drosophila* and the identification of putative target genes. *Nucleic Acids Research*. 31(17): 4973-4980.
- 5 Boutla, A., Delidakis, C., Livadaras, I., Tsagris, M., and Tabler, M. (2001). Short 5'-phosphorylated double-stranded RNAs induce RNA interference in *Drosophila*. *Curr Biol* 11, 1776–1780.
- 10 Brennecke, J., Hipfner, D. R., Stark, A., Russell, R. B., and Cohen, S. M. (2003). bantam Encodes a Developmentally Regulated microRNA that Controls Cell Proliferation and Regulates the Proapoptotic Gene *hid* in *Drosophila*. *Cell* 113, 25-36.
- 15 Bridge, A. J., Pebernard, S., Ducraux, A., Nicoulaz, A. L., and Iggo, R. (2003). Induction of an interferon response by RNAi vectors in mammalian cells. *Nat Genet*.
- 20 Brummelkamp, T. R., Bernards, R., and Agami, R. (2002). A system for stable expression of short interfering RNAs in mammalian cells. *Science* 296, 550-553.
- 25 Calin, G.A., Sevignani, C., Dumitru, C.D., Hyslop, T., Noch, E., Yendamuri, S., Shimizu, M., Rattan, S., Bullrich, R., Negrini, M., and Croce, C.M. (2004). Human microRNA genes are frequently located at fragile sites and genomic regions involved in cancer. *Proceedings of the National Academy of Sciences USA*. 101(9): 2999-3004.
- 30 Calin, G.A., Dumitru, C.D., Shimizu, M., Bichi, R., Zupo, S., Noch, E., Aldler, H., Rattan, S., Keating, M., Rai, K., Rassenti, L., Kipps, T., Negrini, M., Bullrich, F., and Croce, C.M. (2002). Frequent deletions and down-regulation of micro-RNA genes miR15 and miR16 at 13q14 in chronic lymphocytic leukemia. *Proceedings of the National Academy of Sciences USA*. 99(24): 15524-9.
- 35 Carrington, J.C., and Ambros, V. (2003). Role of microRNAs in plant and animal development. *Science*. 301(5631): 336-8.
- 40 Caudy, A. A., Myers, M., Hannon, G. J., and Hammond, S. M. (2002). Fragile X-related protein and VIG associate with the RNA interference machinery. *Genes Dev* 16, 2491-2496.

- Chelladurai, B., Li, H., Zhang, K., and Nicholson, A. W. (1993). Mutational analysis of a ribonuclease III processing signal. *Biochemistry* 32, 7549-7558.
- Chen, C.Z., Li, L., Lodish, H.F., and Bartel, D.P. (2004). MicroRNAs modulate hematopoietic lineage differentiation. *Science*. 303: 63-86.
- 5 Chen, X. (2003). A MicroRNA as a Translational Repressor of APETALA2 in Arabidopsis Flower Development. *Science (Science Express)* 10.1126/science.1088060).
- Chi, J.T., Chang, H.Y., Wang, N.N., Chang, D.S., Dunphy, N., and Brown, P.O. (2003). Genomewide view of gene silencing by small interfering RNAs. *Proceedings of the National Academy of Sciences USA*. 100(11): 6343-6346.
- 10 Chiu, Y.-L., and Rana, T. M. (2002). RNAi in Human Cells: Basic Structural and Functional Features of Small Interfering RNA. *Molecular Cell* 10, 549-561.
- Chiu, Y.L. & Rana, T.M. siRNA function in RNAi: a chemical modification analysis. *RNA* 9, 1034-1048 (2003).
- 15 Crow, J. P., Sampson, J. B., Zhuang, Y., Thompson, J. A., and Beckman, J. S. (1997). Decreased zinc affinity of amyotrophic lateral sclerosis-associated superoxide dismutase mutants leads to enhanced catalysis of tyrosine nitration by peroxynitrite. *J Neurochem* 69, 1936-1944.
- Djikeng, A., Shi, H., Tschudi, C., and Ullu, E. (2001). RNA interference in Trypanosoma brucei: cloning of small interfering RNAs provides evidence for retroposon-derived 24-26-nucleotide RNAs. *RNA* 7, 1522-1530.
- 20 D.H. Mathews, J. S., M. Zuker & D.H. Turner (1999). Expanded Sequence Dependence of Thermodynamic Parameters Improves Prediction of RNA Secondary Structure. *J Mol Biol* 288, 911-940.
- 25 Dalmay, T., Horsefield, R., Braunstein, T. H., and Baulcombe, D. C. (2001). SDE3 encodes an RNA helicase required for post-transcriptional gene silencing in Arabidopsis. *EMBO J* 20, 2069-2078.
- Ding, H. et al. Selective silencing by RNAi of a dominant allele that causes amyotrophic lateral sclerosis. *Aging Cell* 2, 209-217 (2003).
- 30 Doench, J. G., Petersen, C. P., and Sharp, P. A. (2003). siRNAs can function as miRNAs. *Genes Dev* 17, 438-442.

- Doench, J.G. & Sharp, P.A. Specificity of microRNA target selection in translational repression. *Genes Dev.* 18, 504–511 (2004).
- Doi, N., Zenno, S., Ueda, R., Ohki-Hamazaki, H., Ui-Tei, K., and Saigo, K. (2003). Short-Interfering-RNA-Mediated Gene Silencing in Mammalian Cells Requires
 5 Dicer and eIF2C Translation Initiation Factors. *Curr Biol* 13, 41-46.
- Dostie, J., Mourelatos, Z., Yang, M., Sharma, A., and Dreyfuss, G. (2003). Numerous microRNPs in neuronal cells containing novel microRNAs. *RNA.* 9: 180-6.
- Elbashir, S. M., Lendeckel, W., and Tuschl, T. (2001a). RNA interference is mediated by 21- and 22-nucleotide RNAs. *Genes Dev* 15, 188-200.
- 10 Elbashir, S. M., Martinez, J., Patkaniowska, A., Lendeckel, W., and Tuschl, T. (2001b). Functional anatomy of siRNAs for mediating efficient RNAi in *Drosophila melanogaster* embryo lysate. *EMBO J* 20, 6877-6888.
- Elbashir, S.M. et al. Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature* 411, 494–498 (2001).
- 15 Emery, J.F., Floyd, S.K., Alvarez, J., Eshed, Y., Hawker, N.P., Izhaki, A., Baum, S.F., and Bowman, J.L. (2003). Radial patterning of *Arabidopsis* shoots by class III HD-ZIP and KANADI genes. *Current Biology.* 13: 1768-1774.
- Enright, A. et al. MicroRNA targets in *Drosophila*. *Genome Biol.* 5, R1 (2003).
- Fire, A., Xu, S., Montgomery, M. K., Kostas, S. A., Driver, S. E., and Mello, C. C.
 20 (1998). Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 391, 806-811.
- Grad, Y., Aach, J., Hayes, G.D., Reinhart, B.J., Church, G.M., Ruvkun, G., *et. al.*, (2003). Computational and experimental identification of *C. elegans* microRNAs. *Molecular Cell.* 11(5): 1253-63.
- 25 Griffiths-Jones, S. (2004). The microRNA registry. *Nucleic Acids Research.* 32: D109-111.
- Grishok, A., Pasquinelli, A. E., Conte, D., Li, N., Parrish, S., Ha, I., Baillie, D. L., Fire, A., Ruvkun, G., and Mello, C. C. (2001). Genes and Mechanisms Related to
 30 RNA Interference Regulate Expression of the Small Temporal RNAs that Control *C. elegans* Developmental Timing. *Cell* 106, 23-34.
- Hake, S. (2003). MicroRNAs: a role in plant development. *Current Biology.* 13(21): R851-2.

- Haley, B., Tang, G. & Zamore, P.D. In vitro analysis of RNA interference in *Drosophila melanogaster*. *Methods* 30, 330–336 (2003).
- Hamilton, A. J., and Baulcombe, D. C. (1999). A species of small antisense RNA in posttranscriptional gene silencing in plants. *Science* 286, 950-952.
- 5 Hammond, S. M., Bernstein, E., Beach, D., and Hannon, G. J. (2000). An RNA-directed nuclease mediates post-transcriptional gene silencing in *Drosophila* cells. *Nature* 404, 293-296.
- Hammond, S. M., Caudy, A. A., and Hannon, G. J. (2001). Post-transcriptional gene silencing by double-stranded RNA. *Nat Rev Genet* 2, 110-119.
- 10 Hannon, G.J. & Zamore, P.D. Small RNAs, *Big Biology: and Biochemical Studies of RNA Interference*. In *RNAi: A Guide To Gene Silencing* (ed. Hannon, G.J.) 87–108 (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY; 2003).
- Holen, T., Amarzguioui, M., Babaie, E. & Prydz, H. Similar behaviour of single-strand and double-strand siRNAs suggests they act through a common RNAi pathway. *Nucleic Acids Res* 31, 2401–2407 (2003).
- 15 Houbaviy, H.B., Murray, M.F., and Sharp, P.A. (2003). Embryonic stem cell-specific MicroRNAs. *Developmental Cell*. 5(2): 351-8.
- Hunter, C., and Poethig, R.S. (2003). miSSING LINKS: miRNAs and plant development. *Current Opinion in Genetics and Development*. 13(4): 372-8.
- 20 Hutvagner, G., McLachlan, J., Pasquinelli, A. E., Balint, É., Tuschl, T., and Zamore, P. D. (2001). A cellular function for the RNA-interference enzyme Dicer in the maturation of the *let-7* small temporal RNA. *Science* 293, 834-838.
- Hutvagner, G., and Zamore, P. D. (2002). A MicroRNA in a Multiple-Turnover RNAi Enzyme Complex. *Science* 297, 2056-2060.
- 25 Hutvagner, G., Simard, M.J., Mello, C.C. & Zamore, P.D. Sequence-specific inhibition of small RNA function. *PLoS Biology* 2, 1–11 (2004).
- Ishizuka, A., Siomi, M. C., and Siomi, H. (2002). A *Drosophila* fragile X protein interacts with components of RNAi and ribosomal proteins. *Genes Dev* 16, 2497-2508.
- 30 Jackson, A. L., Bartz, S. R., Schelter, J., Kobayashi, S. V., Burchard, J., Mao, M., Li, B.,

- Cavet, G., and Linsley, P. S. (2003). Expression profiling reveals off-target gene regulation by RNAi. *Nat Biotechnol* 21, 635-637.
- Johnston, R.J. & Hobert, O. A microRNA controlling left/right neuronal asymmetry in *Caenorhabditis elegans*. *Nature* 426, 845-849 (2003).
- 5 Kasschau, K. D., Xie, Z., Allen, E., Llave, C., Chapman, E. J., Krizan, K. A., and Carrington, J. C. (2003). P1/HC-Pro, a Viral Suppressor of RNA Silencing, Interferes with Arabidopsis Development and miRNA Function. *Dev Cell* 4, 205-217.
- 10 Kawasaki, H., and Taira, K. (2003). Hes1 is a target of microRNA-23 during retinoic-acid-induced neuronal differentiation of NT2 cells. *Nature*. 423(6942): 838-42.
- Kennerdell, J. R., Yamaguchi, S., and Carthew, R. W. (2002). RNAi is activated during *Drosophila* oocyte maturation in a manner dependent on aubergine and spindle-E. *Genes Dev* 16, 1884-1889.
- 15 Ketting, R. F., Fischer, S. E., Bernstein, E., Sijen, T., Hannon, G. J., and Plasterk, R. H. (2001). Dicer functions in RNA interference and in synthesis of small RNA involved in developmental timing in *C. elegans*. *Genes Dev* 15, 2654-2659.
- Khvorova, A., Reynolds, A. & Jayasena, S.D. Functional siRNAs and miRNAs exhibit strand bias. *Cell* 115, 209-216 (2003).
- 20 Kidner, C.A., and Martienssen, R.A. (2004). Spatially restricted microRNA directs leaf polarity through ARGONAUTE1. *Nature*. 428(6978): 81-4.
- Kidner, C.A., and Martienssen, R.A. (2003). Macro effects of microRNAs in plants. *Trends in Genetics*. 19(1): 13-6.
- 25 Kim, J., Krichevsky, A., Grad, Y., Hayes, G.D., Kosik, K.S., Church, G.M., and Ruvkun, G. (2004). Identification of many microRNAs that copurify with polyribosomes in mammalian neurons. *Proceedings of the National Academy of Sciences*. 101(1): 360-365.
- Knight, S. W., and Bass, B. L. (2001). A role for the RNase III enzyme DCR-1 in RNA interference and germ line development in *Caenorhabditis elegans*. *Science* 293, 2269-2271.
- 30 Krichevsky, A.M, King, K.S., Donahue, C.P., Khrapko, K., and Kosik, K.S. (2003). A

- microRNA array reveals extensive regulation of microRNAs during brain development. *RNA* 9(10): 1274-81.
- Lagos-Quintana, M., Rauhut, R., Lendeckel, W., and Tuschl, T. (2001). Identification of Novel Genes Coding for Small Expressed RNAs. *Science* 294, 853-858.
- 5 Lagos-Quintana, M., Rauhut, R., Yalcin, A., Meyer, J., Lendeckel, W., and Tuschl, T. (2002). Identification of Tissue-Specific MicroRNAs from Mouse. *Curr Biol* 12, 735-739.
- Lagos-Quintana, M., Rauhut, R., Meyer, J., Borkhardt, A., and Tuschl, T. (2003). New microRNAs from mouse and human. *RNA* 9, 175-179.
- 10 Lai, E.C., Tomancak, P., Williams, R.W., and Rubin, G.M. (2003). Computational identification of Drosophila microRNA genes. *Genome Biology* 4: R42, 1-20.
- Lai, E.C. Micro RNAs are complementary to 3' UTR sequence motifs that mediate
- 15 negative post-transcriptional regulation. *Nat Genet* 30, 363-364 (2002).
- Lau, N. C., Lim, L. P., Weinstein, E. G., and Bartel, D. P. (2001). An Abundant Class of Tiny RNAs with Probable Regulatory Roles in *Caenorhabditis elegans*. *Science* 294, 858-862.
- Lee, R. C., and Ambros, V. (2001). An Extensive Class of Small RNAs in
- 20 *Caenorhabditis elegans*. *Science* 294, 862-864.
- Lee, R. C., Feinbaum, R. L., and Ambros, V. (1993). The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell* 75, 843-854.
- Lewis, B., Shih, I., Jones-Rhoades, M., Bartel, D. & Burge, C. Prediction of mammalian
- 25 microRNA targets. *Cell* 115, 787-798 (2003).
- Lim, L. P., Glasner, M. E., Yekta, S., Burge, C. B., and Bartel, D. P. (2003a). Vertebrate microRNA genes. *Science* 299, 1540.
- Lim, L. P., Lau, N. C., Weinstein, E. G., Abdelhakim, A., Yekta, S., Rhoades, M. W., Burge, C. B., and Bartel, D. P. (2003b). The microRNAs of *Caenorhabditis*
- 30 *elegans*. *Genes Dev* 17, 991-1008.
- Lin, S. Y., Johnson, S. M., Abraham, M., Vella, M. C., Pasquinelli, A., Gamberi, C.,

- Gottlieb, E., and Slack, F. J. (2003). The *C. elegans* hunchback Homolog, *hbl-1*, Controls Temporal
- Lima, W.F. & Crooke, S.T. Binding affinity and specificity of *Escherichia coli* RNase H1: impact on the kinetics of catalysis of antisense oligonucleotide-RNA hybrids. *Biochemistry* 36, 390–398 (1997).
- 5 Llave, C., Kasschau, K. D., Rector, M. A., and Carrington, J. C. (2002a). Endogenous and Silencing-Associated Small RNAs in Plants. *Plant Cell* 14, 1605-1619.
- Llave, C., Xie, Z., Kasschau, K. D., and Carrington, J. C. (2002b). Cleavage of *Scarecrow-Like* mRNA Targets Directed by a Class of *Arabidopsis* miRNA.
- 10 Science 297, 2053-2056.
- Mallory, A.C., Dugas, D.V., Bartel, D.P., and Bartel, B. (2004). MicroRNA regulation of NAC-domain targets is required for proper formation and separation of adjacent embryonic, vegetative, and floral organs. *Current Biology*. 14(12): 1035-46.
- 15 Mallory, A. C., Reinhart, B. J., Bartel, D., Vance, V. B., and Bowman, L. H. (2002). A viral suppressor of RNA silencing differentially regulates the accumulation of short interfering RNAs and micro-RNAs in tobacco. *Proc Natl Acad Sci U S A* 99, 15228-15233.
- Martinez, J., Patkaniowska, A., H. H. U., Lührmann, R., and Tuschl, T. (2002). Single stranded antisense siRNA guide target RNA cleavage in RNAi. *Cell* 110, 563-574.
- 20 Martinez, J. & Tuschl, T. RISC is a 5' phosphomonoester-producing RNA endonuclease. *Genes Dev.* Published online, April 22, 2004, 10.1101/gad.1187904 (2004).
- 25 McManus, M. T., Petersen, C. P., Haines, B. B., Chen, J., and Sharp, P. A. (2002). Gene silencing using micro-RNA designed hairpins. *RNA* 8, 842-850.
- McManus, M.T. (2003). MicroRNAs and cancer. *Seminars in Cancer Biology*. 13(4): 253-8.
- 30 Meister, G., Landthaler, M., Dorsett, Y. & Tuschl, T. Sequence-specific inhibition of

- microRNA- and siRNA-induced RNA silencing. *RNA* 10, 544–550 (2004).
- Michael, M.Z., O'Connor, S.M., van Holst Pellekaan, N.G., Young, G.P., and James, R.J. (2003). Reduced accumulation of specific microRNAs in colorectal neoplasia. *Molecular Cancer Research*. 1(12): 882-91.
- 5 Myers, J. W., Jones, J. T., Meyer, T., and Ferrell, J. E. (2003). Recombinant Dicer efficiently converts large dsRNAs into siRNAs suitable for gene silencing. *Nat Biotechnol*.
- Mourelatos, Z., Dostie, J., Paushkin, S., Sharma, A. K., Charroux, B., Abel, L., Rappsilber, J., Mann, M., and Dreyfuss, G. (2002). miRNPs: a novel class of
 10 Ribonucleoproteins containing numerous microRNAs. *Genes Dev* 16, 720-728.
- Nykänen, A., Haley, B., and Zamore, P. D. (2001). ATP Requirements and Small Interfering RNA Structure in the RNA Interference Pathway. *Cell* 107, 309-321.
- Olsen, P. H., and Ambros, V. (1999). The lin-4 regulatory RNA controls developmental timing in *Caenorhabditis elegans* by blocking LIN-14 protein synthesis after the
 15 initiation of translation. *Dev Biol* 216, 671-680.
- Paddison, P. J., Caudy, A. A., Bernstein, E., Hannon, G. J., and Conklin, D. S. (2002). Short hairpin RNAs (shRNAs) induce sequence-specific silencing in mammalian cells. *Genes Dev* 16, 948-958.
- Palatnik, J.F., Allen, E., Wu, X., Schommer, C., Scwab, R., Carrington, J.C., and
 20 Weigel, D. (2003). Control of leaf morphogenesis by microRNAs. *Nature*. 425: 257-263.
- Park, W., Li, J., Song, R., Messing, J., and Chen, X. (2002). CARPEL FACTORY, a Dicer Homolog, and HEN1, a Novel Protein, Act in microRNA Metabolism in *Arabidopsis thaliana*. *Curr Biol* 12, 1484–1495.
- 25 Pasquinelli, A.E., and Ruvkun, G. (2002). Control of developmental timing by microRNAs and their targets. *Annual Reviews in Cellular and Developmental Biology*. 18: 495-513.
- Pasquinelli, A. E., Reinhart, B. J., Slack, F., Martindale, M. Q., Kuroda, M. I., Maller, B., Hayward, D. C., Ball, E. E., Degan, B., Muller, P., *et al.* (2000).
 30 Conservation of the sequence and temporal expression of let-7 heterochronic regulatory RNA. *Nature* 408, 86-89.

- Paul, C. P., Good, P. D., Winer, I., and Engelke, D. R. (2002). Effective expression of small interfering RNA in human cells. *Nat Biotechnol* 20, 505-508.
- Pham, J.W., Pellino, J.L., Lee, Y.S., Carthew, R.W. & Sontheimer, E.J. A Dicer-2-dependent 80S complex cleaves targeted mRNAs during RNAi in *Drosophila*.
5 Cell 117, 83-94 (2004).
- Phipps, K.M., Martinez, A., Lu, J., Heinz, B.A. & Zhao, G. Small interfering RNA molecules as potential anti-human rhinovirus agents: in vitro potency, specificity, and mechanism. *Antiviral Res* 61, 49-55 (2004).
- Provost, P., Dishart, D., Doucet, J., Friendewey, D., Samuelsson, B., and Radmark, O.
10 (2002). Ribonuclease activity and RNA binding of recombinant human Dicer. *EMBO J* 21, 5864-5874.
- Pusch, O., Boden, D., Silberman, R., Lee, F., Tucker, L., and Ramratnam, B. (2003). Nucleotide sequence homology requirements of HV-1-specific short hairpin RNA. *Nucleic Acids Research*. 31: 6444-9.
- 15 Rajewsky, N. & Socci, N.D. Computational identification of microRNA targets. *Dev. Biol.* 267, 529-535 (2004).
- Reinhart, B.J., and Bartel, D.P. (2002). Small RNAs correspond to centromere heterochromatic repeats. *Science*. 297: 1831.
- 20 Reinhart, B. J., Slack, F. J., Basson, M., Pasquinelli, A. E., Bettinger, J. C., Rougvie, A. E., Horvitz, H. R., and Ruvkun, G. (2000). The 21-nucleotide let-7 RNA regulates developmental timing in *Caenorhabditis elegans*. *Nature* 403, 901-906.
- Reinhart, B. J., Weinstein, E. G., Rhoades, M. W., Bartel, B., and Bartel, D. P. (2002). MicroRNAs in plants. *Genes Dev* 16, 1616-1626.
- 25 Reynolds, A. et al. Rational siRNA design for RNA interference. *Nature Biotechnology* 22, 326-330 (2004).
- Rhoades, M. W., Reinhart, B. J., Lim, L. P., Burge, C. B., Bartel, B., and Bartel, D. P. (2002). Prediction of Plant MicroRNA Targets. *Cell* 110, 513-520.
- Ruvkun, G. (2001). Molecular biology. Glimpses of a tiny RNA world. *Science* 294,
30 797-799.

- Schwarz, D.S. and Zamore, P.D. (2002). Why do miRNAs live in the miRNP? *Genes & Development*. 16: 1025-1031.
- Schwarz, D. S., Hutvagner, G., Haley, B., and Zamore, P. D. (2002). Evidence that siRNAs function as guides, not primers, in the *Drosophila* and human RNAi pathways. *Molecular Cell* 10, 537-548.
- Schwarz, D.S. et al. Asymmetry in the assembly of the RNAi enzyme complex. *Cell* 115, 199–208 (2003).
- Schwarz, D.S., Tomari, Y. & Zamore, P.D. The RNA-induced silencing complex is a Mg²⁺-dependent endonuclease. *Curr. Biol.* 14, 787–791 (2004).
- Semizarov, D., Frost, L., Sarthy, A., Kroeger, P., Halbert, D.N., and Fesik, S.W. (2003). Specificity of short interfering RNA determined through gene expression signatures. *Proceedings of the National Academy of Sciences USA*. 100(11): 6347-6342.
- Sempere, L.F., Freemantle, S., Pitha-Rowe, I., Moss, E., Dmitrovsky, E., and Ambros, V. (2004). Expression profiling of mammalian microRNAs uncovers a subset of brain-expressed microRNAs with possible roles in murine and human neuronal differentiation. *Genome Biology*. 5: R13
- Sijen, T., Fleenor, J., Simmer, F., Thijssen, K. L., Parrish, S., Timmons, L., Plasterk, R. H., and Fire, A. (2001). On the Role of RNA Amplification in dsRNA-Triggered Gene Silencing. *Cell* 107, 465-476.
- Silhavy, D., Molnar, A., Lucioli, A., Szitty, G., Hornyik, C., Tavazza, M., and Burgyan, J. (2002). A viral protein suppresses RNA silencing and binds silencing-generated, 21- to 25-nucleotide double-stranded RNAs. *EMBO J* 21, 3070-3080.
- Slack, F. J., Basson, M., Liu, Z., Ambros, V., Horvitz, H. R., and Ruvkun, G. (2000). The *lin-41* RBCC gene acts in the *C. elegans* heterochronic pathway between the *let-7* regulatory RNA and the LIN-29 transcription factor. *Mol Cell* 5, 659-669.
- Stark, A., Brennecke, J., Russel, R. & Cohen, S. Identification of *Drosophila* microRNA targets. *PLoS Biology* 1, 1–13 (2003).
- Stryer, L. *Biochemistry*. (W. H. Freeman and Company, San Francisco; 1981).
- Tabara, H., Yigit, E., Siomi, H., and Mello, C. C. (2002). The dsRNA Binding Protein

- RDE-4 Interacts with RDE-1, DCR-1, and a DexH-Box Helicase to Direct RNAi in *C. elegans*. *Cell* 109, 861-871.
- Takamizawa, J., Konishi, H., Yanagisawa, K., Tomida, S., Osada, H., Endoh, H., Harano, T., Yatabe, Y., Nagino, M., Nimura, Y., Mitsudomi, T., and Takahashi, T. (2004). Reduced expression of the let-7 MicroRNAs in Human Lung Cancers in Association with Shortened Postoperative Survival. *Cancer Research*. 64: 3753-6.
- Tang, G., Reinhart, B. J., Bartel, D. P., and Zamore, P. D. (2003). A biochemical framework for RNA silencing in plants. *Genes Dev* 17, 49-63.
- 10 Tijsterman, M., Ketting, R. F., Okihara, K. L., Sijen, T., and Plasterk, R. H. (2002). RNA helicase MUT-14-dependent gene silencing triggered in *C. elegans* by short antisense RNAs. *Science* 295, 694-697.
- Tomari, Y. et al. RISC assembly defects in the *Drosophila* RNAi mutant armitage. *Cell* 116, 831-841 (2004).
- 15 Tuschl, T., Zamore, P. D., Lehmann, R., Bartel, D. P., and Sharp, P. A. (1999). Targeted mRNA degradation by double-stranded RNA in vitro. *Genes Dev* 13, 3191-3197.
- Vaucheret, H., Vazques, F., Crata, P., and Bartel, D.P. (2004). The action of ARGONAUTE1 in the miRNA pathway and its regulation by the miRNA pathway are crucial for plant development. *Genes & Development*. 18(10):1187-97.
- 20 Vella, M., Choi, E., Lin, S., Reinert, K. & Slack, F. The *C. elegans* microRNA let-7 binds to imperfect let-7 complementary sites from the lin-41 3'UTR. *Genes Dev*. 18, 132-137 (2004).
- Voet, D. & Voet, J.G. *Biochemistry*. (John Wiley & Sons, Hoboken, NJ; 2004).
- 25 Wightman, B., Ha, I., and Ruvkun, G. (1993). Posttranscriptional regulation of the heterochronic gene lin-14 by lin-4 mediates temporal pattern formation in *C. elegans*. *Cell* 75, 855-862.
- Wu, H., Lima, W.F. & Crooke, S.T. Investigating the structure of human RNase H1 by site-directed mutagenesis. *J. Biol. Chem*. 276, 23547-23553. (2001).
- 30 Wu-Scharf, D., Jeong, B., Zhang, C., and Cerutti, H. (2000). Transgene and transposon silencing in *chlamydomonas reinhardtii* by a DEAH-Box RNA helicase. *Science* 290, 1159-1163.

- Xie, Z., Kasschau, K. D., and Carrington, J. C. (2003). Negative Feedback Regulation of Dicer-Like1 in Arabidopsis by microRNA-Guided mRNA Degradation. *Curr Biol* 13, 784-789.
- 5 Xu, P., Vernooy, S. Y., Guo, M., and Hay, B. A. (2003). The Drosophila MicroRNA Mir-14 Suppresses Cell Death and Is Required for Normal Fat Metabolism. *Curr Biol* 13, 790-795.
- Yekta, S., Shih, I., and Bartel, D.P. (2004). MicroRNA-Directed Cleavage of *HOXB8* mRNA. *Science* 304: 594-6.
- 10 Yu, J. Y., DeRuijter, S. L., and Turner, D. L. (2002). RNA interference by expression of short-interfering RNAs and hairpin RNAs in mammalian cells. *Proc Natl Acad Sci U S A* 99, 6047-6052.
- Zamore, P. D., Tuschl, T., Sharp, P. A., and Bartel, D. P. (2000). RNAi: double-stranded RNA directs the ATP-dependent cleavage of mRNA at 21 to 23 nucleotide intervals. *Cell* 101, 25-33.
- 15 Zeng, Y., and Cullen, B.R. (2003). Sequence requirements for micro RNA processing and function in human cells. *RNA*. 9: 112-123.
- Zeng, Y., Yi, R., and Cullen, B.R. (2003). MicroRNAs and small interfering RNAs can inhibit mRNA expression by similar mechanisms. *Proceedings of the National Academy of Sciences*. 100(17): 9779-9784.
- 20 Zeng, Y., Wagner, E.J., and Cullen, B.R. (2002). Both natural and designed microRNAs can inhibit the expression of cognate mRNAs when expressed in human cells. *Molecular Cell*. 9: 1327-33.
- 25 Zhang, H., Kolb, F. A., Brondani, V., Billy, E., and Filipowicz, W. (2002). Human Dicer preferentially cleaves dsRNAs at their termini without a requirement for ATP. *EMBO J* 21, 5875-5885.
- Zhong, R., and Ye, Z.H. (2004). Amphivasal vascular bundle 1, a gain-of-function mutation of the *IFL1/REV* gene, is associated with alterations in the polarity of leaves, stems, and carpels. *Plant Cell Physiology*. 45(4): 369-385.
- 30

Zuker, M. (2003). Mfold web server for nucleic acid folding and hybridization prediction. Nucleic Acids Res 31, 1-10.

Equivalents

- 5 Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

What is claimed:

1. A method of enhancing the ability of a first strand of a siRNA-like duplex to act in mediating RNA silencing, comprising lessening the base pair strength
5 between the 5' end of the first strand and the 3' end of a second strand of the duplex as compared to the base pair strength between the 3' end of the first strand and the 5' end of the second strand.
2. A method of enhancing the efficacy of a siRNA-like duplex, the siRNA-
10 like duplex comprising a miRNA and an miRNA* strand, comprising lessening the base pair strength between the miRNA strand 5' end (miRNA 5') and the miRNA* strand 3' end (miRNA* 3') as compared to the base pair strength between the miRNA strand 3' end (miRNA 3') and the miRNA* strand 5' end (miRNA* 5'), such that efficacy is enhanced.
- 15 3. A method for promoting entry of a desired strand of an siRNA-like duplex into a RISC complex, comprising enhancing the asymmetry of the siRNA duplex, such that entry of the desired strand is promoted.
- 20 4. The method of claim 3, wherein the asymmetry is enhanced by lessening the base pair strength between the 5' end of the desired strand and the 3' end of a complementary strand of the duplex as compared to the base pair strength between the 3' end of the desired strand and the 5' end of the complementary strand.
- 25 5. A siRNA-like duplex comprising a miRNA strand and an miRNA* strand, wherein the base pair strength between the miRNA strand 5' end (miRNA 5') and the miRNA* strand 3' end (miRNA* 3') is less than the base pair strength between the miRNA strand 3' end (miRNA 3') and the miRNA* strand 5' end (miRNA* 5'), such that the miRNA strand preferentially silences a target mRNA.
- 30 6. The duplex of claim 5, wherein the base-pair strength is less due to fewer G:C base pairs between the 5' end of the first or miRNA strand and the 3' end of the

second or miRNA* strand than between the 3' end of the first or miRNA strand and the 5' end of the second or miRNA* strand.

7. The duplex of claim 5, wherein the base pair strength is less due to at least one mismatched base pair between the 5' end of the first or miRNA strand and the 3' end of the second or miRNA* strand.

8. The duplex of claim 5, wherein the base pair strength is less due to at least one wobble base pair, e.g., G:U, between the 5' end of the first or miRNA strand and the 3' end of the second or miRNA* strand.

9. The duplex of claim 5, wherein the base pair strength is less due to at least one base pair comprising a rare nucleotide.

10. The duplex of claim 9, wherein the rare nucleotide is inosine (I).

11. The duplex of claim 5, wherein the base pair strength is less due to at least one base pair comprising a modified nucleotide.

12. The duplex of claim 11, wherein the modified nucleotide is selected from the group consisting of 2-amino-G, 2-amino-A, 2,6-diamino-G, and 2,6-diamino-A.

13. A composition comprising the siRNA-like duplex of any one of claims 5-12 formulated to facilitate entry of the siRNA-like duplex into a cell.

14. A pharmaceutical composition comprising the siRNA-like duplex of any one of claims 5-12.

15. An engineered pre-miRNA comprising the siRNA-like duplex of any one of claims 5-12.

16. A vector encoding the pre-miRNA of claim 15.

17. A pri-miRNA comprising the pre-miRNA of claim 15.
18. A vector encoding the pri-miRNA of claim 17.
- 5 19. A small hairpin RNA (shRNA) comprising nucleotide sequence identical to the miRNA and miRNA* strand of the siRNA-like duplex of any one of claims 5-12.
20. The shRNA of claim 19, wherein the nucleotide sequence identical to the miRNA strand is upstream of the nucleotide sequence identical to the miRNA* strand.
- 10 21. The shRNA of claim 19, wherein the nucleotide sequence identical to the miRNA* strand is upstream of the nucleotide sequence identical to the miRNA strand.
22. A vector encoding the shRNAs of any one of claims 19-21.
- 15 23. A transgene encoding the shRNAs of any one of claims 19-21.
24. A cell comprising the vector of any one of claims 16, 18 and 22.
- 20 25. The cell of claim 24 which is a mammalian cell.
26. The cell of claim 24 which is a human cell.
27. A method of enhancing silencing of a target mRNA, comprising
- 25 contacting a cell having an RNA silencing pathway with an RNA silencing produced according to the methods of any one of claims 1-4, under conditions such that silencing is enhanced.
28. A method of enhancing silencing of a target mRNA in a subject,
- 30 comprising administering to the subject a pharmaceutical composition comprising a RNA silencing agent produced according to the methods of any one of claims 1-4, such that silencing is enhanced.

29. A method of decreasing silencing of an inadvertant target mRNA by a dsRNA silencing agent, the dsRNA silencing agent (e.g., a siRNA-like duplex) comprising a miRNA strand and a miRNA* strand comprising the steps of: (a) detecting a significant degree of complementarity between the miRNA* strand and the inadvertant target; and (b) enhancing the base pair strength between the 5' end of the miRNA* strand and the 3' end of the miRNA strand relative to the base pair strength between the 3' end of the miRNA* strand and the 5' end of the miRNA strand; such that silencing of the inadvertant target mRNA is decreased.

30. The method of claim 29, wherein the silencing of the inadvertant target mRNA is decreased relative to silencing of a desired target mRNA.

31. A method of treating a diseased cell characterized by an miRNA disorder, comprising administering an effective amount of an RNAi agent comprising a sense strand and an antisense strand, wherein entry of the antisense strand into a RISC complex is promoted, such that the disorder is treated.

32. The method of claim 31, wherein the entry of the antisense strand is promoted by lessening the base pair strength between the 5' end of the antisense strand (AS 5') and the 3' end of the sense strand (S 3') as compared to the base pair strength between the 3' end of the antisense strand (AS 3') and the 5' end of the sense strand (S 5').

33. The method of claim 32, wherein the base pair strength between the AS 5' and the S 3' is lessened by introducing fewer G:C base pairs between the AS 5' and the S 3' than between the AS 3' and the S 5'.

34. The method of claim 32, wherein the base pair strength is lessened due to at least one mismatched base pair between the AS 5' and the S 3'.

35. The method of claim 34, wherein the mismatched base pair is selected from the group consisting of G:A, C:A, C:T, C:U, G:G, A:A, C:C, U:T, and U:U.

36. The method of claim 32, wherein the base pair strength is less due to at least one wobble base pair between the AS 5' and the S 3'.

37. The method of claim 36, wherein the wobble base pair is selected from the group consisting of a G:U or G:T.

38. The method of claim 32, wherein the base pair strength is less due to at least one base pair comprising a rare nucleotide.

39. The method of claim 38, wherein the rare nucleotide is inosine (I).

40. The method of claim 38, wherein the base pair is selected from the group consisting of an I:A, I:U and I:C.

41. The method of claim 32, wherein the base pair strength is less due to at least one base pair comprising a modified nucleotide.

42. The method of claim 41, wherein the modified nucleotide is selected from the group consisting of 2-amino-G, 2-amino-A, 2,6-diamino-G, and 2,6-diamino-A.

43. The method of claim 31, wherein the entry of the antisense strand is promoted by enhancing the base pair strength between the 3' end of the antisense strand (AS 3') and the 5' end of the sense strand (S 5') as compared to the base pair strength between the 5' end of antisense strand (AS 5') and the 3' end of the sense strand (S 3').

44. The method of claim 43, wherein the base pair strength between the AS 3' and the S 5' is enhanced by introducing more G:C base pairs between the AS 3' and the S 5' than between the AS 5' and the S 3'.

45. The method of claim 31, wherein the diseased cell is further characterized by the overexpression of an mRNA.

46. The method of claim 45, wherein the antisense strand preferentially guides cleavage of the mRNA in a normal cell.

47. The method of claim 45, wherein the mRNA contains a mutation that suppresses its ability to be cleaved by the antisense strand of a natural miRNA.

48. The method of claim 31, wherein the diseased cell is further characterized by the underexpression of an mRNA.

49. The method of claim 45 or 48, wherein the diseased cell is further
5 characterized by the overexpression of the sense strand of a natural miRNA.

50. The method of claim 49, wherein the sense strand of the natural miRNA preferentially guides cleavage of the mRNA in the diseased cell.

10 51. The method of claim 45 or 48, wherein the disease is further characterized by the underexpression of the antisense strand of a natural miRNA.

52. The method of claim 51, wherein the antisense strand of the natural miRNA preferentially guides cleavage of the mRNA in the diseased cell.
15

53. The method of claim 48, wherein the mRNA contains a mutation that promotes its ability to be cleaved by the antisense strand of a natural miRNA.

54. The method of claim 31, wherein the sequence of the antisense strand of
20 the RNAi agent is homologous to the sequence of the antisense strand of a natural miRNA.

55. The method of claim 54, wherein the sequence of the antisense strand of the RNAi agent is identical to the sequence of the antisense strand of a natural miRNA.
25

56. The method of claim 54, wherein the miRNA disorder is characterized by a mutation within the nucleotide sequence of the natural miRNA that promotes entry of a complementary sense strand of the natural miRNA into a RISC complex.

30 57. The method of claim 56, wherein the mutation lessens the base pair strength between the 5' end of sense strand (S 5') and the 3' end of the antisense strand (AS 3') as compared to the base pair strength between the 5' end of the antisense strand (AS 5') and the 3' end of the sense strand (S '3).

58. The method of claim 56, wherein the mutation enhances the base pair strength between the 5' end of antisense strand (AS 5') and the 3' end of the sense strand (S 3') as compared to the base pair strength between the 3' end of the antisense strand (AS 3') and the 5' end of the sense strand (S 5').

59. The method of claim 31, wherein the RNAi agent is a siRNA duplex.

60. The method of claim 59, wherein the siRNA duplex is chemically synthesized.

61. The method of claim 59, wherein the siRNA is enzymatically synthesized.

62. The method of claim 59, wherein the siRNA is derived from an engineered precursor.

63. The method of claim 59, wherein the siRNA is comprised of modified nucleotides.

64. The method of claim 31, wherein the cell in a subject suffering from a disease characterized by a miRNA disorder.

65. The method of claim 64, wherein the subject is a human.

66. A method of treating a diseased cell characterized by an miRNA disorder, comprising administering an effective amount of an RNAi agent comprising a sense strand and an antisense strand, wherein entry of the sense strand into a RISC complex is promoted, such that the disorder is treated.

67. The method of claim 66, wherein the entry of the sense strand is promoted by lessening the base pair strength between the 5' end of the sense strand (S 5') and the 3' end of the antisense strand (AS 3') as compared to the base pair strength

between the 3' end of the sense strand (S 3') and the 5' end of the antisense strand (AS 5').

68. The method of claim 67, wherein the base pair strength between the S 5' and the AS 3' is lessened by introducing fewer G:C base pairs between the S 5' and the AS 3' than between the S 3' and the AS 5'.

69. The method of claim 67, wherein the base pair strength is lessened due to at least one mismatched base pair between the S 5' and the AS 3'.

70. The method of claim 69, wherein the mismatched base pair is selected from the group consisting of G:A, C:A, C:T, C:U, G:G, A:A, C:C, U:T, and U:U.

71. The method of claim 67, wherein the base pair strength is less due to at least one wobble base pair between the S 5' and the AS 3'.

72. The method of claim 71, wherein the wobble base pair is selected from the group consisting of a G:U or G:T.

73. The method of claim 67, wherein the base pair strength is less due to at least one base pair comprising a rare nucleotide.

74. The method of claim 73, wherein the rare nucleotide is inosine (I).

75. The method of claim 73, wherein the base pair is selected from the group consisting of an I:A, I:U and I:C.

76. The method of claim 67, wherein the base pair strength is less due to at least one base pair comprising a modified nucleotide.

77. The method of claim 76, wherein the modified nucleotide is selected from the group consisting of 2-amino-G, 2-amino-A, 2,6-diamino-G, and 2,6-diamino-A.

78. The method of claim 66, wherein the entry of the sense strand is promoted by enhancing the base pair strength between the 3' end of the sense strand (S

3') and the 5' end of the antisense strand (AS '5) as compared to the base pair strength between the

5' end of sense strand (S 5') and the 3' end of the antisense strand (AS 3').

5 79. The method of claim 78, wherein the base pair strength between the S 3' and the AS 5' is enhanced by introducing more G:C base pairs between the S 3' and the AS 5' than between the S 5' and the AS 3'.

 80. The method of claim 67, wherein the disease is further characterized by
10 the overexpression of an mRNA.

 81. The method of claim 80, wherein the sense strand preferentially guides cleavage of the mRNA in a normal cell.

15 82. The method of claim 80, wherein the mRNA contains a mutation that suppresses its ability to be cleaved by the sense strand of a natural miRNA.

 83. The method of claim 66, wherein the diseased cell is further characterized by the underexpression of an mRNA.

20

 84. The method of claim 81 or 83, wherein the diseased cell is further characterized by the overexpression of the antisense strand of a natural miRNA.

 85. The method of claim 80, wherein the antisense strand of the natural
25 miRNA preferentially guides cleavage of the mRNA in the diseased cell.

 86. The method of claim 81 or 83, wherein the diseased cell is further characterized by the underexpression of the sense strand of a natural miRNA.

30 87. The method of claim 86, wherein the sense strand of the natural miRNA preferentially guides cleavage of the mRNA in the diseased cell.

 88. The method of claim 84, wherein the mRNA contains a mutation that promotes its ability to be cleaved by the sense strand of a natural miRNA.

35

89. The method of claim 66, wherein the sequence of the sense strand of the RNAi agent is homologous to the sequence of the sense strand of a natural miRNA.

90. The method of claim 89, wherein the sequence of the sense strand of the RNAi agent is identical to the sequence of the sense strand of a natural miRNA.

91. The method of claim 89, wherein the miRNA disorder is characterized by a mutation within the nucleotide sequence of the natural miRNA that promotes entry of a complementary antisense strand of the natural miRNA into a RISC complex.

92. The method of claim 91, wherein the mutation lessens the base pair strength between the 5' end of antisense strand (AS 5') and the 3' end of the sense strand (S 3') as compared to the base pair strength between the 5' end of the sense strand (S 5') and the 3' end of the antisense strand (AS 3').

93. The method of claim 91, wherein the mutation enhances the base pair strength between the 5' end of sense strand (S 5') and the 3' end of the antisense strand (AS 3') as compared to the base pair strength between the 3' end of the sense strand (S 3') and the 5' end of the antisense strand (AS 5').

94. The method of claim 66, wherein the RNAi agent is a siRNA duplex.

95. The method of claim 94, wherein the siRNA duplex is chemically synthesized.

96. The method of claim 94, wherein the siRNA is enzymatically synthesized.

97. The method of claim 94, wherein the siRNA is derived from an engineered precursor.

98. The method of claim 94, wherein the siRNA is comprised of modified nucleotides.

99. The method of claim 66, wherein the cell is present in a subject suffering from a disease characterized by a miRNA disorder.

5 100. The method of claim 99, wherein the subject is a human.


1/33

(SEQ ID
Pp luc sense target: 5' -...cgaggugaacaucacguacgcggaauacuucgaaaugucc...-3' (●) NO.: 1)
Pp luc anti-sense target: 3' -...gcuccacuuguagugcaugcgccuuaugaagcuuuacagg...-5' (○) (SEQ ID
 NO.: 2)

Fig. 1A



(SEQ ID
 5' -CGUACGCGGAAUACUUCGAAA-3' (○) NO.: 3)

 3' -GUGCAUGCGCCUUAUGAAGCU-5' (●) (SEQ ID
 NO.: 4)

*Fig. 1B*


(SEQ ID
 5' -pCGUACGCGGAAUACUUCGAAA-3' (○) NO.: 3)

 5' -pUCGAAGUAUCCGCGUACGUG-3' (●) (SEQ ID
 NO.: 4)



*Fig. 1C**Fig. 1D*

(SEQ ID
 5' -UGUACGCGGAAUACUUCGAAA-3' (○) NO.: 5)

 3' -GUGCAUGCGCCUUAUGAAGCU-5' (●) (SEQ ID
 NO.: 4)

*Fig. 1E*

(SEQ ID
Pp luc sense target: 5' -...cgaggugaacaucacguacgcggaauacuucgaaaugucc...-3' (●) NO.: 6)
Pp luc anti-sense target: 3' -...gcuccacuuguagugcaugcgccuuaugaagcuuuacagg...-5' (○) (SEQ ID
 NO.: 7)

Fig. 2A*Fig. 2B**Fig. 2C**Fig. 2D**Fig. 2E*



3/33

(SEQ ID
sod1 sense target: 5' -...agagaggcauguuggagacuugggcaaugugacugcugacaa...3' (●) NO.: 11)
sod1 anti-sense target: 3' -...cuccguacaaccucugaacccguuacacugacgacuguuuc...5' (○) (SEQ ID
 NO.: 12)

Fig. 3A



(SEQ ID (SEQ ID
 5' -GAGACUUGGGCAAUGUGACdTdT-3' (○) NO.: 13) 5' -GAGACUUGGGCAAUGUGAAdTdT-3' (○) NO.: 15)

 3' -dTdTTCUCUGAACCCGUUACACUG-5' (●) (SEQ ID NO.: 14) 3' -dTdTTCUCUGAACCCGUUACACUG-5' (●) (SEQ ID
 NO.: 14)

*Fig. 3B**Fig. 3C*

(SEQ ID (SEQ ID
 5' -GAGACUUGGGCAAUGUGACdTdT-3' (○) NO.: 13) 5' -GAGACUUGGGCAAUGUGAAdTdT-3' (○) NO.: 15)




 3' -dTdTTCUCUGAACCCGUUACACUU-5' (●) (SEQ ID NO.: 16) 3' -dTdTTCUCUGAACCCGUUACACUU-5' (●) (SEQ ID
 NO.: 16)

*Fig. 3D**Fig. 3E*

4/33

(SEQ ID NO.: 13) 5' -GAGACUUGGGCAAUGUGACdTdT-3' (O) (SEQ ID NO.: 18) 5' -UAGACUUGGGCAAUGUGACdTdT-3' (O)



 3' -dTdT TAUCUGAACCCGUUACACUG-5' (●) (SEQ ID NO.: 17) 3' -dTdT TCUCUGAACCCGUUACACUG-5' (●) (SEQ ID NO.: 14)

*Fig. 3F**Fig. 3G**Fig. 3H**Fig. 3I*

5/33



(SEQ ID NO.: 19) 5' -IAGACUUGGGCAAUGUGACdTdT-3' (O) (SEQ ID NO.: 13) 5' -GAGACUUGGGCAAUGUGACdTdT-3' (O)

 (SEQ ID NO.: 14) 3' -dTdTTCUCUGAACCCGUUACACUG-5' (●) (SEQ ID NO.: 20) 3' -dTdTTCUCUGAACCCGUUACACUI-5' (●)

*Fig. 4A**Fig. 4B*

(SEQ ID NO.: 19) 5' -IAGACUUGGGCAAUGUGACdTdT-3' (O) (SEQ ID NO.: 19) 5' -IAGACUUGGGCAAUGUGACdTdT-3' (O)

 (SEQ ID NO.: 20) 3' -dTdTTCUCUGAACCCGUUACACUI-5' (●) (SEQ ID NO.: 20) 5' -IUCACAUUGCCCAAGUCUCdTdT-3' (●)

*Fig. 4C**Fig. 4D*




Fig. 5A







Fig. 5C




miR-9	5' - UCUUGGUUAUCUAGCUGUAUGA-3' (SEQ ID NO.: 47) 3' - AUUGAAGCAUUCGAUCGAAUA-5' (SEQ ID NO.: 48)	miR-1	5' - CCAUGCUUCUUGCAUUAUA-3' (SEQ ID NO.: 24) 3' - GAGUAUGAAGAAUUGUAGGU-5' (SEQ ID NO.: 25)
miR-10	5' - ACC CUGUAGAUCGAAUUUGU-3' (SEQ ID NO.: 49) 3' - UUGGAGAGAUUUUGGUUAAA-5' (SEQ ID NO.: 50)	miR-2a-1	5' - UC-UCAAAG-UGGUUGUAGAAUG-3' (SEQ ID NO.: 26) 3' - CGAGUAGUUUCGACCGACACUAU-5' (SEQ ID NO.: 27)
miR-11	5' - AAGAACUUUCUCUGUGACCGG-3' (SEQ ID NO.: 51) 3' - CGUUCUUGAGUCUGACAGUAC-5' (SEQ ID NO.: 52)	miR-2a-2	5' - CUCAUCAAG--UGGUUGUAGUAG-3' (SEQ ID NO.: 28) 3' - CGAGUAGUUUCGACCGACACUAU-5' (SEQ ID NO.: 27)
miR-12	5' - UGAGUAUAUCAUGAGUACUGGU-3' (SEQ ID NO.: 53) 3' - CGGAUUAUACUGUA-UUCAUGAC-5' (SEQ ID NO.: 54)	miR-2b-1	5' - UCUUCAAG-UGGCAGUGACAUAG-3' (SEQ ID NO.: 29) 3' - CGAGGAGUUUCGACCGACACUAU-5' (SEQ ID NO.: 30)
miR-13a	5' - UCCUCAAG-GGUUGUGAAUGUC-3' (SEQ ID NO.: 55) 3' - UGAGUAGUUUACCGACACUAU-5' (SEQ ID NO.: 56)	miR-2b-2	5' - UCUUCAAG-UGGUUGUAGAAUG-3' (SEQ ID NO.: 31) 3' - CGAGGAGUUUCGACCGACACUAU-5' (SEQ ID NO.: 30)
miR-13b-1	5' - UCGUUAUUAUUUGUAGCUUAUG-3' (SEQ ID NO.: 57) 3' - UGAGCAGUUUACCGACACUAU-3' (SEQ ID NO.: 58)	miR-3	5' - GGAUGCAUCUUGUGCAGUUAUG-3' (SEQ ID NO.: 32) 3' - ACUCUGUGAGAAACGGGUCACU-3' (SEQ ID NO.: 33)
miR-13b-2	5' - GCGUCAAAUAGACUGUGAGCUA-3' (SEQ ID NO.: 59) 3' - UGAGCAGUUUACCGACACUAU-5' (SEQ ID NO.: 60)	miR-4	5' - CUUUGGUGCUGCCUUAUGU-3' (SEQ ID NO.: 34) 3' - AGUUAACCAACAGAUCAUAUA-5' (SEQ ID NO.: 35)
miR-14	5' - GGAGCGAGACGGGACUCUAU-3' (SEQ ID NO.: 61) 3' - AUCCUCUCUCUUUUUUCUGACU-5' (SEQ ID NO.: 62)	miR-5	5' - AAAGGAACGAUCGUUGUAGUAGU-3' (SEQ ID NO.: 36) 3' - UAUUUCUUU-UUAGUGACACUAU-5' (SEQ ID NO.: 37)
miR-124a	5' - GUAUCCACUUGAGGCCUUAUUAUG-3' (SEQ ID NO.: 63) 3' - ACCGUUAGUGGCGCACGGAAU-5' (SEQ ID NO.: 64)	miR-6-1	5' - AGGGAUAGUUGCUGUGGUAU-3' (SEQ ID NO.: 38) 3' - UUUUUUCUUGCUGGACACUAU-5' (SEQ ID NO.: 39)
miR-125b (lin-4)	5' - UCCUGAGACCCUAAAUUGUGA-3' (SEQ ID NO.: 65) 3' - UUAUGGCCUUAUUUUGAACA-5' (SEQ ID NO.: 66)	miR-6-2	5' - AGGGAACUUCUGCUGUGAUAUA-3' (SEQ ID NO.: 40) 3' - UUUUUUCUUGCUGGUGAC-ACUAU-5' (SEQ ID NO.: 41)
miR-133	5' - AGCUGGUUGACAUCGGGUCAGAU-3' (SEQ ID NO.: 67) 3' - UGUCGACCAACUUUC-CCUGGUU-5' (SEQ ID NO.: 68)	miR-6-3	5' - AGGGAACGGUUGCUGUGAUAUA-3' (SEQ ID NO.: 42) 3' - UUUUUUCUUGCUGGUGAC-ACUAU-5' (SEQ ID NO.: 41)
miR-184	5' - CCUUAUCA-UUCUCUCG-CCCGG-3' (SEQ ID NO.: 69) 3' - CGGGAAUAGUCAAGAG-GCAGGU-5' (SEQ ID NO.: 70)	miR-7	5' - UGGAAGACUAGUGAUUU--UGUUGU-3' (SEQ ID NO.: 43) 3' - CAUUGUUCUGUUCUCCUAAUAAUAAU-5' (SEQ ID NO.: 44)
bantam	5' - CGGUUUUCGAUUUGGUUGACU-3' (SEQ ID NO.: 71) 3' - UUAUCGGAAGUUUUUACUAGAGU-5' (SEQ ID NO.: 72)	miR-8	5' - CAUCGG-ACCGGGCAGCAUUAUA-3' (SEQ ID NO.: 45) 3' - CUGUAGAAUUGACUGUCAUAU-5' (SEQ ID NO.: 46)

Fig. 6A

8/33

*Fig. 6B*

9/33

*Fig. 7*

sod1 sense target: 5'-...agagaggaugaggagacuugggcaaugacugcugacaa...-3' (●)(SEQ ID NO.: 73)
sod1 anti-sense target: 3'-...cucoguacaaccucucugaaacccguuacacugacgacuguuuc...-5' (○)(SEQ ID NO.: 74)

Fig. 8A




Fig. 8D

Fig. 8C

5'-UGGAG (○)
3'-UUAACGUC (●)
ΔG = -10.4

5'-CACAU (●)
3'-UUGUGUA (○)
ΔG = -8.2

Fig. 8B

Fig. 8E

hft sense target: 5' - ...ugcagcugaucaugcugacccugaggaacaguuc...-3' (●)(SEQ ID NO.: 78)
hft anti-sense target: 3' - ...acgucgacuaguagcuacacgacugggacuccuugaag...-5' (○)(SEQ ID NO.: 79)

Fig. 9A




Fig. 9B

Fig. 9C

Fig. 9D





$\Delta G = -10.5$


$\Delta G = -10.5$

Fig. 9E



12/33



$$\Delta G = -8.7 \text{ kcal/mol}$$






$$\Delta G = -7.4 \text{ kcal/mol} \quad \Delta G = -7.2 \text{ kcal/mol}$$



$$\Delta G = -8.4 \text{ kcal/mol}$$

Fig. 10


13/33

*Fig. 11A**Fig. 11B**Fig. 11C*

14/33


5' -GAGACUUGGGCAAUGUGACdTdT-3' (○)(SEQ ID NO.: 13)

 3' -dTdTTCUCAGAACCCGUUACACUG-5' (●)(SEQ ID NO.: 87)

*Fig. 11D*


5' -GAGACUUGGGCAAUGUGACdTdT-3' (○)(SEQ ID NO.: 13)

 3' -dTdTTCUGUGAACCCGUUACACUG-5' (●)(SEQ ID NO.: 88)

*Fig. 11E*

5' -GAGACUUGGGCAAUGUGACdTdT-3' (○)(SEQ ID NO.: 13)


 3' -dTdTTCACUGAACCCGUUACACUG-5' (●)(SEQ ID NO.: 89)

*Fig. 11F*

15/33


5' -GAGACUUGGGCAAUGUGGCdTdT-3' (○)(SEQ ID NO.: 90)

 3' -dTdTTCUCUGAACCCGUUACACUG-5' (●)(SEQ ID NO.: 14)

*Fig. 11G*


5' -GAGACUUGGGCAAUGUGACdTdT-3' (○)(SEQ ID NO.: 13)

 3' -dTdTTCUCUGAACCCGUUACAUG-5' (●)(SEQ ID NO.: 91)


*Fig. 11H*

5' -GAGACUUGGGCAAUGUGGCdTdT-3' (○)(SEQ ID NO.: 90)

 3' -dTdTTCUCUGAACCCGUUACAUG-5' (●)(SEQ ID NO.: 91)

*Fig. 11I*

16/33

*Fig. 12*

17/33




Fig. 13C




Fig. 13B




Fig. 13A

Target + perfect match siRNA

3'-... CCAACUCCAUCAUCCAAACAUAUCACUU...-5' (SEQ ID NO.: 92)
|||||
5'-UGAGGUAGUAGGUUGUAUAAGU-3' (SEQ ID NO.: 93)
3'-UUACUCCAUCUCCAACAUAU-5' (SEQ ID NO.: 94)

1 nt 3' mismatched siRNA

3'-... CCAACUCCAUCAUCCAAACAUAUCACUU...-5' (SEQ ID NO.: 92)
|||||
5'-UGAGGUAGUAGGUUGUAUAAGG-3' (SEQ ID NO.: 95)
UUACUCCAUCUCCAACAUAU (SEQ ID NO.: 94)

2 nt 3' mismatched siRNA

3'-... CCAACUCCAUCAUCCAAACAUAUCACUU...-5' (SEQ ID NO.: 92)
|||||
5'-UGAGGUAGUAGGUUGUAUAAG-3' (SEQ ID NO.: 96)
UUCUCCAUCUCCAACAUAU (SEQ ID NO.: 97)

3 nt 3' mismatched siRNA

3'-... CCAACUCCAUCAUCCAAACAUAUCACUU...-5' (SEQ ID NO.: 92)
|||||
5'-UGAGGUAGUAGGUUGUAUCAG-3' (SEQ ID NO.: 98)
UUCUCCAUCUCCAACAUAAG (SEQ ID NO.: 99)

4 nt 3' mismatched siRNA

3'-... CCAACUCCAUCAUCCAAACAUAUCACUU...-5' (SEQ ID NO.: 92)
|||||
5'-UGAGGUAGUAGGUUGUAGCAG-3' (SEQ ID NO.: 100)
UUCUCCAUCUCCAACAUCG (SEQ ID NO.: 101)

18/33






Fig. 14B

Fig. 14A

*Fig. 14C**Fig. 14D*




Fig. 15A




Fig. 15C




Fig. 15B

21/33




Fig. 16A




Fig. 16B




Fig. 16C






Fig. 16D



23/33




Fig. 18

5'-*UGAGGUA GUAGGUU GUUAUAGU-3' (SEQ ID NO.: 93)
3'-UUACUCCAUCAUCCAACAUAU-5' (SEQ ID NO.: 94)

Fig. 19A




Fig. 19B






Fig. 19C

25/33

*Fig. 20A**Fig. 20B*

26/33

Target and siRNA	Figure #
<p>target : fully matched siRNA</p> <p>3'-...CCAACUCCAUAUCCAACAUAUCACUU...-5' (SEQ ID NO.: 92)</p> <p>5'-UGAGGUAGUAGGUUGUAUAGU-3' (SEQ ID NO.: 93)</p> <p>3'-UUACUCCAUAUCCAACAUAU-5' (SEQ ID NO.: 94)</p>	
<p>target : 1 nt 3' mismatched siRNA</p> <p>3'-...CCAACUCCAUAUCCAACAUAUCACUU...-5' (SEQ ID NO.: 92)</p> <p>5'-UGAGGUAGUAGGUUGUAUAGG-3' (SEQ ID NO.: 95)</p> <p>UUACUCCAUAUCCAACAUAU (SEQ ID NO.: 94)</p>	
<p>target : 2 nt 3' mismatched siRNA</p> <p>3'-...CCAACUCCAUAUCCAACAUAUCACUU...-5' (SEQ ID NO.: 92)</p> <p>5'-UGAGGUAGUAGGUUGUAUAAG-3' (SEQ ID NO.: 96)</p> <p>UUCUCCAUAUCCAACAUAU (SEQ ID NO.: 97)</p>	
<p>target : 3 nt 3' mismatched siRNA</p> <p>3'-...CCAACUCCAUAUCCAACAUAUCACUU...-5' (SEQ ID NO.: 92)</p> <p>5'-UGAGGUAGUAGGUUGUAUCAG-3' (SEQ ID NO.: 98)</p> <p>UUCUCCAUAUCCAACAUAU (SEQ ID NO.: 99)</p>	
<p>target : 4 nt 3' mismatched siRNA</p> <p>3'-...CCAACUCCAUAUCCAACAUAUCACUU...-5' (SEQ ID NO.: 92)</p> <p>5'-UGAGGUAGUAGGUUGUAGCAG-3' (SEQ ID NO.: 100)</p> <p>UUCUCCAUAUCCAACAUCG (SEQ ID NO.: 101)</p>	
<p>target : 5 nt 3' mismatched siRNA</p> <p>3'-...CCAACUCCAUAUCCAACAUAUCACUU...-5' (SEQ ID NO.: 92)</p> <p>5'-UGAGGUAGUAGGUUGUUCAG-3' (SEQ ID NO.: 102)</p> <p>UUCUCCAUAUCCAACAACG (SEQ ID NO.: 103)</p>	

Figures 2b-d,
3a,b and 5a

Fig. 21A

27/33




Figure 3c

Fig. 21B

target : 1 nt 5« mismatched siRNA

3' - . . . CCAACUCCAUAUCAACAUAUCACUU . . . - 5' (SEQ ID NO.: 92)
 |||||
5' - AGAGGUAGUAGGUUGUAUAGU - 3' (SEQ ID NO.: 110)
 UUCGACCAUCAUCCAACAUAU (SEQ ID NO.: 111)

target : 3 nt 5« mismatched siRNA

3' - . . . CCAACUCCAUAUCAACAUAUCACUU . . . - 5' (SEQ ID NO.: 92)
 |||||
5' - ACUGGUAGUAGGUUGUAUAGU - 3' (SEQ ID NO.: 112)
 UUCGACCAUCAUCCAACAUAU (SEQ ID NO.: 113)

target : 5 nt 5« mismatched siRNA

3' - . . . CCAACUCCAUAUCAACAUAUCACUU . . . - 5' (SEQ ID NO.: 92)
 |||||
5' - ACUCCUAGUAGGUUGUAUAGU - 3' (SEQ ID NO.: 114)
 UUCGAGGAUCAUCCAACAUAU (SEQ ID NO.: 115)

target : 6 nt 5« mismatched siRNA

3' - . . . CCAACUCCAUAUCAACAUAUCACUU . . . - 5' (SEQ ID NO.: 92)
 |||||
5' - ACUCCAAGUAGGUUGUAUAGU - 3' (SEQ ID NO.: 116)
 UUCGAGGUUCAUCCAACAUAU (SEQ ID NO.: 117)

target : 7 nt 5« mismatched siRNA

3' - . . . CCAACUCCAUAUCAACAUAUCACUU . . . - 5' (SEQ ID NO.: 92)
 |||||
5' - ACUCCAUGUAGGUUGUAUAGU - 3' (SEQ ID NO.: 118)
 UUCGAGGUACAUCCAACAUAU (SEQ ID NO.: 119)

target : 8 nt 5« mismatched siRNA

3' - . . . CCAACUCCAUAUCAACAUAUCACUU . . . - 5' (SEQ ID NO.: 92)
 |||||
5' - ACUCCAUCUAGGUUGUAUAGU - 3' (SEQ ID NO.: 120)
 UUCGAGGUAGA UCCAACAUAU (SEQ ID NO.: 121)

Figures 4a,
d and 5a

Fig. 21C

29/33

8 nt 3« mismatched target : siRNA

3' - . . . CCAACUCCAUCAUCCAUGUACGUCCUU . . . - 5' (SEQ ID NO.: 122)

|||||

5' - UGAGGUAGUAGGUUGUAUAGU - 3' (SEQ ID NO.: 123)

UUCCUCCAUCAUCCAACAUAU (SEQ ID NO.: 111)

8 nt 5« mismatched target : siRNA

3' - . . . CCAUGAGGUAGAUCCAACAUAUCACUU . . . - 5' (SEQ ID NO.: 124)

|||||

5' - UGAGGUAGUAGGUUGUAUAGU - 3' (SEQ ID NO.: 125)

UUCCUCCAUCAUCCAACAUAU (SEQ ID NO.: 111)

8 nt 5« mismatched target : 8 nt 5« mismatch siRNA

3' - . . . CCAUGAGGUAGAUCCAACAUAUCACUU . . . - 5' (SEQ ID NO.: 124)

|||||

5' - ACUCCAUCUAGGUUGUAUAGU - 3' (SEQ ID NO.: 120)

UUCGAGGUAGAUCCAACAUAU (SEQ ID NO.: 121)

Figure 4b

Fig. 21D

30/33




Figure 4c

Fig. 21E

31/33




Figure 4c

Fig. 21F

32/33




Figure 4c

Fig. 21G

33/33




Figure 5c

Fig. 21H

SEQUENCE LISTING

<110> University of Massachusetts et al.

<120> METHODS AND COMPOSITIONS FOR ENHANCING THE EFFICACY AND SPECIFICITY OF RNAI

<130> UMY-066CPPC

<150> 10/912440

<151> 2004-08-04

<160> 164

<170> PatentIn Ver. 3.3

<210> 1

<211> 40

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 1
cgaggugaac aucacguacg cggaauacuu cgaaaugucc 40

<210> 2

<211> 40

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 2
ggacauuucg aaguauuccg cguacgugau guucaccucg 40

<210> 3

<211> 21

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 3
cguacgcgga auacuucgaa a 21

<210> 4

<211> 21

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 4
ucgaaguauu ccgcguacgu g 21

<210> 5
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 5
uguacgcgga auacuucgaa a 21

<210> 6
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 6
cgaggugaac aucacguacg cggaauacuu cgaaaugucc 40

<210> 7
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 7
ggacauuucg aaguauuccg cguacgugau guucaccucg 40

<210> 8
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 8
gaaguauucc gcguacguga u 21

<210> 9
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 9
 uacguacgcg gaauacuucg a 21

<210> 10
 <211> 21
 <212> RNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 10
 gaaguuuucc gcguacguua u 21

<210> 11
 <211> 42
 <212> RNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 11
 agagaggcau guuggagacu ugggcaaugu gacugcugac aa 42

<210> 12
 <211> 41
 <212> RNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 12
 cuuugucagc agucacauug cccaagucuc caacaugccu c 41

<210> 13
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Combined DNA/RNA Molecule:
 Synthetic oligonucleotide

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 13
 gagacuuggg caaugugact t 21

<210> 14
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 14
gucacauugc ccaagucuct t 21

<210> 15
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 15
gagacuuggg caaugugaat t 21

<210> 16
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 16
uucacauugc ccaagucuct t 21

<210> 17
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 17
gucacauugc ccaagucuat t 21

<210> 18
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 18
uagacuuggg caaugugact t 21

<210> 19
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<220>
<221> modified_base
<222> (1)
<223> Inosine

<400> 19
nagacuuggg caaugugact t 21

<210> 20
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<220>
<221> modified_base
<222> (1)

<223> Inosine

<400> 20
nucacauugc ccaagucuct t 21

<210> 21
<211> 72
<212> RNA
<213> Drosophila sp.

<400> 21
ggcaaauga gguaguaggu uguauaguag uaauuacaca ucuaucuaa caaugugcua 60
gcuuucuuug cu 72

<210> 22
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 22
ugagguagua gguuguauag u 21

<210> 23
<211> 22
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 23
uauacaugu gcuagcuuuc uu 22

<210> 24
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 24
ccaugcuucc uugcaucaa ua 22

<210> 25
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 25
uggaauuaa agaaguaugg ag 22

<210> 26
<211> 21
<212> RNA

<213> Drosophila sp.

<400> 26
ucucaaagug guugugaaau g 21

<210> 27
<211> 23
<212> RNA
<213> Drosophila sp.

<400> 27
uaucaacagcc agcuuugaug agc 23

<210> 28
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 28
cucaucaagu gguugugaua ug 22

<210> 29
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 29
ucuucaaagu ggcagugaca ug 22

<210> 30
<211> 23
<212> RNA
<213> Drosophila sp.

<400> 30
uaucaacagcc agcuuugagg agc 23

<210> 31
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 31
ucuucaaagu gguugugaaa ug 22

<210> 32
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 32
ggaugcaucu ugugcaguua ug 22

<210> 33
<211> 22
<212> RNA

<213> Drosophila sp.

<400> 33
ucacugggca aagugugucu ca 22

<210> 34
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 34
cuuuggucgu ccagccuuag gu 22

<210> 35
<211> 21
<212> RNA
<213> Drosophila sp.

<400> 35
auaaaagcuag acaaccuuug a 21

<210> 36
<211> 23
<212> RNA
<213> Drosophila sp.

<400> 36
aaaggaacga ucguugugau aug 23

<210> 37
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 37
uaucaacagug auuuuuccuuu au 22

<210> 38
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 38
agggaaauagu ugcugugcug ua 22

<210> 39
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 39
uaucaacagug gcuguucuuu uu 22

<210> 40
<211> 23
<212> RNA

<213> Drosophila sp.

<400> 40
agggaacuuc ugcugcugau aua 23

<210> 41
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 41
uaucaacagug gcuguucuuu uu 22

<210> 42
<211> 23
<212> RNA
<213> Drosophila sp.

<400> 42
agggaacggu ugcugaugau gua 23

<210> 43
<211> 23
<212> RNA
<213> Drosophila sp.

<400> 43
uggaagacua gugauuuugu ugu 23

<210> 44
<211> 26
<212> RNA
<213> Drosophila sp.

<400> 44
uaacaauaaa ucccuugucu ucuuac 26

<210> 45
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 45
caucggaccg ggcagcaua ga 22

<210> 46
<211> 23
<212> RNA
<213> Drosophila sp.

<400> 46
uaauacuguc agguaaagau guc 23

<210> 47
<211> 23
<212> RNA

<213> Drosophila sp.

<400> 47

ucuuugguua ucuagcugua uga

23

<210> 48

<211> 23

<212> RNA

<213> Drosophila sp.

<400> 48

auaaagcuag cuuaccgaag uua

23

<210> 49

<211> 21

<212> RNA

<213> Drosophila sp.

<400> 49

accuguaga uccgaauug u

21

<210> 50

<211> 22

<212> RNA

<213> Drosophila sp.

<400> 50

aaauucgguu cuagagaggu uu

22

<210> 51

<211> 21

<212> RNA

<213> Drosophila sp.

<400> 51

aagaacuuuc ucugugaccc g

21

<210> 52

<211> 21

<212> RNA

<213> Drosophila sp.

<400> 52

caucacaguc ugaguucuug c

21

<210> 53

<211> 23

<212> RNA

<213> Drosophila sp.

<400> 53

ugaguauuac aucagguacu ggu

23

<210> 54
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 54
caguacuuau gucauacuac gc 22

<210> 55
<211> 23
<212> RNA
<213> Drosophila sp.

<400> 55
uccucaaagg guugugaaau guc 23

<210> 56
<211> 23
<212> RNA
<213> Drosophila sp.

<400> 56
auaucacagc cauuuugaug agu 23

<210> 57
<211> 25
<212> RNA
<213> Drosophila sp.

<400> 57
ucguuaaaaau guuugugaac uuaug 25

<210> 58
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 58
uaucaacagcc auuuugacga gu 22

<210> 59
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 59
gcgucaaaaau gacugugagc ua 22

<210> 60
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 60
uaucaacagcc auuuugacga gu 22

<210> 61
<211> 21
<212> RNA
<213> Drosophila sp.

<400> 61
ggagcgagac ggggacucac u 21

<210> 62
<211> 21
<212> RNA
<213> Drosophila sp.

<400> 62
ucagucuuuu ucucucuccu a 21

<210> 63
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 63
guauccacug uaggccuaua ug 22

<210> 64
<211> 21
<212> RNA
<213> Drosophila sp.

<400> 64
uaaggcacgc ggugaaugcc a 21

<210> 65
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 65
ucccugagac ccuaacuugu ga 22

<210> 66
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 66
acaaguuuug aucuccggua uu 22

<210> 67
<211> 23
<212> RNA
<213> Drosophila sp.

<400> 67
agcugguuga caucggguca gau 23

<210> 68
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 68
uugguccccu ucaaccagcu gu 22

<210> 69
<211> 21
<212> RNA
<213> Drosophila sp.

<400> 69
ccuuaucuu cucucgcccc g 21

<210> 70
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 70
uggacggaga acugauaagg gc 22

<210> 71
<211> 22
<212> RNA
<213> Drosophila sp.

<400> 71
cgguuuucga uuugguuuga cu 22

<210> 72
<211> 23
<212> RNA
<213> Drosophila sp.

<400> 72
ugagaucauu uugaaagcug auu 23

<210> 73
<211> 42
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 73
agagaggcau guuggagacu ugggcaaugu gacugcugac aa 42

<210> 74
<211> 41
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 74
cuuugucagc agucacauug cccaagucuc caacaugccu c 41

<210> 75

<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 75
uggagacuug ggcaaugugt t 21

<210> 76

<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 76
cacauugccc aagucuccat t 21

<210> 77

<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 77
uacauugccc aagucuccat t 21

<210> 78
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 78
ugcagcugau caucgaugug cugaccuga ggaacaguuc 40

<210> 79
<211> 40
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 79
gaacuguucc ucaggguacag cacaucgaug aucagcugca 40

<210> 80
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 80
ugugcugacc cugaggaaca g 21

<210> 81
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 81
guuccucagg gucagcacau c 21

<210> 82
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 82
ugugcugacc cugaggaaaa g 21

<210> 83
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 83
uuuccucagg gucagcacau c 21

<210> 84
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 84
gagacuuggg caaugagact t 21

<210> 85
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 85
gagacuuggg caaugaaact t 21

<210> 86
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 86
gagacuuggg caauguguct t 21

<210> 87
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 87
gucacauugc ccaagacuct t 21

<210> 88
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 88
gucacauugc ccaaguguct t 21

<210> 89
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 89
gucacauugc ccaagucact t 21

<210> 90
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 90
gagacuuggg caauguggct t 21

<210> 91
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 91
guuacauugc ccaagucuct t 21

<210> 92
<211> 27
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 92
uucacuaauac aaccuacuac cucaacc 27

<210> 93
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 93
ugagguagua gguuguauag u 21

<210> 94
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 94
uauacaaccu acuaccucau u 21

<210> 95
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 95
ugagguagua gguuguauag g 21

<210> 96
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 96
ugagguagua gguuguauaa g 21

<210> 97
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 97
uauacaaccu acuaccuccu u 21

<210> 98
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 98
ugagguagua gguuguauca g 21

<210> 99
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 99
gauacaaccu acuaccuccu u 21

<210> 100
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 100
ugagguagua gguuguagca g 21

<210> 101
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 101
gcuacaaccu acuaccuccu u 21

<210> 102
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 102
ugagguagua gguuguugca g 21

<210> 103
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 103
gcaacaaccu acuaccuccu u 21

<210> 104
<211> 21
<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 104

ugagguagua gguugaugca g

21

<210> 105

<211> 21

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 105

gcaucaaccu acuaccuccu u

21

<210> 106

<211> 21

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 106

ugagguagua gguacaugca g

21

<210> 107

<211> 21

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 107

gcaugaaccu acuaccuccu u

21

<210> 108

<211> 21

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 108

ugagguagua gcaacaugca g

21

<210> 109
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 109
gcauguuccu acuaccuccu u 21

<210> 110
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 110
agagguagua gguuguauag u 21

<210> 111
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 111
uauacaaccu acuaccuccu u 21

<210> 112
<211> 21
<212> RNA

<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 112
acugguagua gguuguauag u 21

<210> 113
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 113
uauacaaccu acuaccagcu u 21

<210> 114
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 114
acuccuagua gguuguauag u 21

<210> 115
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 115
uauacaaccu acuaggagcu u 21

<210> 116
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 116
acuccaagua gguuguauag u 21

<210> 117
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 117
uauacaaccu acuuggagcu u 21

<210> 118
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 118
acuccaugua gguuguauag u 21

<210> 119
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 119
uauacaaccu acauggagcu u 21

<210> 120
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 120
acuccaucua gguuguauag u 21

<210> 121
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 121
uauacaaccu agauggagcu u 21

<210> 122
<211> 27
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 122
uuccugcaug uaccuacuac cucaacc 27

<210> 123
<211> 21
<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 123

ugagguagua gguuguauag u

21

<210> 124

<211> 27

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 124

uucacuaauac aaccuagaug gaguacc

27

<210> 125

<211> 21

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 125

ugagguagua gguuguauag u

21

<210> 126

<211> 27

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 126

uuccugcaug aaccuacuac cucaacc

27

<210> 127

<211> 21

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 127

ugagguagua gguuguauag u

21

<210> 128
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 128
agagguagua gguuguauag u 21

<210> 129
<211> 27
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 129
uuccugcaug aaccuacuac cugaacc 27

<210> 130
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 130
acugguagua gguuguauag u 21

<210> 131
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 131
acucguagua gguuguauag u 21

<210> 132
<211> 27
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 132
uuccugcaug uaccuacuac cucaacc 27

<210> 133
<211> 27
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 133
uuccugcaug uaccuacuac cugaacc 27

<210> 134
<211> 27
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 134
uuccugcaug uuccuacuac cucaacc 27

<210> 135
<211> 27
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 135
uuccugcaug uuccuacuac cugaacc 27

<210> 136
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 136
uauacaaccu acuaccagcu u 21

<210> 137
<211> 31
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 137
ucuucacua u acaaccuacu accucaaccu u 31

<210> 138
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 138
uauacaaccu aguaccucau u 21

<210> 139
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 139
uauacaaccu agauggagcu u 21

<210> 140
<211> 31
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 140
aaucacua u acaaccaaga accucaaccu u 31

<210> 141
<211> 31
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 141
ucuucacua u acauggaacu accucaaccu u 31

<210> 142
<211> 31
<212> RNA
<213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 142
 ucuuccaacu acaaccuacu accucaaccu u 31

<210> 143
 <211> 31
 <212> RNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 143
 ucuuccaacc aagaccuacu accucaaccu u 31

<210> 144
 <211> 40
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 primer

<400> 144
 cgctaatacg actcactata gcagttggcg ccgcgaacga 40

<210> 145
 <211> 40
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 primer

<400> 145
 gcgtaatacg actcactata gtcacatctc atctacctcc 40

<210> 146
 <211> 106
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 primer

<400> 146
 cccatttagg tgacactata gatttacatc gcgttgagt tagaacgggt gtataaaagg 60
 ttgaggtagt aggttgata gtgaagagag gagttcatga tcagtg 106

<210> 147
<211> 106
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
primer

<400> 147
cccatttagg tgacactata gatttacatc gcgttgagt tagaacgggt gtataaaagg 60
ttgaggtagt aggttcatgc aggaagagag gagttcatga tcagtg 106

<210> 148
<211> 106
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220>
<223> Description of Artificial Sequence: Synthetic
primer

<400> 148
cccatttagg tgacactata gatttacatc gcgttgagt tagaacgggt gtataaaagg 60
ttgaggtagt aggtacaugc aggaagagag gagttcatga tcagtg 106

<210> 149
<211> 106
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
primer

<400> 149
cccatttagg tgacactata gatttacatc gcgttgagt tagaacgggt gtataaaagg 60
ttgaggtagt aggaacatgc aggaagagag gagttcatga tcagtg 106

<210> 150
<211> 106
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
primer

<400> 150
cccatttagg tgacactata gatttacatc gcgttgagt tagaacgggt gtataaaagg 60
tactccatct aggttgata gtgaagagag gagttcatga tcagtg 106

<210> 151
<211> 106
<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<400> 151

cccatcttagg tgacactata gatttacatc gcgttgagtg tagaacgggt gtataaaagg 60
tactcgtagt aggttgtata gtgaagagag gagttcatga tcagt 106

<210> 152

<211> 42

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<400> 152

gcgtaatacg actcactata ggaacagtat gtctcagaca tc 42

<210> 153

<211> 21

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<400> 153

uucgaaguau uccgcguacg u 21

<210> 154

<211> 42

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<400> 154

gcgtaatacg actcactata ggacaagcct aattagtgat gc 42

<210> 155

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<400> 155

gaacagtatg tctcagacat c 21

<210> 156
 <211> 41
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 primer

<400> 156
 gcgtaatacgc actcactata gggctttgtt agcagccgga t 41

<210> 157
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 primer

<400> 157
 gggagaccac aacggtttcc c 21

<210> 158
 <211> 31
 <212> RNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 158
 acauuucgaa guauuccgcg uacgugaugu u 31

<210> 159
 <211> 31
 <212> RNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 159
 caucacguac gcggaauacu ucgaaauguc c 31

<210> 160
 <211> 16
 <212> RNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 160
 cguacuuuug uacgug 16

<210> 161
<211> 16
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 161
uguacuuuug uacgug 16

<210> 162
<211> 15
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 162
ucgaauuuuu cgaaa 15

<210> 163
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
probe

<400> 163
acaaauucgg aucuacaggg u 21

<210> 164
<211> 22
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
probe

<400> 164
aaaccucucu agaaccgaau uu 22